

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, C12N 15/10, 15/66, C12Q 1/68, G01N 33/50, C07K 16/18, G01N 33/53, A61K 48/00, 38/17	A2	(11) International Publication Number: WO 99/06551 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/IB98/01235 (22) International Filing Date: 31 July 1998 (31.07.98) (30) Priority Data: 08/905,133 1 August 1997 (01.08.97) US (71) Applicant (for all designated States except US): GENSET [FR/FR]; 24, rue Royale, F-75008 Paris (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): DUMAS MILNE EDWARDS, Jean-Baptiste [FR/FR]; 8, rue Grégoire de Tours, F-75006 Paris (FR). DUCLERT, Aymeric [FR/FR]; 6 ter, rue Victorine, F-94100 Saint-Maur (FR). LACROIX, Bruno [FR/FR]; 93, route de Vourles, F-69230 Saint-Genis Laval (FR). (74) Agents: MARTIN, Jean-Jacques et al.; Cabinet Regimbeau, 26, avenue Kléber, F-75116 Paris (FR).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: 5' ESTs FOR SECRETED PROTEINS IDENTIFIED FROM BRAIN TISSUES (57) Abstract <p>The sequence of 5' ESTs derived from mRNAs encoding secreted proteins are disclosed. The 5' ESTs may be to obtain cDNAs, and genomic DNAs corresponding to the 5' ESTs. The 5' ESTs may also be used in diagnostic, forensic, gene therapy, and chromosome mapping procedures. Upstream regulatory sequences may also be obtained using the 5' ESTs. The 5' ESTs may also be used to design expression vectors and secretion vectors.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5' ESTs FOR SECRETED PROTEINS IDENTIFIED FROM BRAIN TISSUES**Background of the Invention**

The estimated 50,000-100,000 genes scattered along the human chromosomes offer tremendous promise for the understanding, diagnosis, and treatment of human diseases. In addition, probes capable of specifically hybridizing to loci distributed throughout the human genome find applications in the construction of high resolution chromosome maps and in the identification of individuals.

In the past, the characterization of even a single human gene was a painstaking process, requiring years of effort. Recent developments in the areas of cloning vectors, DNA sequencing, and computer technology have merged to greatly accelerate the rate at which human genes can be isolated, sequenced, mapped, and characterized. Cloning vectors such as yeast artificial chromosomes (YACs) and bacterial artificial chromosomes (BACs) are able to accept DNA inserts ranging from 300 to 1000 kilobases (kb) or 100-400 kb in length respectively, thereby facilitating the manipulation and ordering of DNA sequences distributed over great distances on the human chromosomes. Automated DNA sequencing machines permit the rapid sequencing of human genes. Bioinformatics software enables the comparison of nucleic acid and protein sequences, thereby assisting in the characterization of human gene products.

Currently, two different approaches are being pursued for identifying and characterizing the genes distributed along the human genome. In one approach, large fragments of genomic DNA are isolated, cloned, and sequenced. Potential open reading frames in these genomic sequences are identified using bioinformatics software. However, this approach entails sequencing large stretches of human DNA which do not encode proteins in order to find the protein encoding sequences scattered throughout the genome. In addition to requiring extensive sequencing, the bioinformatics software may mischaracterize the genomic sequences obtained. Thus, the software may produce false positives in which non-coding DNA is mischaracterized as coding DNA or false negatives in which coding DNA is mislabeled as non-coding DNA.

An alternative approach takes a more direct route to identifying and characterizing human genes. In this approach, complementary DNAs (cDNAs) are synthesized from isolated messenger RNAs (mRNAs) which encode human proteins. Using this approach,

sequencing is only performed on DNA which is derived from protein coding portions of the genome. Often, only short stretches of the cDNAs are sequenced to obtain sequences called expressed sequence tags (ESTs). The ESTs may then be used to isolate or purify extended cDNAs which include sequences adjacent to the EST sequences. The extended cDNAs may
5 contain all of the sequence of the EST which was used to obtain them or only a portion of the sequence of the EST which was used to obtain them. In addition, the extended cDNAs may contain the full coding sequence of the gene from which the EST was derived or, alternatively, the extended cDNAs may include portions of the coding sequence of the gene from which the EST was derived. It will be appreciated that there may be several extended
10 cDNAs which include the EST sequence as a result of alternate splicing or the activity of alternative promoters.

In the past, these short EST sequences were often obtained from oligo-dT primed cDNA libraries. Accordingly, they mainly corresponded to the 3' untranslated region of the mRNA. In part, the prevalence of EST sequences derived from the 3' end of the mRNA is a
15 result of the fact that typical techniques for obtaining cDNAs are not well suited for isolating cDNA sequences derived from the 5' ends of mRNAs. (Adams *et al.*, *Nature* 377:3-174, 1996; Hillier *et al.*, *Genome Res.* 6:807-828, 1996).

In addition, in those reported instances where longer cDNA sequences have been obtained, the reported sequences typically correspond to coding sequences and do not include
20 the full 5' untranslated region of the mRNA from which the cDNA is derived. Such incomplete sequences may not include the first exon of the mRNA, particularly in situations where the first exon is short. Furthermore, they may not include some exons, often short ones, which are located upstream of splicing sites. Thus, there is a need to obtain sequences derived from the 5' ends of mRNAs.

25 While many sequences derived from human chromosomes have practical applications, approaches based on the identification and characterization of those chromosomal sequences which encode a protein product are particularly relevant to diagnostic and therapeutic uses. Of the 50,000-100,000 protein coding genes, those genes encoding proteins which are secreted from the cell in which they are synthesized, as well as the secreted proteins
30 themselves, are particularly valuable as potential therapeutic agents. Such proteins are often

involved in cell to cell communication and may be responsible for producing a clinically relevant response in their target cells.

In fact, several secretory proteins, including tissue plasminogen activator, G-CSF, GM-CSF, erythropoietin, human growth hormone, insulin, interferon- α , interferon- β , interferon- γ , and interleukin-2, are currently in clinical use. These proteins are used to treat a wide range of conditions, including acute myocardial infarction, acute ischemic stroke, anemia, diabetes, growth hormone deficiency, hepatitis, kidney carcinoma, chemotherapy induced neutropenia and multiple sclerosis. For these reasons, extended cDNAs encoding secreted proteins or portions thereof represent a particularly valuable source of therapeutic agents. Thus, there is a need for the identification and characterization of secreted proteins and the nucleic acids encoding them.

In addition to being therapeutically useful themselves, secretory proteins include short peptides, called signal peptides, at their amino termini which direct their secretion. These signal peptides are encoded by the signal sequences located at the 5' ends of the coding sequences of genes encoding secreted proteins. Because these signal peptides will direct the extracellular secretion of any protein to which they are operably linked, the signal sequences may be exploited to direct the efficient secretion of any protein by operably linking the signal sequences to a gene encoding the protein for which secretion is desired. In addition, portions of signal sequences may also be used to direct the intracellular import of a peptide or protein of interest. This may prove beneficial in gene therapy strategies in which it is desired to deliver a particular gene product to cells other than the cell in which it is produced. Signal sequences encoding signal peptides also find application in simplifying protein purification techniques. In such applications, the extracellular secretion of the desired protein greatly facilitates purification by reducing the number of undesired proteins from which the desired protein must be selected. Thus, there exists a need to identify and characterize the 5' portions of the genes for secretory proteins which encode signal peptides.

Public information on the number of human genes for which the promoters and upstream regulatory regions have been identified and characterized is quite limited. In part, this may be due to the difficulty of isolating such regulatory sequences. Upstream regulatory sequences such as transcription factor binding sites are typically too short to be utilized as probes for isolating promoters from human genomic libraries. Recently, some approaches

have been developed to isolate human promoters. One of them consists of making a CpG island library (Cross, *et al.*, *Nature Genetics* 6: 236-244, 1994). The second consists of isolating human genomic DNA sequences containing SpeI binding sites by the use of SpeI binding protein. (Mortlock *et al.*, *Genome Res.* 6:327-335, 1996). Both of these approaches
5 have their limits due to a lack of specificity or of comprehensiveness.

The present 5' ESTs may be used to efficiently identify and isolate upstream regulatory regions which control the location, developmental stage, rate, and quantity of protein synthesis, as well as the stability of the mRNA. (Theil, *BioFactors* 4:87-93, 1993). Once identified and characterized, these regulatory regions may be utilized in gene therapy or
10 protein purification schemes to obtain the desired amount and locations of protein synthesis or to inhibit, reduce, or prevent the synthesis of undesirable gene products.

In addition, ESTs containing the 5' ends of secretory protein genes may include sequences useful as probes for chromosome mapping and the identification of individuals. Thus, there is a need to identify and characterize the sequences upstream of the 5' coding
15 sequences of genes encoding secretory proteins.

Summary of the Invention

The present invention relates to purified, isolated, or recombinant ESTs which include sequences derived from the authentic 5' ends of their corresponding mRNAs. The term
20 "corresponding mRNA" refers to the mRNA which was the template for the cDNA synthesis which produced the 5' EST. These sequences will be referred to hereinafter as "5' ESTs."
As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition. Individual 5' EST clones isolated from a cDNA library have been conventionally purified to electrophoretic homogeneity. The sequences obtained from these
25 clones could not be obtained directly either from the library or from total human DNA. The cDNA clones are not naturally occurring as such, but rather are obtained via manipulation of a partially purified naturally occurring substance (messenger RNA). The conversion of mRNA into a cDNA library involves the creation of a synthetic substance (cDNA) and pure individual cDNA clones can be isolated from the synthetic library by clonal selection. Thus,
30 creating a cDNA library from messenger RNA and subsequently isolating individual clones from that library results in an approximately 10^4 - 10^6 fold purification of the native message.

Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

As used herein, the term "isolated" requires that the material be removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide present in a living animal is not isolated, but the same polynucleotide, separated from some or all of the coexisting materials in the natural system, is isolated.

As used herein, the term "recombinant" means that the 5' EST is adjacent to "backbone" nucleic acid to which it is not adjacent in its natural environment. Additionally, to be "enriched" the 5' ESTs will represent 5% or more of the number of nucleic acid inserts in a population of nucleic acid backbone molecules. Backbone molecules according to the present invention include nucleic acids such as expression vectors, self-replicating nucleic acids, viruses, integrating nucleic acids, and other vectors or nucleic acids used to maintain or manipulate a nucleic acid insert of interest. Preferably, the enriched 5' ESTs represent 15% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. More preferably, the enriched 5' ESTs represent 50% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. In a highly preferred embodiment, the enriched 5' ESTs represent 90% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules.

"Stringent", moderate, and "low" hybridization conditions are as defined in Example 29.

Unless otherwise indicated, a "complementary" sequence is fully complementary.

Thus, 5' ESTs in cDNA libraries in which one or more 5' ESTs make up 5% or more of the number of nucleic acid inserts in the backbone molecules are "enriched recombinant 5' ESTs" as defined herein. Likewise, 5' ESTs in a population of plasmids in which one or more 5' EST of the present invention have been inserted such that they represent 5% or more of the number of inserts in the plasmid backbone are "enriched recombinant 5' ESTs" as defined herein. However, 5' ESTs in cDNA libraries in which 5' ESTs constitute less than 5% of the number of nucleic acid inserts in the population of backbone molecules, such as libraries in

which backbone molecules having a 5' EST insert are extremely rare, are not "enriched recombinant 5' ESTs."

In particular, the present invention relates to 5' ESTs which are derived from genes encoding secreted proteins. As used herein, a "secreted" protein is one which, when
5 expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal peptides in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g. soluble proteins), or partially (e.g. receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

10 Such 5' ESTs include nucleic acid sequences, called signal sequences, which encode signal peptides which direct the extracellular secretion of the proteins encoded by the genes from which the 5' ESTs are derived. Generally, the signal peptides are located at the amino termini of secreted proteins.

Secreted proteins are translated by ribosomes associated with the "rough"
15 endoplasmic reticulum. Generally, secreted proteins are co-translationally transferred to the membrane of the endoplasmic reticulum. Association of the ribosome with the endoplasmic reticulum during translation of secreted proteins is mediated by the signal peptide. The signal peptide is typically cleaved following its co-translational entry into the endoplasmic reticulum. After delivery to the endoplasmic reticulum, secreted proteins may proceed through the
20 Golgi apparatus. In the Golgi apparatus, the proteins may undergo post-translational modification before entering secretory vesicles which transport them across the cell membrane.

The 5' ESTs of the present invention have several important applications. For example, they may be used to obtain and express cDNA clones which include the full protein
25 coding sequences of the corresponding gene products, including the authentic translation start sites derived from the 5' ends of the coding sequences of the mRNAs from which the 5' ESTs are derived. These cDNAs will be referred to hereinafter as "full length cDNAs." These cDNAs may also include DNA derived from mRNA sequences upstream of the translation start site. The full length cDNA sequences may be used to express the proteins
30 corresponding to the 5' ESTs. As discussed above, secreted proteins are therapeutically important. Thus, the proteins expressed from the cDNAs may be useful in treating or

controlling a variety of human conditions. The 5' ESTs may also be used to obtain the corresponding genomic DNA. The term "corresponding genomic DNA" refers to the genomic DNA which encodes the mRNA from which the 5' EST was derived.

Alternatively, the 5' ESTs may be used to obtain and express extended cDNAs
5 encoding portions of the secreted protein. The portions may comprise the signal peptides of the secreted proteins or the mature proteins generated when the signal peptide is cleaved off. The portions may also comprise polypeptides having at least 10 consecutive amino acids encoded by the extended cDNAs or full length cDNAs. Alternatively, the portions may comprise at least 15 consecutive amino acids encoded by the extended cDNAs or full length
10 cDNAs. In some embodiments, the portions may comprise at least 25 consecutive amino acids encoded by the extended cDNAs or full length cDNAs. In other embodiments, the portions may comprise at least 40 amino acids encoded by the extended cDNAs or full length cDNAs.

Antibodies which specifically recognize the entire secreted proteins encoded by the
15 extended cDNAs, full length cDNAs, or fragments thereof having at least 10 consecutive amino acids, at least 15 consecutive amino acids, at least 25 consecutive amino acids, or at least 40 consecutive amino acids may also be obtained as described below. Antibodies which specifically recognize the mature protein generated when the signal peptide is cleaved may also be obtained as described below. Similarly, antibodies which specifically recognize the
20 signal peptides encoded by the extended cDNAs or full length cDNAs may also be obtained.

In some embodiments, the extended cDNAs obtained using the 5' ESTs include the signal sequence. In other embodiments, the extended cDNAs obtained using the 5' ESTs may include the full coding sequence for the mature protein (*i.e.* the protein generated when the signal polypeptide is cleaved off). In addition, the extended cDNAs obtained using the 5'
25 ESTs may include regulatory regions upstream of the translation start site or downstream of the stop codon which control the amount, location, or developmental stage of gene expression.

As discussed above, secreted proteins are therapeutically important. Thus, the proteins expressed from the extended cDNAs or full length cDNAs obtained using the 5'
30 ESTs may be useful in treating or controlling a variety of human conditions.

The 5' ESTs (or cDNAs or genomic DNAs obtained therefrom) may be used in forensic procedures to identify individuals or in diagnostic procedures to identify individuals having genetic diseases resulting from abnormal expression of the genes corresponding to the 5' ESTs. In addition, the present invention is useful for constructing a high resolution map of the human chromosomes.

The present invention also relates to secretion vectors capable of directing the secretion of a protein of interest. Such vectors may be used in gene therapy strategies in which it is desired to produce a gene product in one cell which is to be delivered to another location in the body. Secretion vectors may also facilitate the purification of desired proteins.

The present invention also relates to expression vectors capable of directing the expression of an inserted gene in a desired spatial or temporal manner or at a desired level. Such vectors may include sequences upstream of the 5' ESTs, such as promoters or upstream regulatory sequences.

Finally, the present invention may also be used for gene therapy to control or treat genetic diseases. Signal peptides may also be fused to heterologous proteins to direct their extracellular secretion.

Bacterial clones containing Bluescript plasmids having inserts containing the 5' ESTs of the present invention (SEQ ID NOs: 38-195 are presently stored at 80°C in 4% (v/v) glycerol in the inventor's laboratories under the designations listed next to the SEQ ID NOs in II). The inserts may be recovered from the deposited materials by growing the appropriate clones on a suitable medium. The Bluescript DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a cesium chloride gradient, size exclusion chromatography, or anion exchange chromatography. The plasmid DNA obtained using these procedures may then be manipulated using standard cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the EST insertion. The PCR product which corresponds to the 5' EST can then be manipulated using standard cloning techniques familiar to those skilled in the art.

One aspect of the present invention is a purified or isolated nucleic acid having the sequence of one of SEQ ID NOs: 38-195 or having a sequence complementary thereto. In one embodiment, the nucleic acid is recombinant.

Another aspect of the present invention is a purified or isolated nucleic acid comprising at least 10 consecutive bases of the sequence of one of SEQ ID NOs: 38-195 or one of the sequences complementary thereto.

Yet another aspect of the present invention is a purified or isolated nucleic acid comprising at least 15 consecutive bases of one of the sequences of SEQ ID NOs: 38-195 or one of the sequences complementary thereto. In one embodiment, the nucleic acid is recombinant.

A further aspect of the present invention is a purified or isolated nucleic acid of at least 15 bases capable of hybridizing under stringent conditions to the sequence of one of SEQ ID NOs: 38-195 or one of the sequences complementary to the sequences of SEQ ID NOs: 38-195. In one embodiment, the nucleic acid is recombinant.

Another aspect of the present invention is a purified or isolated nucleic acid encoding a human gene product, said human gene product having a sequence partially encoded by one of the sequences of SEQ ID NO: 38-195.

Still another aspect of the present invention is a method of making a cDNA encoding a human secretory protein, said human secretory protein being partially encoded by one of SEQ ID NOs 38-195, comprising the steps of contacting a collection of mRNA molecules from human cells with a primer comprising at least 15 consecutive nucleotides of a sequence complementary to one of SEQ ID NOs: 38-195; hybridizing said primer to an mRNA in said collection that encodes said protein; reverse transcribing said hybridized primer to make a first cDNA strand from said mRNA; making a second cDNA strand complementary to said first cDNA strand; and isolating the resulting cDNA encoding said protein comprising said first cDNA strand and said second cDNA strand.

Another aspect of the invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the

cDNA comprises the full protein coding sequence of said protein which sequence is partially included in one of the sequences of SEQ ID NOs: 38-195.

Another aspect of the present invention is a method of making a cDNA encoding a human secretory protein that is partially encoded by one of SEQ ID NOs 38-195, comprising the steps of obtaining a cDNA comprising one of the sequences of SEQ ID NOs: 38-195; contacting said cDNA with a detectable probe comprising at least 15 consecutive nucleotides of said sequence of SEQ ID NO: 38-195 or a sequence complementary thereto under conditions which permit said probe to hybridize to said cDNA; identifying a cDNA which hybridizes to said detectable probe; and isolating said cDNA which hybridizes to said probe.

Another aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

Another aspect of the present invention is a method of making a cDNA comprising one of the sequence of SEQ ID NOs: 38-195, comprising the steps of contacting a collection of mRNA molecules from human cells with a first primer capable of hybridizing to the polyA tail of said mRNA; hybridizing said first primer to said polyA tail; reverse transcribing said mRNA to make a first cDNA strand; making a second cDNA strand complementary to said first cDNA strand using at least one primer comprising at least 15 nucleotides of one of the sequences of SEQ ID NOs 38-195; and isolating the resulting cDNA comprising said first cDNA strand and said second cDNA strand.

Another aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

In one embodiment of the method described in the two paragraphs above, the second cDNA strand is made by contacting said first cDNA strand with a first pair of primers, said

first pair of primers comprising a second primer comprising at least 15 consecutive nucleotides of one of the sequences of SEQ ID NOs 38-195 and a third primer having a sequence therein which is included within the sequence of said first primer; performing a first polymerase chain reaction with said first pair of nested primers to generate a first PCR product; contacting said first PCR product with a second pair of primers, said second pair of primers comprising a fourth primer, said fourth primer comprising at least 15 consecutive nucleotides of said sequence of one of SEQ ID NOs: 38-195, and a fifth primer, said fourth and fifth primers being capable of hybridizing to sequences within said first PCR product; and performing a second polymerase chain reaction, thereby generating a second PCR product.

One aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195, or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

Another aspect of the present invention is the method described four paragraphs above in which the second cDNA strand is made by contacting said first cDNA strand with a second primer comprising at least 15 consecutive nucleotides of the sequences of SEQ ID NOs: 38-195; hybridizing said second primer to said first strand cDNA; and extending said hybridized second primer to generate said second cDNA strand.

Another aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein partially encoded by one of SEQ ID NOs 38-195 or comprising a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

Another aspect of the present invention is a method of making a protein comprising one of the sequences of SEQ ID NOs: 196-353, comprising the steps of obtaining a cDNA encoding the full protein sequence partially included in one of the sequences of sequence of SEQ ID NOs: 38-195; inserting said cDNA in an expression vector such that said cDNA is

operably linked to a promoter; introducing said expression vector into a host cell whereby said host cell produces the protein encoded by said cDNA; and isolating said protein.

Another aspect of the present invention is an isolated protein obtainable by the method described in the preceding paragraph.

5 Another aspect of the present invention is a method of obtaining a promoter DNA comprising the steps of obtaining DNAs located upstream of the nucleic acids of SEQ ID NOs: 38-195 or the sequences complementary thereto; screening said upstream DNAs to identify a promoter capable of directing transcription initiation; and isolating said DNA comprising said identified promoter. In one embodiment, the obtaining step comprises
10 chromosome walking from said nucleic acids of SEQ ID NOs: 38-195 or sequences complementary thereto. In another embodiment, the screening step comprises inserting said upstream sequences into a promoter reporter vector. In another embodiment, the screening step comprises identifying motifs in said upstream DNAs which are transcription factor binding sites or transcription start sites.

15 Another aspect of the present invention is an isolated promoter obtainable by the method described above.

Another aspect of the present invention is an isolated or purified protein comprising one of the sequences of SEQ ID NOs: 196-353.

20 Another aspect of the present invention is the inclusion of at least one of the sequences of SEQ ID NOs: 38-195, or one of the sequences complementary to the sequences of SEQ ID NOs: 38-195, or a fragment thereof of at least 15 consecutive nucleotides in an array of discrete ESTs or fragments thereof of at least 15 nucleotides in length. In one embodiment, the array includes at least two of the sequences of SEQ ID NOs: 38-195, the sequences complementary to the sequences of SEQ ID NOs: 38-195, or fragments thereof
25 at least 15 consecutive nucleotides. In another embodiment, the array includes at least five of the sequences of SEQ ID NOs: 38-195, the sequences complementary to the sequences of SEQ ID NOs: 38-195, or fragments thereof of at least 15 consecutive nucleotides.

Another aspect of the present invention is a promoter having a sequence selected from the group consisting of SEQ ID NOs: 31, 34, and 37.

Brief Description of the Drawings

Figure 1 is a summary of a procedure for obtaining cDNAs which have been selected to include the 5' ends of the mRNAs from which they derived.

Figure 2 shows the distribution of Von Heijne scores for 5' ESTs in each of the categories described herein and the probability that these 5' ESTs encode a signal peptide.

Figure 3 summarizes a general method used to clone and sequence extended cDNAs containing sequences adjacent to 5' ESTs.

Figure 4 (description of promoters structure isolated from SignalTag 5' ESTs) provides a schematic description of promoters isolated and the way they are assembled with the corresponding 5' tags.

Detailed Description of the Preferred Embodiment

Table IV is an analysis of the 43 amino acids located at the N terminus of all human SwissProt proteins to determine the frequency of false positives and false negatives using the techniques for signal peptide identification described herein.

Table V shows the distribution of 5' ESTs in each category described herein and the number of 5' ESTs in each category having a given minimum Von Heijne's score.

Table VI shows the distribution of 5' ESTs in each category described herein with respect to the tissue from which the 5' ESTs of the corresponding mRNA were obtained.

Table VII describes the transcription factor binding sites present in each of these promoters.

I. General Methods for Obtaining 5' ESTs derived from mRNAs with intact 5' ends

In order to obtain the 5' ESTs of the present invention, mRNAs with intact 5' ends must be obtained. Currently, there are two approaches for obtaining such mRNAs with intact 5' ends as described below: either chemical (1) or enzymatic (2).

1. Chemical Methods for Obtaining mRNAs having Intact 5' Ends

One of these approaches is a chemical modification method involving derivatization of the 5' ends of the mRNAs and selection of the derivatized mRNAs. The 5' ends of eukaryotic mRNAs possess a structure referred to as a "cap" which comprises a guanosine

methyated at the 7 position. The cap is joined to the first transcribed base of the mRNA by a 5', 5'-triphosphate bond. In some instances, the 5' guanosine is methylated in both the 2 and 7 positions. Rarely, the 5' guanosine is trimethylated at the 2, 7 and 7 positions. In the chemical method for obtaining mRNAs having intact 5' ends, the 5' cap is specifically
5 derivatized and coupled to a reactive group on an immobilizing substrate. This specific derivatization is based on the fact that only the ribose linked to the methylated guanosine at the 5' end of the mRNA and the ribose linked to the base at the 3' terminus of the mRNA, possess 2', 3'-cis diols.

Optionally, the 2', 3'-cis diol of the 3' terminal ribose may be chemically modified,
10 substituted, converted, or eliminated, leaving only the ribose linked to the methylated guanosine at the 5' end of the mRNA with a 2', 3'-cis diol. A variety of techniques are available for eliminating the 2', 3'-cis diol on the 3' terminal ribose. For example, controlled alkaline hydrolysis may be used to generate mRNA fragments in which the 3' terminal ribose is a 3'-phosphate, 2'-phosphate or (2', 3')-cyclophosphate. Thereafter, the fragment which
15 includes the original 3' ribose may be eliminated from the mixture through chromatography on an oligodT column. Alternatively, a base which lacks the 2', 3'-cis diol may be added to the 3' end of the mRNA using an RNA ligase such as T4 RNA ligase. Example 1 below describes a method for ligation of a nucleoside diphosphate to the 3' end of messenger RNA.

20

EXAMPLE 1

Ligation of the Nucleoside Diphosphate pCp to the 3' End of mRNA.

One µg of RNA was incubated in a final reaction medium of 10 µl in the presence of 5 U of T₄ phage RNA ligase in the buffer provided by the manufacturer (Gibco - BRL), 40 U of the RNase inhibitor RNasin (Promega) and, 2 µl of ³²pCp (Amersham #PB 10208). The
25 incubation was performed at 37°C for 2 hours or overnight at 7-8°C.

Following modification or elimination of the 2', 3'-cis diol at the 3' ribose, the 2', 3'-cis diol present at the 5' end of the mRNA may be oxidized using reagents such as NaBH₄, NaBH₃CN, or sodium periodate, thereby converting the 2', 3'-cis diol to a dialdehyde.

Example 2 describes the oxidation of the 2', 3'-cis diol at the 5' end of the mRNA with sodium periodate.

EXAMPLE 2

5 Oxidation of 2', 3'-cis diol at the 5' End of the mRNA with Sodium Periodate

0.1 OD unit of either a capped oligoribonucleotide of 47 nucleotides (including the cap) or an uncapped oligoribonucleotide of 46 nucleotides were treated as follows. The oligoribonucleotides were produced by *in vitro* transcription using the transcription kit "AmpliScribe T7" (Epicentre Technologies). As indicated below, the DNA template for the
10 RNA transcript contained a single cytosine. To synthesize the uncapped RNA, all four NTPs were included in the *in vitro* transcription reaction. To obtain the capped RNA, GTP was replaced by an analogue of the cap, m7G(5')ppp(5')G. This compound, recognized by the polymerase, was incorporated into the 5' end of the nascent transcript during the initiation of transcription but was not incorporated during the extension step. Consequently, the resulting
15 RNA contained a cap at its 5' end. The sequences of the oligoribonucleotides produced by the *in vitro* transcription reaction were:

+Cap:

5'm7GpppGCAUCCUACUCCCAUCCAAUUCCACCCUAACUCCUCCCAUCUCCAC-
3' (SEQ ID NO:1)

20 -Cap:

5'-pppGCAUCCUACUCCCAUCCAAUUCCACCCUAACUCCUCCCAUCUCCAC-3'
(SEQ ID NO:2)

The oligoribonucleotides were dissolved in 9 µl of acetate buffer (0.1 M sodium acetate, pH 5.2) and 3 µl of freshly prepared 0.1 M sodium periodate solution. The mixture
25 was incubated for 1 hour in the dark at 4°C or room temperature. Thereafter, the reaction was stopped by adding 4 µl of 10% ethylene glycol. The product was ethanol precipitated, resuspended in at least 10 µl of water or appropriate buffer and dialyzed against water.

The resulting aldehyde groups may then be coupled to molecules having a reactive
30 amine group, such as hydrazine, carbazide, thiocarbazide or semicarbazide groups, in order to facilitate enrichment of the 5' ends of the mRNAs. Molecules having reactive amine groups

which are suitable for use in selecting mRNAs having intact 5' ends include avidin, proteins, antibodies, vitamins, ligands capable of specifically binding to receptor molecules, or oligonucleotides. Example 3 below describes the coupling of the resulting dialdehyde to biotin.


5

EXAMPLE 3

Coupling of the Dialdehyde at the 5' End of Transcripts with Biotin

The oxidation product obtained in Example 2 was dissolved in 50 μ l of sodium acetate at a pH between 5 and 5.2 and 50 μ l of freshly prepared 0.02 M solution of biotin hydrazide in a methoxyethanol/water mixture (1:1) of formula:

10



In the compound used in these experiments, $n=5$. However, it will be appreciated that other commercially available hydrazides may also be used, such as molecules of the above formula in which n varies from 0 to 5. The mixture was then incubated for 2 hours at 37°C, precipitated with ethanol and dialyzed against distilled water. Example 4 demonstrates the specificity of the biotinylation reaction.

15

EXAMPLE 4

20

Specificity of Biotinylation of Capped Transcripts

The specificity of the biotinylation for capped mRNAs was evaluated by gel electrophoresis of the following samples:

Sample 1. The 46 nucleotide uncapped *in vitro* transcript prepared as in Example 2 and labeled with 32 pCp as described in Example 1.

25

Sample 2. The 46 nucleotide uncapped *in vitro* transcript prepared as in Example 2, labeled with 32 pCp as described in Example 1, treated with the oxidation reaction of Example 2, and subjected to the biotinylation conditions of Example 3.

Sample 3. The 47 nucleotide capped *in vitro* transcript prepared as in Example 2 and labeled with ^{32}pCp as described in Example 1.

Sample 4. The 47 nucleotide capped *in vitro* transcript prepared as in Example 2, labeled with ^{32}pCp as described in Example 1, treated with the oxidation reaction of Example 2, and subjected to the biotinylation conditions of Example 3.

Samples 1 and 2 had identical migration rates, demonstrating that the uncapped RNAs were not oxidized and biotinylated. Sample 3 migrated more slowly than Samples 1 and 2, while Sample 4 exhibited the slowest migration. The difference in migration of the RNAs in Samples 3 and 4 demonstrates that the capped RNAs were specifically biotinylated.

10

In some cases, mRNAs having intact 5' ends may be enriched by binding the molecule containing a reactive amine group to a suitable solid phase substrate such as the inside of the vessel containing the mRNAs, magnetic beads, chromatography matrices, or nylon or nitrocellulose membranes. For example, where the molecule having a reactive amine group is biotin, the solid phase substrate may be coupled to avidin or streptavidin. Alternatively, where the molecule having the reactive amine group is an antibody or receptor ligand, the solid phase substrate may be coupled to the cognate antigen or receptor. Finally, where the molecule having a reactive amine group comprises an oligonucleotide, the solid phase substrate may comprise a complementary oligonucleotide.

The mRNAs having intact 5' ends may be released from the solid phase following the enrichment procedure. For example, where the dialdehyde is coupled to biotin hydrazide and the solid phase comprises streptavidin, the mRNAs may be released from the solid phase by simply heating to 95 degrees Celsius in 2% SDS. In some methods, the molecule having a reactive amine group may also be cleaved from the mRNAs having intact 5' ends following enrichment. Example 5 describes the capture of biotinylated mRNAs with streptavidin coated beads and the release of the biotinylated mRNAs from the beads following enrichment.

25

EXAMPLE 5

Capture and Release of Biotinylated mRNAs Using Streptavidin Coated Beads

The streptavidin coated magnetic beads were prepared according to the manufacturer's instructions (CPG Inc., USA). The biotinylated mRNAs were added to a

30

hybridization buffer (1.5 M NaCl, pH 5 - 6). After incubating for 30 minutes, the unbound and nonbiotinylated material was removed. The beads were then washed several times in water with 1% SDS. The beads thus obtained were incubated for 15 minutes at 95°C in water containing 2% SDS.

- 5 Example 6 demonstrates the efficiency with which biotinylated mRNAs were recovered from the streptavidin coated beads.

EXAMPLE 6

Efficiency of Recovery of Biotinylated mRNAs

- 10 The efficiency of the recovery procedure was evaluated as follows. Capped RNAs were labeled with ³²pCp, oxidized, biotinylated and bound to streptavidin coated beads as described above. Subsequently, the bound RNAs were incubated for 5, 15 or 30 minutes at 95°C in the presence of 2% SDS.

- 15 The products of the reaction were analyzed by electrophoresis on 12% polyacrylamide gels under denaturing conditions (7 M urea). The gels were subjected to autoradiography. During this manipulation, the hydrazone bonds were not reduced.

Increasing amounts of nucleic acids were recovered as incubation times in 2% SDS increased, demonstrating that biotinylated mRNAs were efficiently recovered.

- 20 In an alternative method for obtaining mRNAs having intact 5' ends, an oligonucleotide which has been derivatized to contain a reactive amine group is specifically coupled to mRNAs having an intact cap. Preferably, the 3' end of the mRNA is blocked prior to the step in which the aldehyde groups are joined to the derivatized oligonucleotide, as described above, so as to prevent the derivatized oligonucleotide from being joined to the 3' end of the mRNA. For example, pCp may be attached to the 3' end of the mRNA using T4 RNA ligase as described in example 1. However, as discussed above, blocking the 3' end of the mRNA is an optional step. Derivatized oligonucleotides may be prepared as described in Example 7.
- 25

EXAMPLE 7**Derivatization of Oligonucleotides**

An oligonucleotide phosphorylated at its 3' end was converted to a 3' hydrazide in 3' by treatment with an aqueous solution of hydrazine or of dihydrazide of the formula
5 $H_2N(R1)NH_2$ at about 1 to 3 M, and at pH 4.5 at a temperature of 8°C overnight. This incubation was performed in the presence of a carbodiimide type agent soluble in water such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide at a final concentration of 0.3 M.

The derivatized oligonucleotide was then separated from the other agents and products using a standard technique for isolating oligonucleotides.

10 As discussed above, the mRNAs to be enriched may be treated to eliminate the 3' OH groups which may be present thereon. This may be accomplished by enzymatic ligation of sequences lacking a 3' OH, such as pCp, as described in Example 1. Alternatively, the 3' OH groups may be eliminated by alkaline hydrolysis as described in Example 8 below.

15

EXAMPLE 8**Elimination of 3' OH Groups of mRNA Using Alkaline Hydrolysis**

In a total volume of 100 µl of 0.1 N sodium hydroxide, 1.5 µg mRNA is incubated for 40 to 60 minutes at 4°C. The solution is neutralized with acetic acid and precipitated with ethanol.

20 Following the optional elimination of the 3' OH groups, the diol groups at the 5' ends of the mRNAs are oxidized as described below in Example 9.

EXAMPLE 9**Oxidation of Diols of mRNA**

25 Up to 1 OD unit of RNA was dissolved in 9 µl of buffer (0.1 M sodium acetate, pH 6-7) or water and 3 µl of freshly prepared 0.1 M sodium periodate solution. The reaction was incubated for 1 h in the dark at 4°C or room temperature. Following the incubation, the reaction was stopped by adding 4 µl of 10% ethylene glycol. Thereafter the mixture was incubated at room temperature for 15 minutes. After ethanol precipitation, the product was
30 resuspended in at least 10 µl of water or appropriate buffer and dialyzed against water.

Following oxidation of the diol groups at the 5' ends of the mRNAs, the derivatized oligonucleotide was joined to the resulting aldehydes as described in Example 10.

EXAMPLE 10

5 Ligature of Aldehydes of mRNA to Derivatized Oligonucleotides

The oxidized mRNA was dissolved in an acidic medium such as 50 μ l of sodium acetate pH 4-6. Fifty μ l of a solution of the derivatized oligonucleotide were added in order to obtain an mRNA:derivatized oligonucleotide ratio of 1:20. The mixture was reduced with a borohydride and incubated for 2 h at 37°C or overnight (14 h) at 10°C. The mixture was
10 then ethanol precipitated, resuspended in 10 μ l or more of water or appropriate buffer and dialyzed against distilled water. If desired, the resulting product may be analyzed using acrylamide gel electrophoresis, HPLC analysis, or other conventional techniques.

Following the attachment of the derivatized oligonucleotide to the mRNAs, a reverse
15 transcription reaction may be performed as described in Example 11 below.

EXAMPLE 11

Reverse Transcription of mRNAs Ligatured to Derivatized Oligonucleotides

An oligodeoxyribonucleotide was derivatized as follows. Three OD units of an
20 oligodeoxyribonucleotide of sequence 5' ATCAAGAATTTCGCACGAGACCATTAA3' (SEQ ID NO:3) having 5'-OH and 3'-P ends were dissolved in 70 μ l of a 1.5 M hydroxybenzotriazole solution, pH 5.3, prepared in dimethylformamide/water (75:25) containing 2 μ g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide. The mixture was incubated for 2 h 30 min at 22°C and then precipitated twice in LiClO₄/acetone. The pellet
25 was resuspended in 200 μ l of 0.25 M hydrazine and incubated at 8°C from 3 to 14 h. Following the hydrazine reaction, the mixture was precipitated twice in LiClO₄/acetone.

The messenger RNAs to be reverse transcribed were extracted from blocks of placenta having sides of 2 cm which had been stored at -80°C. The total RNA was extracted using conventional acidic phenol techniques. Oligo-dT chromatography was used to purify
30 the mRNAs. The integrity of the mRNAs was checked by Northern-blotting.

The diol groups on 7 μ g of the placental mRNAs were oxidized as described above in Example 9. The derivatized oligonucleotide was joined to the mRNAs as described in Example 10 above except that the precipitation step was replaced by an exclusion chromatography step to remove derivatized oligodeoxyribonucleotides which were not joined to mRNAs. Exclusion chromatography was performed as follows:

Ten ml of Ultrogel AcA34 (BioSeptra#230151) gel, a mix of agarose and acrylamide, were equilibrated in 50 ml of a solution of 10 mM Tris pH 8.0, 300 mM NaCl, 1 mM EDTA, and 0.05% SDS. The mixture was allowed to sediment. The supernatant was eliminated and the gel was resuspended in 50 ml of buffer. This procedure was repeated 2 or 3 times.

A glass bead (diameter 3 mm) was introduced into a 2 ml disposable pipette (length 25 cm). The pipette was filled with the gel suspension until the height of the gel stabilized at 1 cm from the top of the pipette. The column was then equilibrated with 20 ml of equilibration buffer (10 mM Tris HCl pH 7.4, 20 mM NaCl).

Ten μ l of the mRNA which had reacted with the derivatized oligonucleotide were mixed in 39 μ l of 10 mM urea and 2 μ l of blue-glycerol buffer, which had been prepared by dissolving 5 mg of bromophenol blue in 60% glycerol (v/v), and passing the mixture through a 0.45 μ m diameter filter.

The column was then loaded with the mRNAs coupled to the oligonucleotide. As soon as the sample had penetrated, equilibration buffer was added. Hundred μ l fractions were then collected. Derivatized oligonucleotide which had not been attached to mRNA appeared in fraction 16 and later fractions. Thus, fractions 3 to 15 were combined and precipitated with ethanol.

To determine whether the derivatized oligonucleotide was actually linked to mRNA, one tenth of the combined fractions were spotted twice on a nylon membrane and hybridized to a radioactive probe using conventional techniques. The 32 P labeled probe used in these hybridizations was an oligodeoxyribonucleotide of sequence 5'TAATGGTCTCGTGCGAATTCTTGAT3' (SEQ ID NO:4) anticomplementary to the derivatized oligonucleotide. A signal observed after autoradiography, indicated that the derivatized oligonucleotide had been truly joined to the mRNA.

The remaining nine tenth of the mRNAs which had reacted with the derivatized oligonucleotide was reverse transcribed as follows. A reverse transcription reaction was

carried out with reverse transcriptase following the manufacturer's instructions and 50 pmol of nonamers with random sequence as primers.

To ensure that reverse transcription had been carried out through the cap structure, two types of experiments were performed.

5 In the first approach, after elimination of RNA of the cDNA:RNA heteroduplexes obtained from the reverse transcription reaction by an alkaline hydrolysis, a portion of the resulting single stranded cDNAs was spotted on a positively charged membrane and hybridized, using conventional methods, to a ³²P labeled probe having a sequence identical to that of the derivatized oligonucleotide. Control spots containing, 1 pmol, 100 fmol, 50 fmol,
10 10 fmol and 1 fmol of a control oligodeoxyribonucleotide of sequence identical to that of the derivatized oligonucleotide were included. The signal observed in the spots containing the cDNA indicated that approximately 15 fmol of the derivatized oligonucleotide had been reverse transcribed. These results demonstrate that the reverse transcription can be performed through the cap and, in particular, that reverse transcriptase crosses the 5'-P-P-P-
15 5' bond of the cap of eukaryotic messenger RNAs.

In the second type of experiment, the single stranded cDNAs obtained from the above first strand synthesis were used as template for PCR reactions. Two types of reactions were carried out. First, specific amplification of the mRNAs for alpha globin, dehydrogenase, pp15 and elongation factor E4 were carried out using the following pairs of
20 oligodeoxyribonucleotide primers.

alpha-globin

GLO-S: 5'CCG ACA AGA CCA ACG TCA AGG CCG C3' (SEQ ID NO:5)

GLO-As: 5'TCA CCA GCA GGC AGT GGC TTA GGA G 3' (SEQ ID NO:6)

25

dehydrogenase

3 DH-S: 5'AGT GAT TCC TGC TAC TTT GGA TGG C3' (SEQ ID NO:7)

3 DH-As: 5'GCT TGG TCT TGT TCT GGA GTT TAG A3' (SEQ ID NO:8)

30

pp15

PP15-S: 5'TCC AGA ATG GGA GAC AAG CCA ATT T3' (SEQ ID NO:9)

PP15-As: 5'AGG GAG GAG GAA ACA GCG TGA GTC C3' (SEQ ID NO:10)

Elongation factor E4

EFA1-S: 5'ATG GGA AAG GAA AAG ACT CAT ATC A3' (SEQ ID NO:11)

5 EF1A-As: 5'AGC AGC AAC AAT CAG GAC AGC ACA G3' (SEQ ID NO:12)

Second, non specific amplifications were also carried out with the antisense oligodeoxyribonucleotides of the pairs described above and with a primer derived from the sequence of the derivatized oligodeoxyribonucleotide
10 (5'ATCAAGAATTCGCACGAGACCATT3') (SEQ ID NO:13).

One twentieth of the following RT-PCR product samples were run on a 1.5% agarose gel and stained with ethidium bromide.

Sample 1: The products of a PCR reaction using the globin primers of SEQ ID NOs 5 and 6 in the presence of cDNA.

15 Sample 2: The products of a PCR reaction using the globin primers of SEQ ID NOs 5 and 6 in the absence of added cDNA.

Sample 3: The products of a PCR reaction using the dehydrogenase primers of SEQ ID NOs 7 and 8 in the presence of cDNA.

20 Sample 4: The products of a PCR reaction using the dehydrogenase primers of SEQ ID NOs 7 and 8 in the absence of added cDNA.

Sample 5: The products of a PCR reaction using the pp15 primers of SEQ ID NOs 9 and 10 in the presence of cDNA.

Sample 6: The products of a PCR reaction using the pp15 primers of SEQ ID NOs 9 and 10 in the absence of added cDNA.

25 Sample 7: The products of a PCR reaction using the EIF4 primers of SEQ ID NOs 11 and 12 in the presence of added cDNA.

Sample 8: The products of a PCR reaction using the EIF4 primers of SEQ ID NOs 11 and 12 in the absence of added cDNA.

30 A band of the size expected for the PCR product was observed only in samples 1, 3, 5 and 7, thus indicating the presence of the corresponding sequence in the cDNA population.

PCR reactions were also carried out with the antisense oligonucleotides of the globin and dehydrogenase primers (SEQ ID NOs 6 and 8) and an oligonucleotide whose sequence corresponds to that of the derivatized oligonucleotide. The presence of PCR products of the expected size in the samples equivalent to above samples 1 and 3 indicated that the
5 derivatized oligonucleotide had been linked to mRNA.

The above examples summarize the chemical procedure for enriching mRNAs for those having intact 5' ends as illustrated in Figure 1. Further detail regarding the chemical approaches for obtaining such mRNAs are disclosed in International Application No.
10 WO96/34981, published November 7, 1996, which is incorporated herein by reference. Strategies based on the above chemical modifications to the 5' cap structure may be utilized to generate cDNAs selected to include the 5' ends of the mRNAs from which they derived. In one version of such procedures, the 5' ends of the mRNAs are modified as described above. Thereafter, a reverse transcription reaction is conducted to extend a primer
15 complementary to the 5' end of the mRNA. Single stranded RNAs are eliminated to obtain a population of cDNA/mRNA heteroduplexes in which the mRNA includes an intact 5' end. The resulting heteroduplexes may be captured on a solid phase coated with a molecule capable of interacting with the molecule used to derivatize the 5' end of the mRNA. Thereafter, the strands of the heteroduplexes are separated to recover single stranded first
20 cDNA strands which include the 5' end of the mRNA. Second strand cDNA synthesis may then proceed using conventional techniques. For example, the procedures disclosed in WO 96/34981 or in Carninci. *et al.*, *Genomics* 37:327-336, 1996, the disclosures of which are incorporated herein by reference, may be employed to select cDNAs which include the sequence derived from the 5' end of the coding sequence of the mRNA.

25 Following ligation of the oligonucleotide tag to the 5' cap of the mRNA, a reverse transcription reaction is conducted to extend a primer complementary to the mRNA to the 5' end of the mRNA. Following elimination of the RNA component of the resulting heteroduplex using standard techniques, second strand cDNA synthesis is conducted with a primer complementary to the oligonucleotide tag.
30

2. Enzymatic Methods for Obtaining mRNAs having Intact 5' Ends

Other techniques for selecting cDNAs extending to the 5' end of the mRNA from which they are derived are fully enzymatic. Some versions of these techniques are disclosed in Dumas Milne Edwards J.B. (Doctoral Thesis of Paris VI University, Le clonage des ADNc complets: difficultes et perspectives nouvelles. Apports pour l'etude de la regulation de l'expression de la tryptophane hydroxylase de rat, 20 Dec. 1993), EP0 625572 and Kato *et al.*, *Gene* 150:243-250, 1994, the disclosures of which are incorporated herein by reference.

Briefly, in such approaches, isolated mRNA is treated with alkaline phosphatase to remove the phosphate groups present on the 5' ends of uncapped incomplete mRNAs. Following this procedure, the cap present on full length mRNAs is enzymatically removed with a decapping enzyme such as T4 polynucleotide kinase or tobacco acid pyrophosphatase. An oligonucleotide, which may be either a DNA oligonucleotide or a DNA-RNA hybrid oligonucleotide having RNA at its 3' end, is then ligated to the phosphate present at the 5' end of the decapped mRNA using T4 RNA ligase. The oligonucleotide may include a restriction site to facilitate cloning of the cDNAs following their synthesis. Example 12 below describes one enzymatic method based on the doctoral thesis of Dumas.

EXAMPLE 12

Enzymatic Approach for Obtaining 5' ESTs

Twenty micrograms of PolyA+ RNA were dephosphorylated using Calf Intestinal Phosphatase (Biolabs). After a phenol chloroform extraction, the cap structure of mRNA was hydrolysed using the Tobacco Acid Pyrophosphatase (purified as described by Shinshi *et al.*, *Biochemistry* 15: 2185-2190, 1976) and a hemi 5'DNA/RNA-3' oligonucleotide having an unphosphorylated 5' end, a stretch of adenosine ribophosphate at the 3' end, and an EcoRI site near the 5' end was ligated to the 5'P ends of mRNA using the T4 RNA ligase (Biolabs). Oligonucleotides suitable for use in this procedure are preferably 30 to 50 bases in length. Oligonucleotides having an unphosphorylated 5' end may be synthesized by adding a fluorochrome at the 5' end. The inclusion of a stretch of adenosine ribophosphates at the 3' end of the oligonucleotide increases ligation efficiency. It will be appreciated that the oligonucleotide may contain cloning sites other than EcoRI.

Following ligation of the oligonucleotide to the phosphate present at the 5' end of the decapped mRNA, first and second strand cDNA synthesis is carried out using conventional methods or those specified in EP0 625,572 and Kato *et al. supra*, and Dumas Milne Edwards, *supra*, the disclosures of which are incorporated herein by reference. The resulting cDNA may then be ligated into vectors such as those disclosed in Kato *et al. supra* or other nucleic acid vectors known to those skilled in the art using techniques such as those described in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2d Ed., Cold Spring Harbor Laboratory Press, 1989, the disclosure of which is incorporated herein by reference.

II. Obtention and Characterization of the 5' ESTs of the Present Invention

The 5' ESTs of the present invention were obtained using the aforementioned chemical and enzymatic approaches for enriching mRNAs for those having intact 5' ends as described below.

1. Obtention of 5' ESTS Using mRNAs with Intact 5' Ends

First, mRNAs were prepared as described in Example 13 below.

EXAMPLE 13

Preparation of mRNA With Intact 5' Ends

Total human RNAs or polyA⁺ RNAs derived from 29 different tissues were respectively purchased from LABIMO and CLONTECH and used to generate 44 cDNA libraries as follows. The purchased RNA had been isolated from cells or tissues using acid guanidium thiocyanate-phenol-chloroform extraction (Chomczynski and Sacchi, *Analytical Biochemistry* 162:156-159, 1987). PolyA⁺ RNA was isolated from total RNA (LABIMO) by two passes of oligo dT chromatography, as described by Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69:1408-1412, 1972 in order to eliminate ribosomal RNA.

The quality and the integrity of the polyA⁺ RNAs were checked. Northern blots hybridized with a globin probe were used to confirm that the mRNAs were not degraded. Contamination of the polyA⁺ mRNAs by ribosomal sequences was checked using Northern blots and a probe derived from the sequence of the 28S rRNA. Preparations of mRNAs with

less than 5% of rRNAs were used in library construction. To avoid constructing libraries with RNAs contaminated by exogenous sequences (prokaryotic or fungal), the presence of bacterial 16S ribosomal sequences or of two highly expressed fungal mRNAs was examined using PCR.

5 Following preparation of the mRNAs, the above described chemical and/or the enzymatic procedures for enriching mRNAs for those having intact 5' ends were employed to obtain 5' ESTs from various tissues. In both approaches, an oligonucleotide tag was attached to the 5' ends of the mRNAs. The oligonucleotide tag had an EcoRI site therein to facilitate later cloning procedures. To facilitate the processing of single stranded and double
10 stranded cDNA obtained in the construction of the libraries, the same nucleotidic sequence was used to design the ligated oligonucleotide in both chemical and enzymatic approaches. Nevertheless, in the chemical procedure, the tag used was an oligodeoxyribonucleotide which was linked to the cap of the mRNA whereas in the enzymatic ligation, the tag was a chimeric hemi 5'DNA/RNA3' oligonucleotide which was ligated to the 5' end of decapped mRNA as
15 described in example 12.

 Following attachment of the oligonucleotide tag to the mRNA by either the chemical or enzymatic methods, the integrity of the mRNA was examined by performing a Northern blot with 200 to 500 ng of mRNA using a probe complementary to the oligonucleotide tag before performing the first strand synthesis as described in example 14.

20

EXAMPLE 14

cDNA Synthesis Using mRNA Templates Having Intact 5' Ends

 For the mRNAs joined to oligonucleotide tags using both the chemical and enzymatic methods, first strand cDNA synthesis was performed using the Superscript II (Gibco BRL) or
25 the Rnase H Minus M-MLV (Promega) reverse transcriptase with random nonamers as primers. In order to protect internal EcoRI sites in the cDNA from digestion at later steps in the procedure, methylated dCTP was used for first strand synthesis. After removal of RNA by an alkaline hydrolysis, the first strand of cDNA was precipitated using isopropanol in order to eliminate residual primers.

30 For both the chemical and the enzymatic methods, the second strand of the cDNA was synthesized with a Klenow fragment using a primer corresponding to the 5' end of the

ligated oligonucleotide described in Example 12. Preferably, the primer is 20-25 bases in length. Methylated dCTP was also used for second strand synthesis in order to protect internal EcoRI sites in the cDNA from digestion during the cloning process.

Following cDNA synthesis, the cDNAs were cloned into pBlueScript as described in
5 Example 15 below.

EXAMPLE 15

Cloning of cDNAs derived from mRNA with intact 5' ends into BlueScript

Following second strand synthesis, the ends of the cDNA were blunted with T4 DNA
10 polymerase (Biolabs) and the cDNA was digested with EcoRI. Since methylated dCTP was used during cDNA synthesis, the EcoRI site present in the tag was the only hemi-methylated site, hence the only site susceptible to EcoRI digestion. The cDNA was then size fractionated using exclusion chromatography (AcA, Biosepra) and fractions corresponding to cDNAs of more than 150 bp were pooled and ethanol precipitated. The cDNA was directionally cloned
15 into the SmaI and EcoRI ends of the phagemid pBlueScript vector (Stratagene). The ligation mixture was electroporated into bacteria and propagated under appropriate antibiotic selection.

Clones containing the oligonucleotide tag attached were then selected as described in
Example 16 below.

20

EXAMPLE 16

Selection of Clones Having the Oligonucleotide Tag Attached Thereto

The plasmid DNAs containing 5' EST libraries made as described above were purified (Qiagen). A positive selection of the tagged clones was performed as follows.
25 Briefly, in this selection procedure, the plasmid DNA was converted to single stranded DNA using gene II endonuclease of the phage F1 in combination with an exonuclease (Chang *et al.*, *Gene* 127:95-8, 1993) such as exonuclease III or T7 gene 6 exonuclease. The resulting single stranded DNA was then purified using paramagnetic beads as described by Fry *et al.*, *Biotechniques*, 13: 124-131, 1992. In this procedure, the single stranded DNA was
30 hybridized with a biotinylated oligonucleotide having a sequence corresponding to the 3' end of the oligonucleotide described in Example 13. Preferably, the primer has a length of 20-25

bases. Clones including a sequence complementary to the biotinylated oligonucleotide were captured by incubation with streptavidin coated magnetic beads followed by magnetic selection. After capture of the positive clones, the plasmid DNA was released from the magnetic beads and converted into double stranded DNA using a DNA polymerase such as the ThermoSequenase obtained from Amersham Pharmacia Biotech. Alternatively, protocols such as the one described in the Gene Trapper kit available from Gibco BRL may be used. The double stranded DNA was then electroporated into bacteria. The percentage of positive clones having the 5' tag oligonucleotide was estimated to typically rank between 90 and 98% using dot blot analysis.

Following electroporation, the libraries were ordered in 384-microtiter plates (MTP). A copy of the MTP was stored for future needs. Then the libraries were transferred into 96 MTP and sequenced as described below.

EXAMPLE 17

Sequencing of Inserts in Selected Clones

Plasmid inserts were first amplified by PCR on PE 9600 thermocyclers (Perkin-Elmer, Applied Biosystems Division, Foster City, CA), using standard SETA-A and SETA-B primers (Genset SA), AmpliTaqGold (Perkin-Elmer), dNTPs (Boehringer), buffer and cycling conditions as recommended by the Perkin-Elmer Corporation.

PCR products were then sequenced using automatic ABI Prism 377 sequencers (Perkin Elmer). Sequencing reactions were performed using PE 9600 thermocyclers with standard dye-primer chemistry and ThermoSequenase (Amersham Pharmacia Biotech). The primers used were either T7 or 21M13 (available from Genset SA) as appropriate. The primers were labeled with the JOE, FAM, ROX and TAMRA dyes. The dNTPs and ddNTPs used in the sequencing reactions were purchased from Boehringer. Sequencing buffer, reagent concentrations and cycling conditions were as recommended by Amersham.

Following the sequencing reaction, the samples were precipitated with ethanol, resuspended in formamide loading buffer, and loaded on a standard 4% acrylamide gel. Electrophoresis was performed for 2.5 hours at 3000V on an ABI 377 sequencer, and the sequence data were collected and analyzed using the ABI Prism DNA Sequencing Analysis Software, version 2.1.2.

2. Computer analysis of the Obtained 5' ESTs: Construction of NetGene and SignalTag databases

The sequence data from the 44 cDNA libraries made as described above were transferred to a proprietary database, where quality control and validation steps were performed. A proprietary base-caller, working using a Unix system, automatically flagged suspect peaks, taking into account the shape of the peaks, the inter-peak resolution, and the noise level. The proprietary base-caller also performed an automatic trimming. Any stretch of 25 or fewer bases having more than 4 suspect peaks was considered unreliable and was discarded. Sequences corresponding to cloning vector or ligation oligonucleotides were automatically removed from the EST sequences. However, the resulting EST sequences may contain 1 to 5 bases belonging to the above mentioned sequences at their 5' end. If needed, these can easily be removed on a case to case basis.

Following sequencing as described above, the sequences of the 5' ESTs were entered in NetGene™, a proprietary database called for storage and manipulation as described below. It will be appreciated by those skilled in the art that the data could be stored and manipulated on any medium which can be read and accessed by a computer. Computer readable media include magnetically, optically, or electronically readable media. For example, the computer readable media may be a hard disc, a floppy disc, a magnetic tape, CD-ROM, RAM, or ROM as well as other types of other media known to those skilled in the art.

In addition, the sequence data may be stored and manipulated in a variety of data processor programs in a diversity of formats. For instance, the sequence data may be stored as text in a word processing file, such as Microsoft WORD or WORDPERFECT or as an ASCII file in a variety of database programs familiar to those of skill in the art, such as DB2, SYBASE, or ORACLE.

The computer readable media on which the sequence information is stored may be in a personal computer, a network, a server or other computer systems known to those skilled in the art. The computer or other system preferably includes the storage media described above, and a processor for accessing and manipulating the sequence data. Once the sequence data has been stored, it may be manipulated and searched to locate those stored sequences which contain a desired nucleic acid sequence or which encode a protein having a particular functional domain. For example, the stored sequence information may be compared to other

known sequences to identify homologies, motifs implicated in biological function, or structural motifs.

Programs which may be used to search or compare the stored sequences include the MacPattern (EMBL), BLAST, and BLAST2 program series (NCBI), basic local alignment
5 search tool programs for nucleotide (BLASTN) and peptide (BLASTX) comparisons (Altschul *et al*, *J. Mol. Biol.* 215: 403, 1990) and FASTA (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85: 2444, 1988). The BLAST programs then extend the alignments on the basis of defined match and mismatch criteria.

Motifs which may be detected using the above programs and those described in
10 Example 28 include sequences encoding leucine zippers, helix-turn-helix motifs, glycosylation sites, ubiquitination sites, alpha helices, and beta sheets, signal sequences encoding signal peptides which direct the secretion of the encoded proteins, sequences implicated in transcription regulation such as homeoboxes, acidic stretches, enzymatic active sites, substrate binding sites, and enzymatic cleavage sites.

15 Before searching the cDNAs in the NetGene™ database for sequence motifs of interest, cDNAs derived from mRNAs which were not of interest were identified and eliminated from further consideration as described in Example 18 below.

EXAMPLE 18

20 Elimination of Undesired Sequences from Further Consideration

5' ESTs in the NetGene™ database which were derived from undesired sequences such as transfer RNAs, ribosomal RNAs, mitochondrial RNAs, prokaryotic RNAs, fungal RNAs, Alu sequences, L1 sequences, or repeat sequences were identified using the FASTA and BLASTN programs with the parameters listed in Table I.

25 To eliminate 5' ESTs encoding tRNAs from further consideration, the 5' EST sequences were compared to the sequences of 1190 known tRNAs obtained from EMBL release 38, of which 100 were human. The comparison was performed using FASTA on both strands of the 5' ESTs. Sequences having more than 80% homology over more than 60 nucleotides were identified as tRNA. Of the 144,341 sequences screened, 26 were identified
30 as tRNAs and eliminated from further consideration.

To eliminate 5' ESTs encoding rRNAs from further consideration, the 5' EST sequences were compared to the sequences of 2497 known rRNAs obtained from EMBL release 38, of which 73 were human. The comparison was performed using BLASTN on both strands of the 5' ESTs with the parameter S=108. Sequences having more than 80%
5 homology over stretches longer than 40 nucleotides were identified as rRNAs. Of the 144,341 sequences screened, 3,312 were identified as rRNAs and eliminated from further consideration.

To eliminate 5' ESTs encoding mtRNAs from further consideration, the 5' EST sequences were compared to the sequences of the two known mitochondrial genomes for
10 which the entire genomic sequences are available and all sequences transcribed from these mitochondrial genomes including tRNAs, rRNAs, and mRNAs for a total of 38 sequences. The comparison was performed using BLASTN on both strands of the 5' ESTs with the parameter S=108. Sequences having more than 80% homology over stretches longer than 40 nucleotides were identified as mtRNAs. Of the 144,341 sequences screened, 6,110 were
15 identified as mtRNAs and eliminated from further consideration.

Sequences which might have resulted from exogenous contaminants were eliminated from further consideration by comparing the 5' EST sequences to release 46 of the EMBL bacterial and fungal divisions using BLASTN with the parameter S=144. All sequences having more than 90% homology over at least 40 nucleotides were identified as exogenous
20 contaminants. Of the 42 cDNA libraries examined, the average percentages of prokaryotic and fungal sequences contained therein were 0.2% and 0.5% respectively. Among these sequences, only one could be identified as a sequence specific to fungi. The others were either fungal or prokaryotic sequences having homologies with vertebrate sequences or including repeat sequences which had not been masked during the electronic comparison.

25 In addition, the 5' ESTs were compared to 6093 Alu sequences and 1115 L1 sequences to mask 5' ESTs containing such repeat sequences. 5' ESTs including THE and MER repeats, SSTR sequences or satellite, micro-satellite, or telomeric repeats were also eliminated from further consideration. On average, 11.5% of the sequences in the libraries contained repeat sequences. Of this 11.5%, 7% contained Alu repeats, 3.3% contained L1
30 repeats and the remaining 1.2% were derived from the other screened types of repetitive sequences. These percentages are consistent with those found in cDNA libraries prepared by

other groups. For example, the cDNA libraries of Adams *et al.* contained between 0% and 7.4% Alu repeats depending on the source of the RNA which was used to prepare the cDNA library (Adams *et al.*, *Nature* 377:174, 1996).

- 5 The sequences of those 5' ESTs remaining after the elimination of undesirable sequences were compared with the sequences of known human mRNAs to determine the accuracy of the sequencing procedures described above.

EXAMPLE 19

10 Measurement of Sequencing Accuracy by Comparison to Known Sequences

To further determine the accuracy of the sequencing procedure described above, the sequences of 5' ESTs derived from known sequences were identified and compared to the original known sequences. First, a FASTA analysis with overhangs shorter than 5 bp on both ends was conducted on the 5' ESTs to identify those matching an entry in the public human mRNA database. The 6655 5' ESTs which matched a known human mRNA were then
15 realigned with their cognate mRNA and dynamic programming was used to include substitutions, insertions, and deletions in the list of "errors" which would be recognized. Errors occurring in the last 10 bases of the 5' EST sequences were ignored to avoid the inclusion of spurious cloning sites in the analysis of sequencing accuracy.

- 20 This analysis revealed that the sequences incorporated in the NetGene™ database had an accuracy of more than 99.5%.

To determine the efficiency with which the above selection procedures select cDNAs which include the 5' ends of their corresponding mRNAs, the following analysis was
25 performed.

EXAMPLE 20

Determination of Efficiency of 5' EST Selection

- To determine the efficiency at which the above selection procedures isolated 5' ESTs
30 which included sequences close to the 5' end of the mRNAs from which they derived, the sequences of the ends of the 5' ESTs derived from the elongation factor 1 subunit α and

ferritin heavy chain genes were compared to the known cDNA sequences of these genes. Since the transcription start sites of both genes are well characterized, they may be used to determine the percentage of derived 5' ESTs which included the authentic transcription start sites.

5 For both genes, more than 95% of the obtained 5' ESTs actually included sequences close to or upstream of the 5' end of the corresponding mRNAs.

To extend the analysis of the reliability of the procedures for isolating 5' ESTs from ESTs in the NetGene™ database, a similar analysis was conducted using a database composed of human mRNA sequences extracted from GenBank database release 97 for
10 comparison. The 5' ends of more than 85% of 5' ESTs derived from mRNAs included in the GeneBank database were located close to the 5' ends of the known sequence. As some of the mRNA sequences available in the GenBank database are deduced from genomic sequences, a 5' end matching with these sequences will be counted as an internal match. Thus, the method used here underestimates the yield of ESTs including the authentic 5' ends
15 of their corresponding mRNAs.

The EST libraries made above included multiple 5' ESTs derived from the same mRNA. The sequences of such 5' ESTs were compared to one another and the longest 5' ESTs for each mRNA were identified. Overlapping cDNAs were assembled into continuous
20 sequences (contigs). The resulting continuous sequences were then compared to public databases to gauge their similarity to known sequences, as described in Example 21 below.

EXAMPLE 21

Clustering of the 5' ESTs and Calculation of Novelty Indices for cDNA Libraries

25 For each sequenced EST library, the sequences were clustered by the 5' end. Each sequence in the library was compared to the others with BLASTN2 (direct strand, parameters S=107). ESTs with High Scoring Segment Pairs (HSPs) at least 25 bp long, having 95% identical bases and beginning closer than 10 bp from each EST 5' end were grouped. The longest sequence found in the cluster was used as representative of the group. A global
30 clustering between libraries was then performed leading to the definition of super-contigs.

To assess the yield of new sequences within the EST libraries, a novelty rate (NR) was defined as: $NR = 100 \times (\text{Number of new unique sequences found in the library} / \text{Total number of sequences from the library})$. Typically, novelty rating ranged between 10% and 41% depending on the tissue from which the EST library was obtained. For most of the libraries, the random sequencing of 5' EST libraries was pursued until the novelty rate reached 20%.

Following characterization as described above, the collection of 5' ESTs in NetGene™ was screened to identify those 5' ESTs bearing potential signal sequences as described in Example 22 below.

EXAMPLE 22

Identification of Potential Signal Sequences in 5' ESTs

The 5' ESTs in the NetGene™ database were screened to identify those having an uninterrupted open reading frame (ORF) longer than 45 nucleotides beginning with an ATG codon and extending to the end of the EST. Approximately half of the cDNA sequences in NetGene™ contained such an ORF. The ORFs of these 5' ESTs were then searched to identify potential signal motifs using slight modifications of the procedures disclosed in Von Heijne, *Nucleic Acids Res.* 14:4683-4690, 1986, the disclosure of which is incorporated herein by reference. Those 5' EST sequences encoding a stretch of at least 15 amino acid long with a score of at least 3.5 in the Von Heijne signal peptide identification matrix were considered to possess a signal sequence. Those 5' ESTs which matched a known human mRNA or EST sequence and had a 5' end more than 20 nucleotides downstream of the known 5' end were excluded from further analysis. The remaining cDNAs having signal sequences therein were included in a database called SignalTag™.

To confirm the accuracy of the above method for identifying signal sequences, the analysis of Example 23 was performed.

EXAMPLE 23**Confirmation of Accuracy of Identification of Potential Signal Sequences in 5' ESTs**

The accuracy of the above procedure for identifying signal sequences encoding signal peptides was evaluated by applying the method to the 43 amino acids located at the N terminus of all human SwissProt proteins. The computed Von Heijne score for each protein was compared with the known characterization of the protein as being a secreted protein or a non-secreted protein. In this manner, the number of non-secreted proteins having a score higher than 3.5 (false positives) and the number of secreted proteins having a score lower than 3.5 (false negatives) could be calculated.

Using the results of the above analysis, the probability that a peptide encoded by the 5' region of the mRNA is in fact a genuine signal peptide based on its Von Heijne's score was calculated based on either the assumption that 10% of human proteins are secreted or the assumption that 20% of human proteins are secreted. The results of this analysis are shown in Figure 2 and table IV.

Using the above method of identification of secretory proteins, 5' ESTs of the following polypeptides known to be secreted were obtained: human glucagon, gamma interferon induced monokine precursor, secreted cyclophilin-like protein, human pleiotropin, and human biotinidase precursor. Thus, the above method successfully identified those 5' ESTs which encode a signal peptide.

To confirm that the signal peptide encoded by the 5' ESTs actually functions as a signal peptide, the signal sequences from the 5' ESTs may be cloned into a vector designed for the identification of signal peptides. Such vectors are designed to confer the ability to grow in selective medium only to host cells containing a vector with an operably linked signal sequence. For example, to confirm that a 5' EST encodes a genuine signal peptide, the signal sequence of the 5' EST may be inserted upstream and in frame with a non-secreted form of the yeast invertase gene in signal peptide selection vectors such as those described in U.S. Patent No. 5,536,637, the disclosure of which is incorporated herein by reference. Growth of host cells containing signal sequence selection vectors with the correctly inserted 5' EST signal sequence confirms that the 5' EST encodes a genuine signal peptide.

Alternatively, the presence of a signal peptide may be confirmed by cloning the extended cDNAs obtained using the ESTs into expression vectors such as pXT1 (as described below in example 30), or by constructing promoter-signal sequence-reporter gene vectors which encode fusion proteins between the signal peptide and an assayable reporter protein. After introduction of these vectors into a suitable host cell, such as COS cells or NIH 3T3 cells, the growth medium may be harvested and analyzed for the presence of the secreted protein. The medium from these cells is compared to the medium from control cells containing vectors lacking the signal sequence or extended cDNA insert to identify vectors which encode a functional signal peptide or an authentic secreted protein.

Those 5' ESTs which encoded a signal peptide, as determined by the method of Example 22 above, were further grouped into four categories based on their homology to known sequences as described in Example 24 below.

EXAMPLE 24

Categorization of 5' ESTs Encoding a Signal Peptide

Those 5' ESTs having a sequence not matching any known vertebrate sequence nor any publicly available EST sequence were designated "new." Of the sequences in the SignalTag™ database, 947 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those 5' ESTs having a sequence not matching any vertebrate sequence but matching a publicly known EST were designated "EST-ext", provided that the known EST sequence was extended by at least 40 nucleotides in the 5' direction. Of the sequences in the SignalTag™ database, 150 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those ESTs not matching any vertebrate sequence but matching a publicly known EST without extending the known EST by at least 40 nucleotides in the 5' direction were designated "EST." Of the sequences in the SignalTag™ database, 599 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those 5' ESTs matching a human mRNA sequence but extending the known sequence by at least 40 nucleotides in the 5' direction were designated "VERT-ext." Of the sequences in the SignalTag™ database, 23 of the 5' ESTs having a Von Heijne's score of at

least 3.5 fell into this category. Included in this category was a 5' EST which extended the known sequence of the human translocase mRNA by more than 200 bases in the 5' direction.

A 5' EST which extended the sequence of a human tumor suppressor gene in the 5' direction was also identified.

- 5 Table V shows the distribution of 5' ESTs in each category and the number of 5' ESTs in each category having a given minimum von Heijne's score.

3. Evaluation of Spatial and Temporal Expression of mRNAs Corresponding to the 5'ESTs or Extended cDNAs

- 10 Each of the 5' ESTs was also categorized based on the tissue from which its corresponding mRNA was obtained, as described below in Example 25.

EXAMPLE 25

Categorization of Expression Patterns

- 15 Table VI shows the distribution of 5' ESTs in each of the above defined category with respect to the tissue from which the 5'ESTs of the corresponding mRNA were obtained.

- Table II provides the sequence identification numbers of 5' EST sequences derived from brain, the categories in which these sequences fall, and the von Heijne's score of the signal peptides which they encode. The 5' EST sequences and the amino acid sequences they encode are provided in the appended sequence listings. Table III provides the sequence ID
20 numbers of the 5' ESTs and the sequences of the signal peptides which they encode. The sequences of the 5' ESTs and the polypeptides they encode are provided in the sequence listing appended hereto.

- The sequences of DNA SEQ ID NOs: 38-195 can readily be screened for any errors
25 therein and any sequence ambiguities can be resolved by resequencing a fragment containing such errors or ambiguities on both strands. Such fragments may be obtained from the plasmids stored in the inventors' laboratory or can be isolated using the techniques described herein. Resolution of any such ambiguities or errors may be facilitated by using primers which hybridize to sequences located close to the ambiguous or erroneous sequences. For
30 example, the primers may hybridize to sequences within 50-75 bases of the ambiguity or

error. Upon resolution of an error or ambiguity, the corresponding corrections can be made in the protein sequences encoded by the DNA containing the error or ambiguity.

In addition to categorizing the 5' ESTs with respect to their tissue of origin, the spatial and temporal expression patterns of the mRNAs corresponding to the 5' ESTs, as well as their expression levels, may be determined as described in Example 26 below. Characterization of the spatial and temporal expression patterns and expression levels of these mRNAs is useful for constructing expression vectors capable of producing a desired level of gene product in a desired spatial or temporal manner, as will be discussed in more detail below.

Furthermore, 5' ESTs whose corresponding mRNAs are associated with disease states may also be identified. For example, a particular disease may result from the lack of expression, over expression, or under expression of an mRNA corresponding to a 5' EST. By comparing mRNA expression patterns and quantities in samples taken from healthy individuals with those from individuals suffering from a particular disease, 5' ESTs responsible for the disease may be identified.

It will be appreciated that the results of the above characterization procedures for 5' ESTs also apply to extended cDNAs (obtainable as described below) which contain sequences adjacent to the 5' ESTs. It will also be appreciated that if desired, characterization may be delayed until extended cDNAs have been obtained rather than characterizing the ESTs themselves.

EXAMPLE 26

Evaluation of Expression Levels and Patterns of mRNAs

Corresponding to 5' ESTs or Extended cDNAs

Expression levels and patterns of mRNAs corresponding to 5' ESTs or extended cDNAs (obtainable as described below in example 27) may be analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277, the entire contents of which are hereby incorporated by reference. Briefly, a 5' EST, extended cDNA, or fragment thereof corresponding to the gene encoding the mRNA to be characterized is inserted at a cloning site immediately downstream of a bacteriophage (T3,

T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the 5' EST or extended cDNA has 100 or more nucleotides. The plasmid is linearized and transcribed in the presence of ribonucleotides comprising modified ribonucleotides (*i.e.* biotin-UTP and DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50°C for 16 hours in an 80% formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (*i.e.* RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence of the DIG modification enables the hybrid to be detected and quantified by ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

The 5' ESTs, extended cDNAs, or fragments thereof may also be tagged with nucleotide sequences for the serial analysis of gene expression (SAGE) as disclosed in UK Patent Application No. 2 305 241 A, the entire contents of which are incorporated by reference. In this method, cDNAs are prepared from a cell, tissue, organism or other source of nucleic acid for which gene expression patterns must be determined. The resulting cDNAs are separated into two pools. The cDNAs in each pool are cleaved with a first restriction endonuclease, called an anchoring enzyme, having a recognition site which is likely to be present at least once in most cDNAs. The fragments which contain the 5' or 3' most region of the cleaved cDNA are isolated by binding to a capture medium such as streptavidin coated beads. A first oligonucleotide linker having a first sequence for hybridization of an amplification primer and an internal restriction site for a so-called tagging endonuclease is ligated to the digested cDNAs in the first pool. Digestion with the second endonuclease produces short tag fragments from the cDNAs.

A second oligonucleotide having a second sequence for hybridization of an amplification primer and an internal restriction site is ligated to the digested cDNAs in the second pool. The cDNA fragments in the second pool are also digested with the tagging endonuclease to generate short tag fragments derived from the cDNAs in the second pool. The tags resulting from digestion of the first and second pools with the anchoring enzyme and the tagging endonuclease are ligated to one another to produce so-called ditags. In some embodiments, the ditags are concatamerized to produce ligation products containing from 2

to 200 ditags. The tag sequences are then determined and compared to the sequences of the 5' ESTs or extended cDNAs to determine which 5' ESTs or extended cDNAs are expressed in the cell, tissue, organism, or other source of nucleic acids from which the tags were derived. In this way, the expression pattern of the 5' ESTs or extended cDNAs in the cell, tissue, organism, or other source of nucleic acids is obtained.

Quantitative analysis of gene expression may also be performed using arrays. As used herein, the term array means a one dimensional, two dimensional, or multidimensional arrangement of full length cDNAs (*i.e.* extended cDNAs which include the coding sequence for the signal peptide, the coding sequence for the mature protein, and a stop codon), extended cDNAs, 5' ESTs or fragments thereof of sufficient length to permit specific detection of gene expression. Preferably, the fragments are at least 15 nucleotides in length. More preferably, the fragments are at least 100 nucleotide long. More preferably, the fragments are more than 100 nucleotides in length. In some embodiments, the fragments may be more than 500 nucleotide long.

For example, quantitative analysis of gene expression may be performed with full length cDNAs as defined below, extended cDNAs, 5' ESTs, or fragments thereof in a complementary DNA microarray as described by Schena *et al.* (*Science* 270:467-470, 1995; *Proc. Natl. Acad. Sci. U.S.A.* 93:10614-10619, 1996). Full length cDNAs, extended cDNAs, 5' ESTs or fragments thereof are amplified by PCR and arrayed from 96-well microtiter plates onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2 min at 95°C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25°C.

Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm² microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60°C. Arrays are washed for 5 min at 25°C in low stringency wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a fluorescence laser scanning device fitted with a custom filter set. Accurate differential

expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

Quantitative analysis of the expression of genes may also be performed with full length cDNAs, extended cDNAs, 5' ESTs, or fragments thereof in complementary DNA arrays as described by Pietu *et al.* (*Genome Research* 6:492-503, 1996). The full length cDNAs, extended cDNAs, 5' ESTs or fragments thereof are PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis of the 5' ESTs or extended cDNAs can be done through high density nucleotide arrays as described by Lockhart *et al.* (*Nature Biotechnology* 14: 1675-1680, 1996) and Sosnowsky *et al.* (*Proc. Natl. Acad. Sci.* 94:1119-1123, 1997). Oligonucleotides of 15-50 nucleotides corresponding to sequences of the 5' ESTs or extended cDNAs are synthesized directly on the chip (Lockhart *et al.*, *supra*) or synthesized and then addressed to the chip (Sosnowsky *et al.*, *supra*). Preferably, the oligonucleotides are about 20 nucleotides in length.

cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the chip. After washing as described in Lockhart *et al.*, *supra* and application of different electric fields (Sonowsky *et al.*, *supra.*), the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed. Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of the mRNA corresponding to the 5' EST or extended cDNA from which the oligonucleotide sequence has been designed.

III. Use of 5' ESTs to Clone Extended cDNAs and to Clone the Corresponding Genomic DNAs

Once 5' ESTs which include the 5' end of the corresponding mRNAs have been selected using the procedures described above, they can be utilized to isolate extended cDNAs which contain sequences adjacent to the 5' ESTs. The extended cDNAs may include the entire coding sequence of the protein encoded by the corresponding mRNA, including the authentic translation start site, the signal sequence, and the sequence encoding the mature protein remaining after cleavage of the signal peptide. Such extended cDNAs are referred to herein as "full length cDNAs." Alternatively, the extended cDNAs may include only the sequence encoding the mature protein remaining after cleavage of the signal peptide, or only the sequence encoding the signal peptide.

Example 27 below describes a general method for obtaining extended cDNAs using 5' ESTs. Example 28 below provides experimental results, using the method explained in example 27, describing several extended cDNAs including the entire coding sequence and authentic 5' end of the corresponding mRNA for several secreted proteins.

The methods of Examples 27, 28, and 29 can also be used to obtain extended cDNAs which encode less than the entire coding sequence of the secreted proteins encoded by the genes corresponding to the 5' ESTs. In some embodiments, the extended cDNAs isolated using these methods encode at least 10 amino acids of one of the proteins encoded by the sequences of SEQ ID NOs: 38-195. In further embodiments, the extended cDNAs encode at least 20 amino acids of the proteins encoded by the sequences of SEQ ID NOs: 38-195. In further embodiments, the extended cDNAs encode at least 30 amino acids of the sequences of SEQ ID NOs: 38-195. In a preferred embodiment, the extended cDNAs encode a full length protein sequence, which includes the protein coding sequences of SEQ ID NOs: 38-195.

EXAMPLE 27

General Method for Using 5' ESTs to Clone and Sequence cDNAs which Include the Entire Coding Region and the Authentic 5' End of the Corresponding mRNA

The following general method has been used to quickly and efficiently isolate extended cDNAs having the authentic 5' ends of their corresponding mRNAs as well as

the full protein coding sequence and including sequence adjacent to the sequences of the 5' ESTs used to obtain them. This method may be applied to obtain extended cDNAs for any 5' EST in the NetGene™ database, including those 5' ESTs encoding polypeptides belonging to secreted proteins. The method is summarized in figure 3.

5

1. Obtention of Extended cDNAs

a) First strand synthesis

The method takes advantage of the known 5' sequence of the mRNA. A reverse transcription reaction is conducted on purified mRNA with a poly 14dT primer containing a 49 nucleotide sequence at its 5' end allowing the addition of a known sequence at the end of the cDNA which corresponds to the 3' end of the mRNA. For example, the primer may have the following sequence: 5'-ATC GTT GAG ACT CGT ACC AGC AGA GTC ACG AGA GAG ACT ACA CGG TAC TGG TTT TTT TTT TTT TTVN -3' (SEQ ID NO:14). Those skilled in the art will appreciate that other sequences may also be added to the poly dT sequence and used to prime the first strand synthesis. Using this primer and a reverse transcriptase such as the Superscript II (Gibco BRL) or Rnase H Minus M-MLV (Promega) enzyme, a reverse transcript anchored at the 3' polyA site of the RNAs is generated.

15

After removal of the mRNA hybridized to the first cDNA strand by alkaline hydrolysis, the products of the alkaline hydrolysis and the residual poly dT primer are eliminated with an exclusion column such as an AcA34 (Biosepra) matrix as explained in Example 11.

20

b) Second strand synthesis

A pair of nested primers on each end is designed based on the known 5' sequence from the 5' EST and the known 3' end added by the poly dT primer used in the first strand synthesis. Softwares used to design primers are either based on GC content and melting temperatures of oligonucleotides, such as OSP (Illier and Green, *PCR Meth. Appl.* 1:124-128, 1991), or based on the octamer frequency disparity method (Griffais *et al.*, *Nucleic Acids Res.* 19: 3887-3891, 1991) such as PC-Rare (<http://bioinformatics.weizmann.ac.il/software/PC-Rare/doc/manuel.html>).

25

Preferably, the nested primers at the 5' end are separated from one another by four to nine bases. The 5' primer sequences may be selected to have melting temperatures and specificities suitable for use in PCR.

Preferably, the nested primers at the 3' end are separated from one another by four to nine bases. For example, the nested 3' primers may have the following sequences: (5'- CCA GCA GAG TCA CGA GAG AGA CTA CAC GG -3'(SEQ ID NO:15), and 5'- CAC GAG AGA GAC TAC ACG GTA CTG G -3' (SEQ ID NO:16). These primers were selected because they have melting temperatures and specificities compatible with their use in PCR. However, those skilled in the art will appreciate that other sequences may also be used as primers.

The first PCR run of 25 cycles is performed using the Advantage Tth Polymerase Mix (Clontech) and the outer primer from each of the nested pairs. A second 20 cycle PCR using the same enzyme and the inner primer from each of the nested pairs is then performed on 1/2500 of the first PCR product. Thereafter, the primers and nucleotides are removed.

2. Sequencing of Full Length Extended cDNAs or Fragments Thereof

Due to the lack of position constraints on the design of 5' nested primers compatible for PCR use using the OSP software, amplicons of two types are obtained. Preferably, the second 5' primer is located upstream of the translation initiation codon thus yielding a nested PCR product containing the whole coding sequence. Such a full length extended cDNA undergoes a direct cloning procedure as described in section a. However, in some cases, the second 5' primer is located downstream of the translation initiation codon, thereby yielding a PCR product containing only part of the ORF. Such incomplete PCR products are submitted to a modified procedure described in section b.

a) Nested PCR products containing complete ORFs

When the resulting nested PCR product contains the complete coding sequence, as predicted from the 5'EST sequence, it is cloned in an appropriate vector such as pED6dpc2, as described in section 3.

b) Nested PCR products containing incomplete ORFs

When the amplicon does not contain the complete coding sequence, intermediate steps are necessary to obtain both the complete coding sequence and a PCR product containing the full coding sequence. The complete coding sequence can be assembled from several partial sequences determined directly from different PCR products as described in the following section.

Once the full coding sequence has been completely determined, new primers compatible for PCR use are designed to obtain amplicons containing the whole coding region. However, in such cases, 3' primers compatible for PCR use are located inside the 3' UTR of the corresponding mRNA, thus yielding amplicons which lack part of this region, *i.e.* the polyA tract and sometimes the polyadenylation signal, as illustrated in figure 3. Such full length extended cDNAs are then cloned into an appropriate vector as described in section 3.

c) Sequencing extended cDNAs

Sequencing of extended cDNAs is performed using a Die Terminator approach with the AmpliTaq DNA polymerase FS kit available from Perkin Elmer.

In order to sequence PCR fragments, primer walking is performed using software such as OSP to choose primers and automated computer software such as ASMG (Sutton *et al.*, *Genome Science Technol.* 1: 9-19, 1995) to construct contigs of walking sequences including the initial 5' tag using minimum overlaps of 32 nucleotides. Preferably, primer walking is performed until the sequences of full length cDNAs are obtained.

Completion of the sequencing of a given extended cDNA fragment is assessed as follows. Since sequences located after a polyA tract are difficult to determine precisely in the case of uncloned products, sequencing and primer walking processes for PCR products are interrupted when a polyA tract is identified in extended cDNAs obtained as described in case b. The sequence length is compared to the size of the nested PCR product obtained as described above. Due to the limited accuracy of the determination of the PCR product size by gel electrophoresis, a sequence is considered complete if the size of the obtained sequence is at least 70 % the size of the first nested PCR product. If the length of the sequence determined from the computer analysis is not at least 70% of the length of the nested PCR product, these PCR products are cloned and the sequence of the insertion is determined. When Northern blot data are available, the size of the mRNA detected for a given PCR

product is used to finally assess that the sequence is complete. Sequences which do not fulfill the above criteria are discarded and will undergo a new isolation procedure.

Sequence data of all extended cDNAs are then transferred to a proprietary database, where quality controls and validation steps are carried out as described in
5 example 15.

3. Cloning of Full Length Extended cDNAs

The PCR product containing the full coding sequence is then cloned in an appropriate vector. For example, the extended cDNAs can be cloned into the expression vector
10 pED6dpc2 (DiscoverEase, Genetics Institute, Cambridge, MA) as follows. pED6dpc2 vector DNA is prepared with blunt ends by performing an EcoRI digestion followed by a fill in reaction. The blunt ended vector is dephosphorylated. After removal of PCR primers and ethanol precipitation, the PCR product containing the full coding sequence or the extended
15 cDNA obtained as described above is phosphorylated with a kinase subsequently removed by phenol-Sevag extraction and precipitation. The double stranded extended cDNA is then ligated to the vector and the resulting expression plasmid introduced into appropriate host cells.

Since the PCR products obtained as described above are blunt ended molecules that can be cloned in either direction, the orientation of several clones for each PCR product is
20 determined. Then, 4 to 10 clones are ordered in microtiter plates and subjected to a PCR reaction using a first primer located in the vector close to the cloning site and a second primer located in the portion of the extended cDNA corresponding to the 3' end of the mRNA. This second primer may be the antisense primer used in anchored PCR in the case of direct cloning (case a) or the antisense primer located inside the 3'UTR in the case of indirect cloning (case
25 b). Clones in which the start codon of the extended cDNA is operably linked to the promoter in the vector so as to permit expression of the protein encoded by the extended cDNA are conserved and sequenced. In addition to the ends of cDNA inserts, approximately 50 bp of vector DNA on each side of the cDNA insert are also sequenced.

The cloned PCR products are then entirely sequenced according to the
30 aforementioned procedure. In this case, contiguation of long fragments is then performed on walking sequences that have already contiguated for uncloned PCR products during

primer walking. Sequencing of cloned amplicons is complete when the resulting contigs include the whole coding region as well as overlapping sequences with vector DNA on both ends.

5 4. Computer analysis of Full Length Extended cDNA

Sequences of all full length extended cDNAs are then submitted to further analysis as described below. Before searching the extended full length cDNAs for sequences of interest, extended cDNAs which are not of interest (vector RNAs, transfer RNAs, ribosomal RNAs, mitochondrial RNAs, prokaryotic RNAs and fungal RNAs) are discarded using methods essentially similar to those described for 5'ESTs in Example 18.

10 *a) Identification of structural features*

Structural features, e.g. polyA tail and polyadenylation signal, of the sequences of full length extended cDNAs are subsequently determined as follows.

15 A polyA tail is defined as a homopolymeric stretch of at least 11 A with at most one alternative base within it. The polyA tail search is restricted to the last 100 nt of the sequence and limited to stretches of 11 consecutive A's because sequencing reactions are often not readable after such a polyA stretch. Stretches having more than 90% homology over 8 nucleotides are identified as polyA tails using BLAST2N.

20 To search for a polyadenylation signal, the polyA tail is clipped from the full-length sequence. The 50 bp preceding the polyA tail are first searched for the canonic polyadenylation AAUAAA signal and, if the canonic signal is not detected, for the alternative AUUAAA signal (Sheets *et al.*, *Nuc. Acids Res.* 18: 5799-5805, 1990). If neither of these consensus polyadenylation signals is found, the canonic motif is searched again allowing one mismatch to account for possible sequencing errors. More than 85 %
25 of identified polyadenylation signals of either type actually ends 10 to 30 bp from the polyA tail. Alternative AUUAAA signals represents approximately 15 % of the total number of identified polyadenylation signals.

b) Identification of functional features

30 Functional features, e.g. ORFs and signal sequences, of the sequences of full length extended cDNAs were subsequently determined as follows.

The 3 upper strand frames of extended cDNAs are searched for ORFs defined as the maximum length fragments beginning with a translation initiation codon and ending with a stop codon. ORFs encoding at least 20 amino acids are preferred.

Each found ORF is then scanned for the presence of a signal peptide in the first 50 amino-acids or, where appropriate, within shorter regions down to 20 amino acids or less in the ORF, using the matrix method of von Heijne (*Nuc. Acids Res.* 14: 4683-4690, 1986), the disclosure of which is incorporated herein by reference as described in Example 22.

c) Homology to either nucleotidic or proteic sequences

Categorization of full-length sequences may be achieved using procedures essentially similar to those described for 5'ESTs in Example 24.

Extended cDNAs prepared as described above may be subsequently engineered to obtain nucleic acids which include desired portions of the extended cDNA using conventional techniques such as subcloning, PCR, or *in vitro* oligonucleotide synthesis. For example, nucleic acids which include only the full coding sequences (*i.e.* the sequences encoding the signal peptide and the mature protein remaining after the signal peptide is cleaved off) may be obtained using techniques known to those skilled in the art. Alternatively, conventional techniques may be applied to obtain nucleic acids which contain only the coding sequences for the mature protein remaining after the signal peptide is cleaved off or nucleic acids which contain only the coding sequences for the signal peptides.

Similarly, nucleic acids containing any other desired portion of the coding sequences for the secreted protein may be obtained. For example, the nucleic acid may contain at least 10 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. In another embodiment, the nucleic acid may contain at least 15 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. Alternatively, the nucleic acid may contain at least 20 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. In another embodiment, the nucleic acid may contain at least 25 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. In yet another embodiment, the nucleic acid may contain at least 40

consecutive bases of an extended cDNA such as one of the extended cDNAs described below.

Once an extended cDNA has been obtained, it can be sequenced to determine the amino acid sequence it encodes. Once the encoded amino acid sequence has been
5 determined, one can create and identify any of the many conceivable cDNAs that will encode that protein by simply using the degeneracy of the genetic code. For example, allelic variants or other homologous nucleic acids can be identified as described below. Alternatively, nucleic acids encoding the desired amino acid sequence can be synthesized *in vitro*.

In a preferred embodiment, the coding sequence may be selected using the known
10 codon or codon pair preferences for the host organism in which the cDNA is to be expressed.

The extended cDNAs derived from the 5' ESTS of the present invention were obtained as described in Example 28 below.

EXAMPLE 28

15 Characterization of cloned extended cDNAs obtained using 5' ESTs

The procedure described in Example 27 above was used to obtain the extended cDNAs derived from the 5' ESTs of the present invention in a variety of tissues. The following list provides a few examples of thus obtained extended cDNAs.

Using this approach, the full length cDNA of SEQ ID NO:17 (internal identification
20 number 48-19-3-G1-FL1) was obtained. This cDNA falls into the "EST-ext" category described above and encodes the signal peptide MKKVLLLITAILAVAVG (SEQ ID NO:18) having a von Heijne score of 8.2.

The full length cDNA of SEQ ID NO:19 (internal identification number 58-34-2-E7-
25 FL2) was also obtained using this procedure. This cDNA falls into the "EST-ext" category described above and encodes the signal peptide MWWFQQGLSFLPSALVIWTS (SEQ ID NO:20) having a von Heijne score of 5.5.

Another full length cDNA obtained using the procedure described above has the sequence of SEQ ID NO:21 (internal identification number 51-27-1-E8-FL1). This cDNA, falls into the "EST-ext" category described above and encodes the signal peptide
30 MVLTTLPANSANSPVNMPTTGPNLSYASSALSPCLT (SEQ ID NO:22) having a von Heijne score of 5.9.

The above procedure was also used to obtain a full length cDNA having the sequence of SEQ ID NO:23 (internal identification number 76-4-1-G5-FL1). This cDNA falls into the "EST-ext" category described above and encodes the signal peptide ILSTVTALTFAXA (SEQ ID NO:24) having a von Heijne score of 5.5.

5 The full length cDNA of SEQ ID NO:25 (internal identification number 51-3-3-B10-FL3) was also obtained using this procedure. This cDNA falls into the "new" category described above and encodes a signal peptide LVLTLCTLPLAVA (SEQ ID NO:26) having a von Heijne score of 10.1.

10 The full length cDNA of SEQ ID NO:27 (internal identification number 58-35-2-F10-FL2) was also obtained using this procedure. This cDNA falls into the "new" category described above and encodes a signal peptide LWLLFFLVTAIHA (SEQ ID NO:28) having a von Heijne score of 10.7.

15 Bacterial clones containing plasmids containing the full length cDNAs described above are presently stored in the inventor's laboratories under the internal identification numbers provided above. The inserts may be recovered from the stored materials by growing an aliquot of the appropriate bacterial clone in the appropriate medium. The plasmid DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a cesium chloride gradient, size exclusion chromatography, or anion exchange chromatography. The plasmid DNA
20 obtained using these procedures may then be manipulated using standard cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the cDNA insertion. The PCR product which corresponds to the cDNA can then be manipulated using standard cloning techniques familiar to those skilled in the art.

25 The polypeptides encoded by the extended cDNAs may be screened for the presence of known structural or functional motifs or for the presence of signatures, small amino acid sequences which are well conserved amongst the members of a protein family. The conserved regions have been used to derive consensus patterns or matrices included in the PROSITE data bank, in particular in the file prosite.dat (Release 13.0 of November 1995,
30 located at <http://expasy.hcuge.ch/sprot/prosite.html>. Prosite_convert and prosite_scan

programs (http://ulrec3.unil.ch/ftpserveur/prosite_scan) may be used to find signatures on the extended cDNAs.

For each pattern obtained with the `prosite_convert` program from the `prosite.dat` file, the accuracy of the detection on a new protein sequence may be assessed by evaluating the frequency of irrelevant hits on the population of human secreted proteins included in the data bank SWISSPROT. The ratio between the number of hits on shuffled proteins (with a window size of 20 amino acids) and the number of hits on native (unshuffled) proteins may be used as an index. Every pattern for which the ratio is greater than 20% (one hit on shuffled proteins for 5 hits on native proteins) may be skipped during the search with `prosite_scan`.
The program used to shuffle protein sequences (`db_shuffled`) and the program used to determine the statistics for each pattern in the protein data banks (`prosite_statistics`) are available on the ftp site http://ulrec3.unil.ch/ftpserveur/prosite_scan.

In addition to PCR based methods for obtaining extended cDNAs, traditional hybridization based methods may also be employed. These methods may also be used to obtain the genomic DNAs which encode the mRNAs from which the 5' ESTs were derived, mRNAs corresponding to the extended cDNAs, or nucleic acids which are homologous to extended cDNAs or 5' ESTs. Example 29 below provides examples of such methods.

EXAMPLE 29

Methods for Obtaining cDNAs which include the Entire Coding Region and the Authentic 5' End of the Corresponding mRNA

A full length cDNA library can be made using the strategies described in Examples 13, 14, 15, and 16 above by replacing the random nonamer used in Example 14 with an oligo-dT primer. For instance, the oligonucleotide of SEQ ID NO:14 may be used.

Alternatively, a cDNA library or genomic DNA library may be obtained from a commercial source or made using techniques familiar to those skilled in the art. Such cDNA or genomic DNA libraries may be used to isolate extended cDNAs obtained from 5' EST or nucleic acids homologous to extended cDNAs or 5' EST as follows. The cDNA library or genomic DNA library is hybridized to a detectable probe comprising at least 10 consecutive nucleotides from the 5' EST or extended cDNA using conventional techniques. Preferably,

the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST or extended cDNA. More preferably, the probe comprises at least 20 to 30 consecutive nucleotides from the 5' EST or extended cDNA. In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST or extended cDNA.

5 Techniques for identifying cDNA clones in a cDNA library which hybridize to a given probe sequence are disclosed in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual 2d Ed.*, Cold Spring Harbor Laboratory Press, 1989, the disclosure of which is incorporated herein by reference. The same techniques may be used to isolate genomic DNAs.

 Briefly, cDNA or genomic DNA clones which hybridize to the detectable probe are
10 identified and isolated for further manipulation as follows. A probe comprising at least 10 consecutive nucleotides from the 5' EST or extended cDNA is labeled with a detectable label such as a radioisotope or a fluorescent molecule. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST or extended cDNA. More preferably, the probe comprises 20 to 30 consecutive nucleotides from the 5' EST or extended cDNA. In
15 some embodiments, the probe comprises more than 30 nucleotides from the 5' EST or extended cDNA.

 Techniques for labeling the probe are well known and include phosphorylation with polynucleotide kinase, nick translation, *in vitro* transcription, and non radioactive techniques. The cDNAs or genomic DNAs in the library are transferred to a nitrocellulose or nylon filter
20 and denatured. After blocking of non specific sites, the filter is incubated with the labeled probe for an amount of time sufficient to allow binding of the probe to cDNAs or genomic DNAs containing a sequence capable of hybridizing thereto.

 By varying the stringency of the hybridization conditions used to identify extended cDNAs or genomic DNAs which hybridize to the detectable probe, extended
25 cDNAs having different levels of homology to the probe can be identified and isolated as described below.

1. Identification of Extended cDNA or Genomic cDNA Sequences Having a High Degree of Homology to the Labeled Probe

To identify extended cDNAs or genomic DNAs having a high degree of homology to the probe sequence, the melting temperature of the probe may be calculated using the following formulas:

For probes between 14 and 70 nucleotides in length the melting temperature (T_m) is calculated using the formula: $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (600/N)$ where N is the length of the probe.

If the hybridization is carried out in a solution containing formamide, the melting temperature may be calculated using the equation $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (0.63\% \text{ formamide}) - (600/N)$ where N is the length of the probe.

Prehybridization may be carried out in 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100 μ g denatured fragmented salmon sperm DNA or 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100 μ g denatured fragmented salmon sperm DNA, 50% formamide. The formulas for SSC and Denhardt's solutions are listed in Sambrook *et al.*, *supra*.

Hybridization is conducted by adding the detectable probe to the prehybridization solutions listed above. Where the probe comprises double stranded DNA, it is denatured before addition to the hybridization solution. The filter is contacted with the hybridization solution for a sufficient period of time to allow the probe to hybridize to extended cDNAs or genomic DNAs containing sequences complementary thereto or homologous thereto. For probes over 200 nucleotides in length, the hybridization may be carried out at 15-25°C below the T_m . For shorter probes, such as oligonucleotide probes, the hybridization may be conducted at 15-25°C below the T_m . Preferably, for hybridizations in 6X SSC, the hybridization is conducted at approximately 68°C. Preferably, for hybridizations in 50% formamide containing solutions, the hybridization is conducted at approximately 42°C.

All of the foregoing hybridizations would be considered to be under "stringent" conditions.

Following hybridization, the filter is washed in 2X SSC, 0.1% SDS at room temperature for 15 minutes. The filter is then washed with 0.1X SSC, 0.5% SDS at room temperature for 30 minutes to 1 hour. Thereafter, the solution is washed at the hybridization

temperature in 0.1X SSC, 0.5% SDS. A final wash is conducted in 0.1X SSC at room temperature.

Extended cDNAs, nucleic acids homologous to extended cDNAs or 5' ESTs, or genomic DNAs which have hybridized to the probe are identified by autoradiography or other conventional techniques.

2. Obtention of Extended cDNA or Genomic cDNA Sequences Having Lower Degrees of Homology to the Labeled Probe

The above procedure may be modified to identify extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs having decreasing levels of homology to the probe sequence. For example, to obtain extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs of decreasing homology to the detectable probe, less stringent conditions may be used. For example, the hybridization temperature may be decreased in increments of 5°C from 68°C to 42°C in a hybridization buffer having a sodium concentration of approximately 1M. Following hybridization, the filter may be washed with 2X SSC, 0.5% SDS at the temperature of hybridization. These conditions are considered to be "moderate" conditions above 50°C and "low" conditions below 50°C.

Alternatively, the hybridization may be carried out in buffers, such as 6X SSC, containing formamide at a temperature of 42°C. In this case, the concentration of formamide in the hybridization buffer may be reduced in 5% increments from 50% to 0% to identify clones having decreasing levels of homology to the probe. Following hybridization, the filter may be washed with 6X SSC, 0.5% SDS at 50°C. These conditions are considered to be "moderate" conditions above 25% formamide and "low" conditions below 25% formamide.

Extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs which have hybridized to the probe are identified by autoradiography.

3. Determination of the Degree of Homology Between the Obtained Extended cDNAs and the Labeled Probe

If it is desired to obtain nucleic acids homologous to extended cDNAs, such as allelic variants thereof or nucleic acids encoding proteins related to the proteins encoded by the extended cDNAs, the level of homology between the hybridized nucleic acid and the

extended cDNA or 5' EST used as the probe may be further determined using BLAST2N; parameters may be adapted depending on the sequence length and degree of homology studied. To determine the level of homology between the hybridized nucleic acid and the extended cDNA or 5'EST from which the probe was derived, the nucleotide sequences of the hybridized nucleic acid and the extended cDNA or 5'EST from which the probe was derived are compared. For example, using the above methods, nucleic acids having at least 95% nucleic acid homology to the extended cDNA or 5'EST from which the probe was derived may be obtained and identified. Similarly, by using progressively less stringent hybridization conditions one can obtain and identify nucleic acids having at least 90%, at least 85%, at least 80% or at least 75% homology to the extended cDNA or 5'EST from which the probe was derived.

To determine whether a clone encodes a protein having a given amount of homology to the protein encoded by the extended cDNA or 5' EST, the amino acid sequence encoded by the extended cDNA or 5' EST is compared to the amino acid sequence encoded by the hybridizing nucleic acid. Homology is determined to exist when an amino acid sequence in the extended cDNA or 5' EST is closely related to an amino acid sequence in the hybridizing nucleic acid. A sequence is closely related when it is identical to that of the extended cDNA or 5' EST or when it contains one or more amino acid substitutions therein in which amino acids having similar characteristics have been substituted for one another. Using the above methods and algorithms such as FASTA with parameters depending on the sequence length and degree of homology studied, one can obtain nucleic acids encoding proteins having at least 95%, at least 90%, at least 85%, at least 80% or at least 75% homology to the proteins encoded by the extended cDNA or 5'EST from which the probe was derived.

In addition to the above described methods, other protocols are available to obtain extended cDNAs using 5' ESTs as outlined in the following paragraphs.

Extended cDNAs may be prepared by obtaining mRNA from the tissue, cell, or organism of interest using mRNA preparation procedures utilizing polyA selection procedures or other techniques known to those skilled in the art. A first primer capable of hybridizing to the polyA tail of the mRNA is hybridized to the mRNA and a reverse transcription reaction is performed to generate a first cDNA strand.

The first cDNA strand is hybridized to a second primer containing at least 10 consecutive nucleotides of the sequences of SEQ ID NOs 38-195. Preferably, the primer comprises at least 12, 15, or 17 consecutive nucleotides from the sequences of SEQ ID NOs 38-195. More preferably, the primer comprises 20 to 30 consecutive nucleotides from the sequences of SEQ ID NOs 38-195. In some embodiments, the primer comprises more than 30 nucleotides from the sequences of SEQ ID NOs 38-195. If it is desired to obtain extended cDNAs containing the full protein coding sequence, including the authentic translation initiation site, the second primer used contains sequences located upstream of the translation initiation site. The second primer is extended to generate a second cDNA strand complementary to the first cDNA strand. Alternatively, RT-PCR may be performed as described above using primers from both ends of the cDNA to be obtained.

Extended cDNAs containing 5' fragments of the mRNA may be prepared by hybridizing an mRNA comprising the sequence of the 5'EST for which an extended cDNA is desired with a primer comprising at least 10 consecutive nucleotides of the sequences complementary to the 5'EST and reverse transcribing the hybridized primer to make a first cDNA strand from the mRNAs. Preferably, the primer comprises at least 12, 15, or 17 consecutive nucleotides from the 5'EST. More preferably, the primer comprises 20 to 30 consecutive nucleotides from the 5'EST.

Thereafter, a second cDNA strand complementary to the first cDNA strand is synthesized. The second cDNA strand may be made by hybridizing a primer complementary to sequences in the first cDNA strand to the first cDNA strand and extending the primer to generate the second cDNA strand.

The double stranded extended cDNAs made using the methods described above are isolated and cloned. The extended cDNAs may be cloned into vectors such as plasmids or viral vectors capable of replicating in an appropriate host cell. For example, the host cell may be a bacterial, mammalian, avian, or insect cell.

Techniques for isolating mRNA, reverse transcribing a primer hybridized to mRNA to generate a first cDNA strand, extending a primer to make a second cDNA strand complementary to the first cDNA strand, isolating the double stranded cDNA and cloning the double stranded cDNA are well known to those skilled in the art and are described in *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc. 1997 and Sambrook *et al.*,

Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, 1989, the entire disclosures of which are incorporated herein by reference.

Alternatively, procedures such as the one described in Example 29 may be used for obtaining full length cDNAs or extended cDNAs. In this approach, full length or extended cDNAs are prepared from mRNA and cloned into double stranded phagemids as follows. The cDNA library in the double stranded phagemids is then rendered single stranded by treatment with an endonuclease, such as the Gene II product of the phage F1, and an exonuclease (Chang *et al.*, *Gene* 127:95-8, 1993). A biotinylated oligonucleotide comprising the sequence of a 5' EST, or a fragment containing at least 10 nucleotides thereof, is hybridized to the single stranded phagemids. Preferably, the fragment comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST. More preferably, the fragment comprises 20-30 consecutive nucleotides from the 5' EST. In some procedures, the fragment may comprise more than 30 consecutive nucleotides from the 5' EST.

Hybrids between the biotinylated oligonucleotide and phagemids having inserts containing the 5' EST sequence are isolated by incubating the hybrids with streptavidin coated paramagnetic beads and retrieving the beads with a magnet (Fry *et al.*, *Biotechniques*, 13: 124-131, 1992). Thereafter, the resulting phagemids containing the 5' EST sequence are released from the beads and converted into double stranded DNA using a primer specific for the 5' EST sequence. Alternatively, protocols such as the Gene Trapper kit (Gibco BRL) may be used. The resulting double stranded DNA is transformed into bacteria. Extended cDNAs containing the 5' EST sequence are identified by colony PCR or colony hybridization.

Using any of the above described methods in section III, a plurality of extended cDNAs containing full length protein coding sequences or sequences encoding only the mature protein remaining after the signal peptide is cleaved off may be provided as cDNA libraries for subsequent evaluation of the encoded proteins or use in diagnostic assays as described below.

IV. Expression of Proteins Encoded by Extended cDNAs Isolated Using 5' ESTs

Extended cDNAs containing the full protein coding sequences of their corresponding mRNAs or portions thereof, such as cDNAs encoding the mature protein, may be used to

express the encoded secreted proteins or portions thereof as described in Example 30 below. If desired, the extended cDNAs may contain the sequences encoding the signal peptide to facilitate secretion of the expressed protein. It will be appreciated that a plurality of extended cDNAs containing the full protein coding sequences or portions thereof may be simultaneously cloned into expression vectors to create an expression library for analysis of the encoded proteins as described below.

EXAMPLE 30

Expression of the Proteins Encoded by the Genes Corresponding to 5'ESTS or Portions Thereof

To express the proteins encoded by the genes corresponding to 5' ESTs (or portions thereof), full length cDNAs containing the entire protein coding region or extended cDNAs containing sequences adjacent to the 5' ESTs (or portions thereof) are obtained as described in Examples 27-29 and cloned into a suitable expression vector. If desired, the nucleic acids may contain the sequences encoding the signal peptide to facilitate secretion of the expressed protein. The nucleic acids inserted into the expression vectors may also contain sequences upstream of the sequences encoding the signal peptide, such as sequences which regulate expression levels or sequences which confer tissue specific expression.

The nucleic acid encoding the protein or polypeptide to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The expression vector may be any of the mammalian, yeast, insect or bacterial expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and facilitate proper protein folding, the codon context and codon pairing of the sequence may be optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, *et al.*, U.S. Patent No. 5,082,767, incorporated herein by this reference.

The cDNA cloned into the expression vector may encode the entire protein (*i.e.* the signal peptide and the mature protein), the mature protein (*i.e.* the protein created by cleaving the signal peptide off), only the signal peptide or any other portion thereof.

The following is provided as one exemplary method to express the proteins encoded by the extended cDNAs corresponding to the 5' ESTs or the nucleic acids described above. First, the methionine initiation codon for the gene and the polyA signal of the gene are identified. If the nucleic acid encoding the polypeptide to be expressed lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the extended cDNA lacks a polyA signal, this sequence can be added to the construct by, for example, splicing out the polyA signal from pSG5 (Stratagene) using BglII and SalI restriction endonuclease enzymes and incorporating it into the mammalian expression vector pXT1 (Stratagene). pXT1 contains the LTRs and a portion of the *gag* gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex thymidine kinase promoter and the selectable neomycin gene. The extended cDNA or portion thereof encoding the polypeptide to be expressed is obtained by PCR from the bacterial vector using oligonucleotide primers complementary to the extended cDNA or portion thereof and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and BglII at the 5' end of the corresponding cDNA 3' primer, taking care to ensure that the extended cDNA is positioned with the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1 containing a poly A signal and prepared for this ligation (blunt/BglII).

The ligated product is transfected into mouse NIH 3T3 cells using Lipofectin (Life Technologies, Inc., Grand Island, New York) under conditions outlined in the product specification. Positive transfectants are selected after growing the transfected cells in 600 µg/ml G418 (Sigma, St. Louis, Missouri). Preferably the expressed protein is released into the culture medium, thereby facilitating purification.

Alternatively, the extended cDNAs may be cloned into pED6dpc2 as described above. The resulting pED6dpc2 constructs may be transfected into a suitable host cell, such as COS 1 cells. Methotrexate resistant cells are selected and expanded. Preferably, the protein expressed from the extended cDNA is released into the culture medium thereby facilitating purification.

Proteins in the culture medium are separated by gel electrophoresis. If desired, the proteins may be ammonium sulfate precipitated or separated based on size or charge prior to electrophoresis.

As a control, the expression vector lacking a cDNA insert is introduced into host cells
5 or organisms and the proteins in the medium are harvested. The secreted proteins present in the medium are detected using techniques familiar to those skilled in the art such as Coomassie blue or silver staining or using antibodies against the protein encoded by the extended cDNA.

Antibodies capable of specifically recognizing the protein of interest may be generated
10 using synthetic 15-mer peptides having a sequence encoded by the appropriate 5' EST, extended cDNA, or portion thereof. The synthetic peptides are injected into mice to generate antibody to the polypeptide encoded by the 5' EST, extended cDNA, or portion thereof.

Secreted proteins from the host cells or organisms containing an expression vector which contains the extended cDNA derived from a 5' EST or a portion thereof are compared
15 to those from the control cells or organism. The presence of a band in the medium from the cells containing the expression vector which is absent in the medium from the control cells indicates that the extended cDNA encodes a secreted protein. Generally, the band corresponding to the protein encoded by the extended cDNA will have a mobility near that expected based on the number of amino acids in the open reading frame of the extended
20 cDNA. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic cleavage.

Alternatively, if the protein expressed from the above expression vectors does not contain sequences directing its secretion, the proteins expressed from host cells containing an expression vector with an insert encoding a secreted protein or portion thereof can be
25 compared to the proteins expressed in control host cells containing the expression vector without an insert. The presence of a band in samples from cells containing the expression vector with an insert which is absent in samples from cells containing the expression vector without an insert indicates that the desired protein or portion thereof is being expressed. Generally, the band will have the mobility expected for the secreted protein or portion
30 thereof. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic cleavage.

The protein encoded by the extended cDNA may be purified using standard immunochromatography techniques. In such procedures, a solution containing the secreted protein, such as the culture medium or a cell extract, is applied to a column having antibodies against the secreted protein attached to the chromatography matrix. The secreted protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically bound secreted protein is then released from the column and recovered using standard techniques.

If antibody production is not possible, the extended cDNA sequence or portion thereof may be incorporated into expression vectors designed for use in purification schemes employing chimeric polypeptides. In such strategies, the coding sequence of the extended cDNA or portion thereof is inserted in frame with the gene encoding the other half of the chimera. The other half of the chimera may be β -globin or a nickel binding polypeptide. A chromatography matrix having antibody to β -globin or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites may be engineered between the β -globin gene or the nickel binding polypeptide and the extended cDNA or portion thereof. Thus, the two polypeptides of the chimera may be separated from one another by protease digestion.

One useful expression vector for generating β -globin chimerics is pSG5 (Stratagene), which encodes rabbit β -globin. Intron II of the rabbit β -globin gene facilitates splicing of the expressed transcript, and the polyadenylation signal incorporated into the construct increases the level of expression. These techniques as described are well known to those skilled in the art of molecular biology. Standard methods are published in methods texts such as Davis *et al.*, (*Basic Methods in Molecular Biology*, Davis, Dibner, and Battey, ed., Elsevier Press, NY, 1986) and many of the methods are available from Stratagene, Life Technologies, Inc., or Promega. Polypeptide may additionally be produced from the construct using *in vitro* translation systems such as the *In vitro* ExpressTM Translation Kit (Stratagene).

Following expression and purification of the secreted proteins encoded by the 5' ESTs, extended cDNAs, or fragments thereof, the purified proteins may be tested for the ability to bind to the surface of various cell types as described in Example 31 below. It will be appreciated that a plurality of proteins expressed from these cDNAs may be included in a

panel of proteins to be simultaneously evaluated for the activities specifically described below, as well as other biological roles for which assays for determining activity are available.

EXAMPLE 31

5 Analysis of Secreted Proteins to Determine Whether they Bind to the Cell Surface

The proteins encoded by the 5' ESTs, extended cDNAs, or fragments thereof are cloned into expression vectors such as those described in Example 30. The proteins are purified by size, charge, immunochromatography or other techniques familiar to those skilled in the art. Following purification, the proteins are labeled using techniques known to those
10 skilled in the art. The labeled proteins are incubated with cells or cell lines derived from a variety of organs or tissues to allow the proteins to bind to any receptor present on the cell surface. Following the incubation, the cells are washed to remove non-specifically bound protein. The labeled proteins are detected by autoradiography. Alternatively, unlabeled proteins may be incubated with the cells and detected with antibodies having a detectable
15 label, such as a fluorescent molecule, attached thereto.

Specificity of cell surface binding may be analyzed by conducting a competition analysis in which various amounts of unlabeled protein are incubated along with the labeled protein. The amount of labeled protein bound to the cell surface decreases as the amount of competitive unlabeled protein increases. As a control, various amounts of an unlabeled
20 protein unrelated to the labeled protein is included in some binding reactions. The amount of labeled protein bound to the cell surface does not decrease in binding reactions containing increasing amounts of unrelated unlabeled protein, indicating that the protein encoded by the cDNA binds specifically to the cell surface.

25 As discussed above, secreted proteins have been shown to have a number of important physiological effects and, consequently, represent a valuable therapeutic resource. The secreted proteins encoded by the extended cDNAs or portions thereof made according to Examples 27-29 may be evaluated to determine their physiological activities as described below.

EXAMPLE 32

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Cytokine,
Cell Proliferation or Cell Differentiation Activity

As discussed above, secreted proteins may act as cytokines or may affect cellular
5 proliferation or differentiation. Many protein factors discovered to date, including all known
cytokines, have exhibited activity in one or more factor dependent cell proliferation assays,
and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a
protein encoded by the extended cDNAs is evidenced by any one of a number of routine
factor dependent cell proliferation assays for cell lines including, without limitation, 32D,
10 DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M⁺ (preB M⁺), 2E8, RB5, DA1, 123, T1165,
HT2, CTLL2, TF-1, Mo7c and CMK. The proteins encoded by the above extended cDNAs
or portions thereof may be evaluated for their ability to regulate T cell or thymocyte
proliferation in assays such as those described above or in the following references, which are
incorporated herein by reference: *Current Protocols in Immunology*, Ed. by Coligan *et al.*,
15 Greene Publishing Associates and Wiley-Interscience; Takai *et al.* *J. Immunol.* 137:3494-
3500, 1986.; Bertagnolli *et al.*, *J. Immunol.* 145:1706-1712, 1990.; Bertagnolli *et al.*, *Cell.*
Immunol. 133:327-341, 1991; Bertagnolli, *et al.*, *J. Immunol.* 149:3778-3783, 1992;
Bowman *et al.*, *J. Immunol.* 152:1756-1761, 1994.

In addition, numerous assays for cytokine production and/or the proliferation of
20 spleen cells, lymph node cells and thymocytes are known. These include the techniques
disclosed in *Current Protocols in Immunology*, *supra* 1:3.12.1-3.12.14; and Schreiber In
Current Protocols in Immunology, *supra* 1 : 6.8.1-6.8.8.

The proteins encoded by the cDNAs may also be assayed for the ability to regulate
the proliferation and differentiation of hematopoietic or lymphopoietic cells. Many assays for
25 such activity are familiar to those skilled in the art, including the assays in the following
references, which are incorporated herein by reference: Bottomly *et al.*, In *Current Protocols*
in Immunology, *supra* 1 : 6.3.1-6.3.12.; deVries *et al.*, *J. Exp. Med.* 173:1205-1211, 1991;
Moreau *et al.*, *Nature* 36:690-692, 1988; Greenberger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*
80:2931-2938, 1983; Nordan, R., In *Current Protocols in Immunology*, *supra* 1 : 6.6.1-
30 6.6.5; Smith *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; Bennett *et al.*, in

Current Protocols in Immunology supra 1 : 6.15.1; Ciarletta et al., In Current Protocols in Immunology. supra 1 : 6.13.1.

The proteins encoded by the cDNAs may also be assayed for their ability to regulate T-cell responses to antigens. Many assays for such activity are familiar to those skilled in the art, including the assays described in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function), Chapter 6 (Cytokines and Their Cellular Receptors) and Chapter 7, (Immunologic Studies in Humans) in *Current Protocols in Immunology supra*; Weinberger *et al.*, *Proc. Natl. Acad. Sci. USA* 77:6091-6095, 1980; Weinberger *et al.*, *Eur. J. Immun.* 11:405-411, 1981; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988.

Those proteins which exhibit cytokine, cell proliferation, or cell differentiation activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which induction of cell proliferation or differentiation is beneficial. Alternatively, as described in more detail below, genes encoding these proteins or nucleic acids regulating the expression of these proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 33

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Activity as Immune System Regulators

The proteins encoded by the cDNAs may also be evaluated for their effects as immune regulators. For example, the proteins may be evaluated for their activity to influence thymocyte or splenocyte cytotoxicity. Numerous assays for such activity are familiar to those skilled in the art including the assays described in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function 3.1-3.19) and Chapter 7 (Immunologic studies in Humans) in *Current Protocols in Immunology*, Coligan *et al.*, Eds, Greene Publishing Associates and Wiley-Interscience; Herrmann *et al.*, *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann *et al.*, *J. Immunol.* 128:1968-1974, 1982; Handa *et al.*, *J. Immunol.* 135:1564-1572, 1985; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988;

Bowman *et al.*, *J. Virology* 61:1992-1998; Bertagnolli *et al.*, *Cell. Immunol.* 133:327-341, 1991; Brown *et al.*, *J. Immunol.* 153:3079-3092, 1994.

5 The proteins encoded by the cDNAs may also be evaluated for their effects on T-cell dependent immunoglobulin responses and isotype switching. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Maliszewski, *J. Immunol.* 144:3028-3033, 1990; Mond *et al.* in *Current Protocols in Immunology*, 1 : 3.8.1-3.8.16, *supra*.

10 The proteins encoded by the cDNAs may also be evaluated for their effect on immune effector cells, including their effect on Th1 cells and cytotoxic lymphocytes. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function 3.1-3.19) and Chapter 7 (Immunologic Studies in Humans) in *Current Protocols in Immunology*, *supra*; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988; Bertagnolli *et al.*, *J. Immunol.* 149:3778-3783, 15 1992.

The proteins encoded by the cDNAs may also be evaluated for their effect on dendritic cell mediated activation of naive T-cells. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Guery *et al.*, *J. Immunol.* 134:536-544, 1995; 20 Inaba *et al.*, *J. Exp. Med.* 173:549-559, 1991; Macatonia *et al.*, *J. Immunol.* 154:5071-5079, 1995; Porgador *et al.*, *J. Exp. Med.* 182:255-260, 1995; Nair *et al.*, *J. Virol.* 67:4062-4069, 1993; Huang *et al.*, *Science* 264:961-965, 1994; Macatonia *et al.*, *J. Exp. Med.* 169:1255-1264, 1989; Bhardwaj *et al.*, *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba *et al.*, *J. Exp. Med.* 172:631-640, 1990.

25 The proteins encoded by the cDNAs may also be evaluated for their influence on the lifetime of lymphocytes. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Darzynkiewicz *et al.*, *Cytometry* 13:795-808, 1992; Gorczyca *et al.*, *Leukemia* 7:659-670, 1993; Gorczyca *et al.*, *Cancer Res.* 53:1945-1951, 1993; Itoh *et al.*, *Cell* 66:233-30 243, 1991; Zacharchuk, *J. Immunol.* 145:4037-4045, 1990; Zamai *et al.*, *Cytometry* 14:891-897, 1993; Gorczyca *et al.*, *Int. J. Oncol.* 1:639-648, 1992.

The proteins encoded by the cDNAs may also be evaluated for their influence on early steps of T-cell commitment and development. Numerous assays for such activity are familiar to those skilled in the art, including without limitation the assays disclosed in the following references, which are incorporated herein by references: Antica *et al.*, *Blood* 84:111-117, 1994; Fine *et al.*, *Cell. Immunol.* 155:111-122, 1994; Galy *et al.*, *Blood* 85:2770-2778, 1995; Toki *et al.*, *Proc. Nat. Acad Sci. USA* 88:7548-7551, 1991.

Those proteins which exhibit activity as immune system regulators activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of immune activity is beneficial. For example, the protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, *Leishmania* spp., *Plasmodium* and various fungal infections such as candidiasis. Of course, in this regard, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the treatment of cancer.

Alternatively, proteins encoded by extended cDNAs derived from the 5' ESTs of the present invention may be used in treatment of autoimmune disorders including, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses either up or down.

Down regulation may involve inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of
5 activated T-cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active non-antigen-specific process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and
10 persists after the end of exposure to the tolerizing agent. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions, such as, for example, B7 costimulation), e.g.,
15 preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the
20 transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation, can lead to the binding of the molecule to the natural
25 ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may
30 avoid the necessity of repeated administration of these blocking reagents. To achieve

sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans.

5 Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, *Science* 257:789-792, 1992 and Turka *et al.*, *Proc. Natl. Acad. Sci USA*, 89:11102-11105, 1992. In addition, murine models of GVHD (see Paul ed., *Fundamental*
10 *Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and
15 autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor/ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which potentially involved in the disease process.

20 Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/pr/pr mice or
25 NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in OD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., *supra*, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may involve either enhancing an existing immune response or eliciting an
30 initial immune response as shown by the following examples. For instance, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases

of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory form of B lymphocyte antigens systemically.

Alternatively, antiviral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention or together with a stimulatory form of a soluble peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention and reintroducing the *in vitro* primed T cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to T cells *in vivo*, thereby activating the T cells.

In another application, upregulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules can be transfected with nucleic acids encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I α chain and β_2 microglobulin or an MHC class II α chain and an MHC class II β chain to thereby express MHC class I or MHC class II proteins on the cell surface, respectively. Expression of the appropriate MHC class I or class II

molecules in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA
5 encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject. Alternatively, as described in more detail below, genes encoding these immune system regulator proteins or nucleic acids regulating the expression of
10 such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 34

Assaying the Proteins Expressed from Extended cDNAs 15 or Portions Thereof for Hematopoiesis Regulating Activity

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their hematopoiesis regulating activity. For example, the effect of the proteins on embryonic stem cell differentiation may be evaluated. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following
20 references, which are incorporated herein by reference: Johansson *et al.* *Cell. Biol.* 15:141-151, 1995; Keller *et al.*, *Mol. Cell. Biol.* 13:473-486, 1993; McClanahan *et al.*, *Blood* 81:2903-2915, 1993.

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their influence on the lifetime of stem cells and stem cell differentiation.
25 Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Freshney, Methylcellulose Colony Forming Assays, in *Culture of Hematopoietic Cells.*, Freshney, *et al.* Eds. pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama *et al.*, *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; McNiece and Briddell, in *Culture of Hematopoietic Cells*,
30 *supra*; Neben *et al.*, *Exp. Hematol.* 22:353-359, 1994; Ploemacher and Cobblestone In

*Culture of Hematopoietic Cells, supra*1-21, Spooncer *et al*, in *Culture of Hematopoietic Cells, supra*163-179 and Sutherland in *Culture of Hematopoietic Cells, supra*. 139-162.

Those proteins which exhibit hematopoiesis regulatory activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of hematopoiesis is beneficial, such as in the treatment of myeloid or lymphoid cell deficiencies. Involvement in regulating hematopoiesis is indicated even by marginal biological activity in support of colony forming cells or of factor-dependent cell lines. For example, proteins supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, indicates utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells. Proteins supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (*i.e.*, traditional CSF activity) may be useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelosuppression. Proteins supporting the growth and proliferation of megakaryocytes and consequently of platelets allows prevention or treatment of various platelet disorders such as thrombocytopenia, and generally may be used in place of or complementary to platelet transfusions. Proteins supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells may therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in vivo* or *ex vivo* (*i.e.*, in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy. Alternatively, as described in more detail below, genes encoding hematopoiesis regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 35

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof
for Regulation of Tissue Growth

The proteins encoded by the extended cDNAs or portions thereof may also be
5 evaluated for their effect on tissue growth. Numerous assays for such activity are familiar to
those skilled in the art, including the assays disclosed in International Patent Publication No.
WO95/16035, International Patent Publication No. WO95/05846 and International Patent
Publication No. WO91/07491, which are incorporated herein by reference.

Assays for wound healing activity include, without limitation, those described in:
10 Winter, *Epidermal Wound Healing*, pps. 71-112, Maibach and Rovee, eds., Year Book
Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, *J. Invest. Dermatol.*
71:382-84, 1978, which are incorporated herein by reference.

Those proteins which are involved in the regulation of tissue growth may then be
formulated as pharmaceuticals and used to treat clinical conditions in which regulation of
15 tissue growth is beneficial. For example, a protein encoded by extended cDNAs derived from
the 5' ESTs of the present invention also may have utility in compositions used for bone,
cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound
healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present
20 invention, which induces cartilage and/or bone growth in circumstances where bone is not
normally formed, has application in the healing of bone fractures and cartilage damage or
defects in humans and other animals. Such a preparation employing a protein of the invention
may have prophylactic use in closed as well as open fracture reduction and also in the
improved fixation of artificial joints. *De novo* bone synthesis induced by an osteogenic agent
25 contributes to the repair of congenital, trauma induced, or oncologic resection induced
craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease,
and in other tooth repair processes. Such agents may provide an environment to attract bone-
forming cells, stimulate growth of bone-forming cells or induce differentiation of bone-
30 forming cell progenitors. A protein of the invention may also be useful in the treatment of
osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or

by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein encoded by extended cDNAs derived from the 5' ESTs of the present invention is tendon/ligament formation. A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition encoded by extended cDNAs derived from the 5' ESTs of the present invention contributes to the repair of tendon or ligaments defects of congenital, traumatic or other origin and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions encoded by extended cDNAs derived from the 5' ESTs of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.*, for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and

Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium) muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to generate. A protein of the invention may also exhibit angiogenic activity.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

Alternatively, as described in more detail below, genes encoding tissue growth regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 36

Assaying the Proteins Expressed from Extended cDNAs or Portions
Thereof for Regulation of Reproductive Hormones

The proteins encoded by the extended cDNAs or portions thereof may also be
5 evaluated for their ability to regulate reproductive hormones, such as follicle stimulating
hormone. Numerous assays for such activity are familiar to those skilled in the art, including
the assays disclosed in the following references, which are incorporated herein by reference:
Vale *et al.*, *Endocrinol.* **91**:562-572, 1972; Ling *et al.*, *Nature* **321**:779-782, 1986; Vale *et*
al., *Nature* **321**:776-779, 1986; Mason *et al.*, *Nature* **318**:659-663, 1985; Forage *et al.*,
10 *Proc. Natl. Acad. Sci. USA* **83**:3091-3095, 1986, Chapter 6.12 in *Current Protocols in*
Immunology, Coligan *et al.* Eds. Greene Publishing Associates and Wiley-Interscience ; Taub
et al., *J. Clin. Invest.* **95**:1370-1376, 1995; Lind *et al.*, *APMIS* **103**:140-146, 1995; Muller *et*
al., *Eur. J. Immunol.* **25**:1744-1748; Gruber *et al.*, *J. Immunol.* **152**:5860-5867, 1994;
Johnston *et al.*, *J Immunol.* **153**:1762-1768, 1994.

15 Those proteins which exhibit activity as reproductive hormones or regulators of cell
movement may then be formulated as pharmaceuticals and used to treat clinical conditions in
which regulation of reproductive hormones are beneficial. For example, a protein encoded by
extended cDNAs derived from the 5' ESTs of the present invention may also exhibit activin-
or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of
20 follicle stimulating hormone (FSH), while activins are characterized by their ability to
stimulate the release of FSH. Thus, a protein encoded by extended cDNAs derived from the
5' ESTs of the present invention, alone or in heterodimers with a member of the inhibin α
family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in
female mammals and decrease spermatogenesis in male mammals. Administration of
25 sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the
protein of the invention, as a homodimer or as a heterodimer with other protein subunits of
the inhibin-B group, may be useful as a fertility inducing therapeutic, based upon the ability of
activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for
example, United States Patent 4,798,885, the disclosure of which is incorporated herein by
30 reference. A protein of the invention may also be useful for advancement of the onset of

fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

Alternatively, as described in more detail below, genes encoding reproductive hormone regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 37

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Chemotactic/Chemokinetic Activity

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for chemotactic/chemokinetic activity. For example, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of

cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: *Current Protocols in Immunology*, Ed by Coligan, Kruisbeek, Margulies, Shevach and Strober, Pub. Greene Publishing Associates and Wiley-Interscience, Chapter 6.12: 6.12.1-6.12.28; Taub *et al.*, *J. Clin. Invest.* 95:1370-1376, 1995; Lind *et al.*, *APMIS* 103:140-146, 1995; Mueller *et al.*, *Eur. J. Immunol.* 25:1744-1748; Gruber *et al.*, *J. Immunol.* 152:5860-5867, 1994; Johnston *et al.* *J. Immunol.*, 153:1762-1768, 1994.

10

EXAMPLE 38

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Regulation of Blood Clotting

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their effects on blood clotting. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Linet *et al.*, *J. Clin. Pharmacol.* 26:131-140, 1986; Burdick *et al.*, *Thrombosis Res.* 45:413-419, 1987; Humphrey *et al.*, *Fibrinolysis* 5:71-79, 1991; Schaub, *Prostaglandins* 35:467-474, 1988.

Those proteins which are involved in the regulation of blood clotting may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of blood clotting is beneficial. For example, a protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulations disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as infarction of cardiac and central nervous system vessels (e.g., stroke)). Alternatively, as described in more detail below, genes encoding blood clotting activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 39

Assaying the Proteins Expressed from Extended cDNAs or
Portions Thereof for Involvement in Receptor/Ligand Interactions

The proteins encoded by the extended cDNAs or a portion thereof may also be
5 evaluated for their involvement in receptor/ligand interactions. Numerous assays for such
involvement are familiar to those skilled in the art, including the assays disclosed in the
following references, which are incorporated herein by reference: Chapter 7. 7.28.1-7.28.22
in *Current Protocols in Immunology*, Coligan *et al.* Eds. Greene Publishing Associates and
Wiley-Interscience; Takai *et al.*, *Proc. Natl. Acad. Sci. USA* 84:6864-6868, 1987; Bierer *et*
10 *al.*, *J. Exp. Med.* 168:1145-1156, 1988; Rosenstein *et al.*, *J. Exp. Med.* 169:149-160, 1989;
Stoltenborg *et al.*, *J. Immunol. Methods* 175:59-68, 1994; Stitt *et al.*, *Cell* 80:661-670, 1995;
Gyuris *et al.*, *Cell* 75:791-803, 1993.

For example, the proteins encoded by extended cDNAs derived from the 5' ESTs of
the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors
15 or agonists of receptor/ligand interactions. Examples of such receptors and ligands include,
without limitation, cytokine receptors and their ligands, receptor kinases and their ligands,
receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their
ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins
and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen
20 recognition and development of cellular and humoral immune responses). Receptors and
ligands are also useful for screening of potential peptide or small molecule inhibitors of the
relevant receptor/ligand interaction. A protein encoded by extended cDNAs derived from the
5' ESTs of the present invention (including, without limitation, fragments of receptors and
ligands) may themselves be useful as inhibitors of receptor/ligand interactions. Alternatively,
25 as described in more detail below, genes encoding proteins involved in receptor/ligand
interactions or nucleic acids regulating the expression of such proteins may be introduced into
appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 40

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof
for Anti-Inflammatory Activity

The proteins encoded by the extended cDNAs or a portion thereof may also be
5 evaluated for anti-inflammatory activity. The anti-inflammatory activity may be achieved by
providing a stimulus to cells involved in the inflammatory response, by inhibiting or
promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or
promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting
cell extravasation, or by stimulating or suppressing production of other factors which more
10 directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can
be used to treat inflammatory conditions including chronic or acute conditions, including
without limitation inflammation associated with infection (such as septic shock, sepsis or
systemic inflammatory response syndrome), ischemia-reperfusion injury, endotoxin lethality,
arthritis, complement-mediated hyperacute rejection, nephritis, cytokine- or chemokine-
15 induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over
production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to
treat anaphylaxis and hypersensitivity to an antigenic substance or material. Alternatively, as
described in more detail below, genes encoding anti-inflammatory activity proteins or nucleic
acids regulating the expression of such proteins may be introduced into appropriate host cells
20 to increase or decrease the expression of the proteins as desired.

EXAMPLE 41

Assaying the Proteins Expressed from Extended cDNAs or
Portions Thereof for Tumor Inhibition Activity

25 The proteins encoded by the extended cDNAs or a portion thereof may also be
evaluated for tumor inhibition activity. In addition to the activities described above for
immunological treatment or prevention of tumors, a protein of the invention may exhibit other
anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for
example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor
30 tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor
growth (such as, for example, by inhibiting angiogenesis), by causing production of other

factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth. Alternatively, as described in more detail below, genes tumor inhibition activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to
5 increase or decrease the expression of the proteins as desired.

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting
10 (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism,
15 processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and
20 growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune
25 response against such protein or another material or entity which is cross-reactive with such protein. Alternatively, as described in more detail below, genes encoding proteins involved in any of the above mentioned activities or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

EXAMPLE 42**Identification of Proteins which Interact with
Polypeptides Encoded by Extended cDNAs**

Proteins which interact with the polypeptides encoded by cDNAs derived from the 5' ESTs or fragments thereof, such as receptor proteins, may be identified using two hybrid systems such as the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech). As described in the manual accompanying the kit which is incorporated herein by reference, the the cDNAs derived from 5' ESTs, or fragments thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA binding domain of the yeast transcriptional activator GAL4. cDNAs in a cDNA library which encode proteins which might interact with the polypeptides encoded by the extended cDNAs or portions thereof are inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into yeast and the yeast are plated on selection medium which selects for expression of selectable markers on each of the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay contain plasmids encoding proteins which interact with the polypeptide encoded by the extended cDNAs or portions thereof.

Alternatively, the system described in Lustig *et al.*, *Methods in Enzymology* 283: 83-99, 1997, and in U.S. Patent No. 5,654,150, the disclosure of which is incorporated herein by reference, may be used for identifying molecules which interact with the polypeptides encoded by extended cDNAs. In such systems, *in vitro* transcription reactions are performed on a pool of vectors containing extended cDNA inserts cloned downstream of a promoter which drives *in vitro* transcription. The resulting pools of mRNAs are introduced into *Xenopus laevis* oocytes. The oocytes are then assayed for a desired activity.

Alternatively, the pooled *in vitro* transcription products produced as described above may be translated *in vitro*. The pooled *in vitro* translation products can be assayed for a desired activity or for interaction with a known polypeptide.

Proteins or other molecules interacting with polypeptides encoded by extended cDNAs can be found by a variety of additional techniques. In one method, affinity

columns containing the polypeptide encoded by the extended cDNA or a portion thereof can be constructed. In some versions, of this method the affinity column contains chimeric proteins in which the protein encoded by the extended cDNA or a portion thereof is fused to glutathione S-transferase. A mixture of cellular proteins or pool of expressed proteins as described above and is applied to the affinity column. Proteins interacting with the polypeptide attached to the column can then be isolated and analyzed on 2-D electrophoresis gel as described in Ramunsen *et al.*, *Electrophoresis* 18:588-598, 1997, the disclosure of which is incorporated herein by reference. Alternatively, the proteins retained on the affinity column can be purified by electrophoresis based methods and sequenced. The same method can be used to isolate antibodies, to screen phage display products, or to screen phage display human antibodies.

Proteins interacting with polypeptides encoded by extended cDNAs or portions thereof can also be screened by using an Optical Biosensor as described in Edwards and Leatherbarrow, *Analytical Biochemistry* 246:1-6, 1997, the disclosure of which is incorporated herein by reference. The main advantage of the method is that it allows the determination of the association rate between the protein and other interacting molecules. Thus, it is possible to specifically select interacting molecules with a high or low association rate. Typically a target molecule is linked to the sensor surface (through a carboxymethyl dextran matrix) and a sample of test molecules is placed in contact with the target molecules. The binding of a test molecule to the target molecule causes a change in the refractive index and/ or thickness. This change is detected by the Biosensor provided it occurs in the evanescent field (which extend a few hundred nanometers from the sensor surface). In these screening assays, the target molecule can be one of the polypeptides encoded by extended cDNAs or a portion thereof and the test sample can be a collection of proteins extracted from tissues or cells, a pool of expressed proteins, combinatorial peptide and/ or chemical libraries, or phage displayed peptides. The tissues or cells from which the test proteins are extracted can originate from any species.

In other methods, a target protein is immobilized and the test population is a collection of unique polypeptides encoded by the extended cDNAs or portions thereof.

To study the interaction of the proteins encoded by the extended cDNAs or portions thereof with drugs, the microdialysis coupled to HPLC method described by Wang *et al.*, *Chromatographia* 44:205-208, 1997 or the affinity capillary electrophoresis method described by Busch *et al.*, *J. Chromatogr.* 777:311-328, 1997, the disclosures of which are incorporated herein by reference can be used.

It will be appreciated by those skilled in the art that the proteins expressed from the extended cDNAs or portions may be assayed for numerous activities in addition to those specifically enumerated above. For example, the expressed proteins may be evaluated for applications involving control and regulation of inflammation, tumor proliferation or metastasis, infection, or other clinical conditions. In addition, the proteins expressed from the extended cDNAs or portions thereof may be useful as nutritional agents or cosmetic agents.

The proteins expressed from the cDNAs or portions thereof may be used to generate antibodies capable of specifically binding to the expressed protein or fragments thereof as described in Example 40 below. The antibodies may be capable of binding a full length protein encoded by a cDNA derived from a 5' EST, a mature protein (*i.e.* the protein generated by cleavage of the signal peptide) encoded by a cDNA derived from a 5' EST, or a signal peptide encoded by a cDNA derived from a 5' EST. Alternatively, the antibodies may be capable of binding fragments of at least 10 amino acids of the proteins encoded by the above cDNAs. In some embodiments, the antibodies may be capable of binding fragments of at least 15 amino acids of the proteins encoded by the above cDNAs. In other embodiments, the antibodies may be capable of binding fragments of at least 25 amino acids of the proteins expressed from the extended cDNAs which comprise at least 25 amino acids of the proteins encoded by the above cDNAs. In further embodiments, the antibodies may be capable of binding fragments of at least 40 amino acids of the proteins encoded by the above cDNAs.

EXAMPLE 43

Production of an Antibody to a Human Protein

Substantially pure protein or polypeptide is isolated from the transfected or transformed cells as described in Example 30. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the

level of a few $\mu\text{g/ml}$. Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

1. Monoclonal Antibody Production by Hybridoma Fusion

5 Monoclonal antibody to epitopes of any of the peptides identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler, and Milstein, *Nature* 256:495, 1975 or derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein or peptides derived therefrom over a period of a few weeks. The mouse is then sacrificed,
10 and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing
15 clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, *Meth. Enzymol.* 70:419, 1980, the disclosure of which is incorporated herein by reference and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal
20 antibody production are described in Davis *et al.* in *Basic Methods in Molecular Biology* Elsevier, New York. Section 21-2, the disclosure of which is incorporated herein by reference.

2. Polyclonal Antibody Production by Immunization

25 Polyclonal antiserum containing antibodies to heterogenous epitopes of a single protein can be prepared by immunizing suitable animals with the expressed protein or peptides derived therefrom, which can be unmodified or modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less
30 immunogenic than others and may require the use of carriers and adjuvant. Also, host animals response vary depending on site of inoculations and doses, with both inadequate or

excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis. *et al.*, *J. Clin. Endocrinol. Metab.* 33:988-991 (1971), the disclosure of which is incorporated herein by reference.

5 Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, *et al.*, Chap. 19 in: *Handbook of Experimental Immunology* D. Wier (ed) Blackwell (1973), the disclosure of which is incorporated herein by reference. Plateau
10 concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12 μ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, D., Chap. 42 in: *Manual of Clinical Immunology*, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980), the disclosure of which is incorporated herein by reference.

15 Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of
20 the protein in the body.

V. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof as Reagents

25 The 5' ESTs of the present invention (or cDNAs or genomic DNAs obtainable therefrom) may be used as reagents in isolation procedures, diagnostic assays, and forensic procedures. For example, sequences from the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be detectably labeled and used as probes to isolate other sequences capable of hybridizing to them. In addition, sequences from 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be used to design PCR primers to
30 be used in isolation, diagnostic, or forensic procedures.

1. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof in Isolation,
Diagnostic and Forensic Procedures

EXAMPLE 44

Preparation of PCR Primers and Amplification of DNA

5 The 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) may be used to prepare PCR primers for a variety of applications, including isolation procedures for cloning nucleic acids capable of hybridizing to such sequences, diagnostic techniques and forensic techniques. The PCR primers are at least 10 bases, and preferably at least 12, 15, or 17 bases in length. More preferably, the PCR primers are at least 20-30 bases in length. In
10 some embodiments, the PCR primers may be more than 30 bases in length. It is preferred that the primer pairs have approximately the same G/C ratio, so that melting temperatures are approximately the same. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see *Molecular Cloning to Genetic Engineering*, White Ed. in *Methods in Molecular Biology* 67: Humana Press, Totowa 1997, the disclosure of which
15 is incorporated herein by reference. In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample.
20 The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites.

EXAMPLE 45

Use of 5'ESTs as Probes

25 Probes derived from 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom), including full length cDNAs or genomic sequences, may be labeled with detectable labels familiar to those skilled in the art, including radioisotopes and non-radioactive labels, to provide a detectable probe. The detectable probe may be single stranded or double stranded
30 and may be made using techniques known in the art, including *in vitro* transcription, nick translation, or kinase reactions. A nucleic acid sample containing a sequence capable of

hybridizing to the labeled probe is contacted with the labeled probe. If the nucleic acid in the sample is double stranded, it may be denatured prior to contacting the probe. In some applications, the nucleic acid sample may be immobilized on a surface such as a nitrocellulose or nylon membrane. The nucleic acid sample may comprise nucleic acids obtained from a variety of sources, including genomic DNA, cDNA libraries, RNA, or tissue samples.

Procedures used to detect the presence of nucleic acids capable of hybridizing to the detectable probe include well known techniques such as Southern blotting, Northern blotting, dot blotting, colony hybridization, and plaque hybridization. In some applications, the nucleic acid capable of hybridizing to the labeled probe may be cloned into vectors such as expression vectors, sequencing vectors, or *in vitro* transcription vectors to facilitate the characterization and expression of the hybridizing nucleic acids in the sample. For example, such techniques may be used to isolate and clone sequences in a genomic library or cDNA library which are capable of hybridizing to the detectable probe as described in Example 30 above.

PCR primers made as described in Example 44 above may be used in forensic analyses, such as the DNA fingerprinting techniques described in Examples 46-50 below. Such analyses may utilize detectable probes or primers based on the sequences of the the 5' ESTs or of cDNAs or genomic DNAs isolated using the 5' ESTs.

EXAMPLE 46

Forensic Matching by DNA Sequencing

In one exemplary method, DNA samples are isolated from forensic specimens of, for example, hair, semen, blood or skin cells by conventional methods. A panel of PCR primers based on a number of the 5' ESTs of Example 25, or cDNAs or genomic DNAs isolated therefrom as described above, is then utilized in accordance with Example 44 to amplify DNA of approximately 100-200 bases in length from the forensic specimen. Corresponding sequences are obtained from a test subject. Each of these identification DNAs is then sequenced using standard techniques, and a simple database comparison determines the differences, if any, between the sequences from the subject and those from the sample. Statistically significant differences between the suspect's DNA sequences and those from the sample conclusively prove a lack of identity. This lack of identity can be proven, for example, with only one sequence. Identity, on the other hand, should be demonstrated with a large

number of sequences, all matching. Preferably, a minimum of 50 statistically identical sequences of 100 bases in length are used to prove identity between the suspect and the sample.

5

EXAMPLE 47

Positive Identification by DNA Sequencing

The technique outlined in the previous example may also be used on a larger scale to provide a unique fingerprint-type identification of any individual. In this technique, primers are prepared from a large number of 5'EST sequences from Example 25, or cDNA or genomic DNA sequences obtainable therefrom. Preferably, 20 to 50 different primers are used. These primers are used to obtain a corresponding number of PCR-generated DNA segments from the individual in question in accordance with Example 44. Each of these DNA segments is sequenced, using the methods set forth in Example 46. The database of sequences generated through this procedure uniquely identifies the individual from whom the sequences were obtained. The same panel of primers may then be used at any later time to absolutely correlate tissue or other biological specimen with that individual.

15

EXAMPLE 48

Southern Blot Forensic Identification

The procedure of Example 47 is repeated to obtain a panel of at least 10 amplified sequences from an individual and a specimen. Preferably, the panel contains at least 50 amplified sequences. More preferably, the panel contains 100 amplified sequences. In some embodiments, the panel contains 200 amplified sequences. This PCR-generated DNA is then digested with one or a combination of, preferably, four base specific restriction enzymes. Such enzymes are commercially available and known to those of skill in the art. After digestion, the resultant gene fragments are size separated in multiple duplicate wells on an agarose gel and transferred to nitrocellulose using Southern blotting techniques well known to those with skill in the art. For a review of Southern blotting see Davis *et al.* (Basic Methods in Molecular Biology, 1986, Elsevier Press. pp 62-65), the disclosure of which is incorporated herein by reference.

20

25

30

A panel of probes based on the sequences of 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom), or fragments thereof of at least 10 bases, are radioactively or colorimetrically labeled using methods known in the art, such as nick translation or end labeling, and hybridized to the Southern blot using techniques known in the art (Davis *et al.*,
5 *supra*). Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom). More preferably, the probe comprises at least 20-30 consecutive nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom). In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom).

10 Preferably, at least 5 to 10 of these labeled probes are used, and more preferably at least about 20 or 30 are used to provide a unique pattern. The resultant bands appearing from the hybridization of a large sample of 5' EST (or cDNAs or genomic DNAs obtainable therefrom) will be a unique identifier. Since the restriction enzyme cleavage will be different for every individual, the band pattern on the Southern blot will also be unique. Increasing the
15 number of 5' EST (or cDNAs or genomic DNAs obtainable therefrom) probes will provide a statistically higher level of confidence in the identification since there will be an increased number of sets of bands used for identification.

EXAMPLE 49

20 Dot Blot Identification Procedure

Another technique for identifying individuals using the 5' EST sequences disclosed herein utilizes a dot blot hybridization technique.

Genomic DNA is isolated from nuclei of subject to be identified. Oligonucleotide probes of approximately 30 bp in length are synthesized that correspond to at least 10,
25 preferably 50 sequences from the 5' ESTs or cDNAs or genomic DNAs obtainable therefrom. The probes are used to hybridize to the genomic DNA through conditions known to those in the art. The oligonucleotides are end labeled with P³² using polynucleotide kinase (Pharmacia). Dot Blots are created by spotting the genomic DNA onto nitrocellulose or the like using a vacuum dot blot manifold (BioRad, Richmond California). The nitrocellulose
30 filter containing the genomic sequences is baked or UV linked to the filter, prehybridized and hybridized with labeled probe using techniques known in the art (Davis *et al.*, *supra*). The ³²P

labeled DNA fragments are sequentially hybridized with successively stringent conditions to detect minimal differences between the 30 bp sequence and the DNA. Tetramethylammonium chloride is useful for identifying clones containing small numbers of nucleotide mismatches (Wood *et al.*, *Proc. Natl. Acad. Sci. USA* **82**(6):1585-1588, 1985) which is hereby incorporated by reference. A unique pattern of dots distinguishes one individual from another individual.

5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) or oligonucleotides containing at least 10 consecutive bases from these sequences can be used as probes in the following alternative fingerprinting technique. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom). More preferably, the probe comprises at least 20-30 consecutive nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom). In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom).

Preferably, a plurality of probes having sequences from different genes are used in the alternative fingerprinting technique. Example 50 below provides a representative alternative fingerprinting procedure in which the probes are derived from 5'EST.

EXAMPLE 50

Alternative "Fingerprint" Identification Technique

20-mer oligonucleotides are prepared from a large number, e.g. 50, 100, or 200, of 5'EST using commercially available oligonucleotide services such as Genset, Paris, France. Cell samples from the test subject are processed for DNA using techniques well known to those with skill in the art. The nucleic acid is digested with restriction enzymes such as EcoRI and XbaI. Following digestion, samples are applied to wells for electrophoresis. The procedure, as known in the art, may be modified to accommodate polyacrylamide electrophoresis, however in this example, samples containing 5 ug of DNA are loaded into wells and separated on 0.8% agarose gels. The gels are transferred onto nitrocellulose using standard Southern blotting techniques.

10 ng of each of the oligonucleotides are pooled and end-labeled with ^{32}P . The nitrocellulose is prehybridized with blocking solution and hybridized with the labeled probes.

Following hybridization and washing, the nitrocellulose filter is exposed to X-Omat AR X-ray film. The resulting hybridization pattern will be unique for each individual.

It is additionally contemplated within this example that the number of probe sequences used can be varied for additional accuracy or clarity.

5

The proteins encoded by the extended cDNAs may also be used to generate antibodies as explained in Examples 30 and 43 in order to identify the tissue type or cell species from which a sample is derived as described in example 51.

10

EXAMPLE 51

Identification of Tissue Types or Cell Species by Means of Labeled Tissue Specific Antibodies

Identification of specific tissues is accomplished by the visualization of tissue specific antigens by means of antibody preparations according to Examples 30 and 43 which are
15 conjugated, directly or indirectly to a detectable marker. Selected labeled antibody species bind to their specific antigen binding partner in tissue sections, cell suspensions, or in extracts of soluble proteins from a tissue sample to provide a pattern for qualitative or semi-qualitative interpretation.

Antisera for these procedures must have a potency exceeding that of the native
20 preparation, and for that reason, antibodies are concentrated to a mg/ml level by isolation of the gamma globulin fraction, for example, by ion-exchange chromatography or by ammonium sulfate fractionation. Also, to provide the most specific antisera, unwanted antibodies, for example to common proteins, must be removed from the gamma globulin fraction, for example by means of insoluble immunoabsorbents, before the antibodies are
25 labeled with the marker. Either monoclonal or heterologous antisera is suitable for either procedure.

A. Immunohistochemical techniques

Purified, high-titer antibodies, prepared as described above, are conjugated to a detectable marker, as described, for example, by Fudenberg, Chap. 26 in: *Basic and Clinical*
30 *Immunology*, 3rd Ed. Lange, Los Altos, California, 1980, or Rose, *et al.*, Chap. 12 in:

Methods in Immunodiagnosis, 2d Ed. John Wiley and Sons, New York (1980), the disclosures of which are incorporated herein by reference.

A fluorescent marker, either fluorescein or rhodamine, is preferred, but antibodies can also be labeled with an enzyme that supports a color producing reaction with a substrate, such as horseradish peroxidase. Markers can be added to tissue-bound antibody in a second step, as described below. Alternatively, the specific antitissue antibodies can be labeled with ferritin or other electron dense particles, and localization of the ferritin coupled antigen-antibody complexes achieved by means of an electron microscope. In yet another approach, the antibodies are radiolabeled, with, for example ^{125}I , and detected by overlaying the antibody treated preparation with photographic emulsion.

Preparations to carry out the procedures can comprise monoclonal or polyclonal antibodies to a single protein or peptide identified as specific to a tissue type, for example, brain tissue, or antibody preparations to several antigenically distinct tissue specific antigens can be used in panels, independently or in mixtures, as required.

Tissue sections and cell suspensions are prepared for immunohistochemical examination according to common histological techniques. Multiple cryostat sections (about 4 μm , unfixed) of the unknown tissue and known control, are mounted and each slide covered with different dilutions of the antibody preparation. Sections of known and unknown tissues should also be treated with preparations to provide a positive control, a negative control, for example, pre-immune sera, and a control for non-specific staining, for example, buffer.

Treated sections are incubated in a humid chamber for 30 min at room temperature, rinsed, then washed in buffer for 30-45 min. Excess fluid is blotted away, and the marker developed.

If the tissue specific antibody was not labeled in the first incubation, it can be labeled at this time in a second antibody-antibody reaction, for example, by adding fluorescein- or enzyme-conjugated antibody against the immunoglobulin class of the antiserum-producing species, for example, fluorescein labeled antibody to mouse IgG. Such labeled sera are commercially available.

The antigen found in the tissues by the above procedure can be quantified by measuring the intensity of color or fluorescence on the tissue section, and calibrating that signal using appropriate standards.

B. Identification of tissue specific soluble proteins

5 The visualization of tissue specific proteins and identification of unknown tissues from that procedure is carried out using the labeled antibody reagents and detection strategy as described for immunohistochemistry; however the sample is prepared according to an electrophoretic technique to distribute the proteins extracted from the tissue in an orderly array on the basis of molecular weight for detection.

10 A tissue sample is homogenized using a Virtis apparatus; cell suspensions are disrupted by Dounce homogenization or osmotic lysis, using detergents in either case as required to disrupt cell membranes, as is the practice in the art. Insoluble cell components such as nuclei, microsomes, and membrane fragments are removed by ultracentrifugation, and the soluble protein-containing fraction concentrated if necessary and reserved for analysis.

15 A sample of the soluble protein solution is resolved into individual protein species by conventional SDS polyacrylamide electrophoresis as described, for example, by Davis, *et al.*, Section 19-2 in: *Basic Methods in Molecular Biology*, Leder ed., Elsevier, New York, 1986, the disclosure of which is incorporated herein by reference, using a range of amounts of polyacrylamide in a set of gels to resolve the entire molecular weight range of proteins to be
20 detected in the sample. A size marker is run in parallel for purposes of estimating molecular weights of the constituent proteins. Sample size for analysis is a convenient volume of from 5 to 55 μ l, and containing from about 1 to 100 μ g protein. An aliquot of each of the resolved proteins is transferred by blotting to a nitrocellulose filter paper, a process that maintains the pattern of resolution. Multiple copies are prepared. The procedure, known as Western Blot
25 Analysis, is well described in Davis, L. *et al.*, *supra* Section 19-3. One set of nitrocellulose blots is stained with Coomassie blue dye to visualize the entire set of proteins for comparison with the antibody bound proteins. The remaining nitrocellulose filters are then incubated with a solution of one or more specific antisera to tissue specific proteins prepared as described in Examples 30 and 43. In this procedure, as in procedure A above, appropriate positive and
30 negative sample and reagent controls are run.

In either procedure A or B, a detectable label can be attached to the primary tissue antigen-primary antibody complex according to various strategies and permutations thereof. In a straightforward approach, the primary specific antibody can be labeled; alternatively, the unlabeled complex can be bound by a labeled secondary anti-IgG antibody. In other approaches, either the primary or secondary antibody is conjugated to a biotin molecule, which can, in a subsequent step, bind an avidin conjugated marker. According to yet another strategy, enzyme labeled or radioactive protein A, which has the property of binding to any IgG, is bound in a final step to either the primary or secondary antibody.

The visualization of tissue specific antigen binding at levels above those seen in control tissues to one or more tissue specific antibodies, prepared from the gene sequences identified from extended cDNA sequences, can identify tissues of unknown origin, for example, forensic samples, or differentiated tumor tissue that has metastasized to foreign bodily sites.

In addition to their applications in forensics and identification, 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be mapped to their chromosomal locations. Example 52 below describes radiation hybrid (RH) mapping of human chromosomal regions using 5'ESTs. Example 53 below describes a representative procedure for mapping an 5' EST to its location on a human chromosome. Example 54 below describes mapping of 5' ESTs on metaphase chromosomes by Fluorescence In Situ Hybridization (FISH). Those skilled in the art will appreciate that the method of Examples 52-54 may also be used to map cDNAs or genomic DNAs obtainable from the 5' ESTs to their chromosomal locations.

2. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof in Chromosome Mapping

EXAMPLE 52

Radiation hybrid mapping of 5'ESTs to the human genome

Radiation hybrid (RH) mapping is a somatic cell genetic approach that can be used for high resolution mapping of the human genome. In this approach, cell lines containing one or more human chromosomes are lethally irradiated, breaking each chromosome into fragments whose size depends on the radiation dose. These fragments are rescued by fusion

with cultured rodent cells, yielding subclones containing different portions of the human genome. This technique is described by Benham *et al.*, *Genomics* 4:509-517, 1989; and Cox *et al.*, *Science* 250:245-250, 1990, the entire contents of which are hereby incorporated by reference. The random and independent nature of the subclones permits efficient mapping of
5 any human genome marker. Human DNA isolated from a panel of 80-100 cell lines provides a mapping reagent for ordering 5'EST. In this approach, the frequency of breakage between markers is used to measure distance, allowing construction of fine resolution maps as has been done using conventional ESTs (Schuler *et al.*, *Science* 274:540-546, 1996, hereby incorporated by reference).

10 RH mapping has been used to generate a high-resolution whole genome radiation hybrid map of human chromosome 17q22-q25.3 across the genes for growth hormone (GH) and thymidine kinase (TK) (Foster *et al.*, *Genomics* 33:185-192, 1996), the region surrounding the Gorlin syndrome gene (Obermayr *et al.*, *Eur. J. Hum. Genet.* 4:242-245, 1996), 60 loci covering the entire short arm of chromosome 12 (Raeymaekers *et al.*,
15 *Genomics* 29:170-178, 1995), the region of human chromosome 22 containing the neurofibromatosis type 2 locus (Frazer *et al.*, *Genomics* 14:574-584, 1992) and 13 loci on the long arm of chromosome 5 (Warrington *et al.*, *Genomics* 11:701-708, 1991).

EXAMPLE 53

20 Mapping of 5'ESTs to Human Chromosomes using PCR techniques

5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be assigned to human chromosomes using PCR based methodologies. In such approaches, oligonucleotide primer pairs are designed from the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) to minimize the chance of amplifying through an intron. Preferably, the
25 oligonucleotide primers are 18-23 bp in length and are designed for PCR amplification. The creation of PCR primers from known sequences is well known to those with skill in the art. For a review of PCR technology see Erlich in *PCR Technology, Principles and Applications for DNA Amplification*, Freeman and Co., New York, 1992, the disclosure of which is incorporated herein by reference.

30 The primers are used in polymerase chain reactions (PCR) to amplify templates from total human genomic DNA. PCR conditions are as follows: 60 ng of genomic DNA is used

as a template for PCR with 80 ng of each oligonucleotide primer, 0.6 unit of Taq polymerase, and 1 μ Ci of a 32 P-labeled deoxycytidine triphosphate. The PCR is performed in a microplate thermocycler (Techne) under the following conditions: 30 cycles of 94°C, 1.4 min; 55°C, 2 min; and 72°C, 2 min; with a final extension at 72°C for 10 min. The amplified products are analyzed on a 6% polyacrylamide sequencing gel and visualized by autoradiography. If the length of the resulting PCR product is identical to the distance between the ends of the primer sequences in the extended cDNA from which the primers are derived, then the PCR reaction is repeated with DNA templates from two panels of human-rodent somatic cell hybrids, BIOS PCRable DNA (BIOS Corporation) and NIGMS Human-Rodent Somatic Cell Hybrid Mapping Panel Number 1 (NIGMS, Camden, NJ).

PCR is used to screen a series of somatic cell hybrid cell lines containing defined sets of human chromosomes for the presence of a given 5' EST (or cDNA or genomic DNA obtainable therefrom). DNA is isolated from the somatic hybrids and used as starting templates for PCR reactions using the primer pairs from the 5' EST (or cDNA or genomic DNA obtainable therefrom). Only those somatic cell hybrids with chromosomes containing the human gene corresponding to the 5' EST (or cDNA or genomic DNA obtainable therefrom) will yield an amplified fragment. The 5' EST (or cDNA or genomic DNA obtainable therefrom) are assigned to a chromosome by analysis of the segregation pattern of PCR products from the somatic hybrid DNA templates. The single human chromosome present in all cell hybrids that give rise to an amplified fragment is the chromosome containing that 5'EST (or cDNA or genomic DNA obtainable therefrom). For a review of techniques and analysis of results from somatic cell gene mapping experiments, see Ledbetter *et al.*, *Genomics* 6:475-481, 1990, the disclosure of which is incorporated herein by reference.

25

EXAMPLE 54

Mapping of Extended 5' ESTs to Chromosomes Using Fluorescence *In Situ*

Hybridization

Fluorescence in situ hybridization allows the 5'EST (or cDNA or genomic DNA obtainable therefrom) to be mapped to a particular location on a given chromosome. The chromosomes to be used for fluorescence in situ hybridization techniques may be obtained from a variety of sources including cell cultures, tissues, or whole blood.

30

In a preferred embodiment, chromosomal localization of an 5'EST (or cDNA or genomic DNA obtainable therefrom) is obtained by FISH as described by Cherif *et al.* (*Proc. Natl. Acad. Sci. U.S.A.*, 87:6639-6643, 1990), the disclosure of which is incorporated herein by reference.. Metaphase chromosomes are prepared from phytohemagglutinin (PHA)-stimulated blood cell donors. PHA-stimulated lymphocytes from healthy males are cultured for 72 h in RPMI-1640 medium. For synchronization, methotrexate (10 μ M) is added for 17 h, followed by addition of 5-bromodeoxyuridine (5-BrdU, 0.1 mM) for 6 h. Colcemid (1 μ g/ml) is added for the last 15 min before harvesting the cells. Cells are collected, washed in RPMI, incubated with a hypotonic solution of KCl (75 mM) at 37°C for 15 min and fixed in three changes of methanol:acetic acid (3:1). The cell suspension is dropped onto a glass slide and air dried. The 5'EST (or cDNA or genomic DNA obtainable therefrom) is labeled with biotin-16 dUTP by nick translation according to the manufacturer's instructions (Bethesda Research Laboratories, Bethesda, MD), purified using a Sephadex G-50 column (Pharmacia, Upsala, Sweden) and precipitated. Just prior to hybridization, the DNA pellet is dissolved in hybridization buffer (50% formamide, 2 X SSC, 10% dextran sulfate, 1 mg/ml sonicated salmon sperm DNA, pH 7) and the probe is denatured at 70°C for 5-10 min.

Slides kept at -20°C are treated for 1 h at 37°C with RNase A (100 μ g/ml), rinsed three times in 2 X SSC and dehydrated in an ethanol series. Chromosome preparations are denatured in 70% formamide, 2 X SSC for 2 min at 70°C, then dehydrated at 4°C. The slides are treated with proteinase K (10 μ g/100 ml in 20 mM Tris-HCl, 2 mM CaCl₂) at 37°C for 8 min and dehydrated. The hybridization mixture containing the probe is placed on the slide, covered with a coverslip, sealed with rubber cement and incubated overnight in a humid chamber at 37°C. After hybridization and post-hybridization washes, the biotinylated probe is detected by avidin-FITC and amplified with additional layers of biotinylated goat anti-avidin and avidin-FITC. For chromosomal localization, fluorescent R-bands are obtained as previously described (Cherif *et al.*, *supra.*). The slides are observed under a LEICA fluorescence microscope (DMRXA). Chromosomes are counterstained with propidium iodide and the fluorescent signal of the probe appears as two symmetrical yellow-green spots on both chromatids of the fluorescent R-band chromosome (red). Thus, a particular 5'EST (or cDNA or genomic DNA obtainable therefrom) may be localized to a particular cytogenetic R-band on a given chromosome.

Once the 5'EST (or cDNA or genomic DNA obtainable therefrom) have been assigned to particular chromosomes using the techniques described in Examples 52-54 above, they may be utilized to construct a high resolution map of the chromosomes on which they are located or to identify the chromosomes in a sample.

5

EXAMPLE 55

Use of 5'EST to Construct or Expand Chromosome Maps

Chromosome mapping involves assigning a given unique sequence to a particular chromosome as described above. Once the unique sequence has been mapped to a given chromosome, it is ordered relative to other unique sequences located on the same chromosome. One approach to chromosome mapping utilizes a series of yeast artificial chromosomes (YACs) bearing several thousand long inserts derived from the chromosomes of the organism from which the extended cDNAs (or genomic DNAs obtainable therefrom) are obtained. This approach is described in Nagaraja *et al.*, *Genome Research* 7:210-222, 1997, the disclosure of which is incorporated herein by reference. Briefly, in this approach each chromosome is broken into overlapping pieces which are inserted into the YAC vector. The YAC inserts are screened using PCR or other methods to determine whether they include the 5'EST (or cDNA or genomic DNA obtainable therefrom) whose position is to be determined. Once an insert has been found which includes the 5'EST (or cDNA or genomic DNA obtainable therefrom), the insert can be analyzed by PCR or other methods to determine whether the insert also contains other sequences known to be on the chromosome or in the region from which the 5'EST (or cDNA or genomic DNA obtainable therefrom) was derived. This process can be repeated for each insert in the YAC library to determine the location of each of the extended cDNAs (or genomic DNAs obtainable therefrom) relative to one another and to other known chromosomal markers. In this way, a high resolution map of the distribution of numerous unique markers along each of the organisms chromosomes may be obtained.

As described in Example 56 below extended cDNAs (or genomic DNAs obtainable therefrom) may also be used to identify genes associated with a particular phenotype, such as hereditary disease or drug response.

3. Use of 5'ESTs or Sequences Obtained Therefrom or Fragments Thereof in Gene Identification

EXAMPLE 56

Identification of genes associated with hereditary diseases or drug response

- 5 This example illustrates an approach useful for the association of 5'ESTs (or cDNA or genomic DNA obtainable therefrom) with particular phenotypic characteristics. In this example, a particular 5'EST (or cDNA or genomic DNA obtainable therefrom) is used as a test probe to associate that 5'EST (or cDNA or genomic DNA obtainable therefrom) with a particular phenotypic characteristic.
- 10 5'ESTs (or cDNA or genomic DNA obtainable therefrom) are mapped to a particular location on a human chromosome using techniques such as those described in Examples 52 and 53 or other techniques known in the art. A search of Mendelian Inheritance in Man (McKusick in *Mendelian Inheritance in Man* (available on line through Johns Hopkins University Welch Medical Library) reveals the region of the human chromosome which
- 15 contains the 5'EST (or cDNA or genomic DNA obtainable therefrom) to be a very gene rich region containing several known genes and several diseases or phenotypes for which genes have not been identified. The gene corresponding to this 5'EST (or cDNA or genomic DNA obtainable therefrom) thus becomes an immediate candidate for each of these genetic diseases.
- 20 Cells from patients with these diseases or phenotypes are isolated and expanded in culture. PCR primers from the 5'EST (or cDNA or genomic DNA obtainable therefrom) are used to screen genomic DNA, mRNA or cDNA obtained from the patients. 5'ESTs (or cDNA or genomic DNA obtainable therefrom) that are not amplified in the patients can be positively associated with a particular disease by further
- 25 analysis. Alternatively, the PCR analysis may yield fragments of different lengths when the samples are derived from an individual having the phenotype associated with the disease than when the sample is derived from a healthy individual, indicating that the gene containing the 5'EST may be responsible for the genetic disease.

VI. Use of 5'EST (or cDNA or Genomic DNA Obtainable Therefrom) to Construct Vectors

The present 5'ESTs (or cDNA or genomic DNA obtainable therefrom) may also be used to construct secretion vectors capable of directing the secretion of the proteins encoded by genes therein. Such secretion vectors may facilitate the purification or enrichment of the proteins encoded by genes inserted therein by reducing the number of background proteins from which the desired protein must be purified or enriched. Exemplary secretion vectors are described in Example 57 below.

1. Construction of Secretion Vectors

EXAMPLE 57

Construction of Secretion Vectors

The secretion vectors include a promoter capable of directing gene expression in the host cell, tissue, or organism of interest. Such promoters include the Rous Sarcoma Virus promoter, the SV40 promoter, the human cytomegalovirus promoter, and other promoters familiar to those skilled in the art.

A signal sequence from a 5' EST (or cDNAs or genomic DNAs obtainable therefrom) is operably linked to the promoter such that the mRNA transcribed from the promoter will direct the translation of the signal peptide. The host cell, tissue, or organism may be any cell, tissue, or organism which recognizes the signal peptide encoded by the signal sequence in the 5' EST (or cDNA or genomic DNA obtainable therefrom). Suitable hosts include mammalian cells, tissues or organisms, avian cells, tissues, or organisms, insect cells, tissues or organisms, or yeast.

In addition, the secretion vector contains cloning sites for inserting genes encoding the proteins which are to be secreted. The cloning sites facilitate the cloning of the insert gene in frame with the signal sequence such that a fusion protein in which the signal peptide is fused to the protein encoded by the inserted gene is expressed from the mRNA transcribed from the promoter. The signal peptide directs the extracellular secretion of the fusion protein.

The secretion vector may be DNA or RNA and may integrate into the chromosome of the host, be stably maintained as an extrachromosomal replicon in the host, be an artificial chromosome, or be transiently present in the host. Many nucleic acid backbones suitable for

use as secretion vectors are known to those skilled in the art, including retroviral vectors, SV40 vectors, Bovine Papilloma Virus vectors, yeast integrating plasmids, yeast episomal plasmids, yeast artificial chromosomes, human artificial chromosomes, P element vectors, baculovirus vectors, or bacterial plasmids capable of being transiently introduced into the host.

The secretion vector may also contain a polyA signal such that the polyA signal is located downstream of the gene inserted into the secretion vector.

After the gene encoding the protein for which secretion is desired is inserted into the secretion vector, the secretion vector is introduced into the host cell, tissue, or organism using calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection, viral particles or as naked DNA. The protein encoded by the inserted gene is then purified or enriched from the supernatant using conventional techniques such as ammonium sulfate precipitation, immunoprecipitation, immunochromatography, size exclusion chromatography, ion exchange chromatography, and HPLC. Alternatively, the secreted protein may be in a sufficiently enriched or pure state in the supernatant or growth media of the host to permit it to be used for its intended purpose without further enrichment.

The signal sequences may also be inserted into vectors designed for gene therapy. In such vectors, the signal sequence is operably linked to a promoter such that mRNA transcribed from the promoter encodes the signal peptide. A cloning site is located downstream of the signal sequence such that a gene encoding a protein whose secretion is desired may readily be inserted into the vector and fused to the signal sequence. The vector is introduced into an appropriate host cell. The protein expressed from the promoter is secreted extracellularly, thereby producing a therapeutic effect.

The 5' ESTs may also be used to clone sequences located upstream of the 5' ESTs which are capable of regulating gene expression, including promoter sequences, enhancer sequences, and other upstream sequences which influence transcription or translation levels. Once identified and cloned, these upstream regulatory sequences may be used in expression vectors designed to direct the expression of an inserted gene in a desired spatial, temporal, developmental, or quantitative fashion. Example 58 describes a method for cloning sequences upstream of the extended cDNAs or 5' ESTs.

2. Identification of Upstream Sequences With Promoting or Regulatory Activities

EXAMPLE 58

Use of Extended cDNAs or 5' ESTs to Clone Upstream Sequences from Genomic DNA

Sequences derived from extended cDNAs or 5' ESTs may be used to isolate the
5 promoters of the corresponding genes using chromosome walking techniques. In one
chromosome walking technique, which utilizes the GenomeWalker™ kit available from
Clontech, five complete genomic DNA samples are each digested with a different restriction
enzyme which has a 6 base recognition site and leaves a blunt end. Following digestion,
oligonucleotide adapters are ligated to each end of the resulting genomic DNA fragments.

10 For each of the five genomic DNA libraries, a first PCR reaction is performed
according to the manufacturer's instructions (which are incorporated herein by reference)
using an outer adaptor primer provided in the kit and an outer gene specific primer. The gene
specific primer should be selected to be specific for the extended cDNA or 5' EST of interest
and should have a melting temperature, length, and location in the extended cDNA or 5' EST
15 which is consistent with its use in PCR reactions. Each first PCR reaction contains 5 ng of
genomic DNA, 5 µl of 10X Tth reaction buffer, 0.2 mM of each dNTP, 0.2 µM each of outer
adaptor primer and outer gene specific primer, 1.1 mM of Mg(OAc)₂, and 1 µl of the Tth
polymerase 50X mix in a total volume of 50 µl. The reaction cycle for the first PCR reaction
is as follows: 1 min - 94°C / 2 sec - 94°C, 3 min - 72°C (7 cycles) / 2 sec - 94°C, 3 min -
20 67°C (32 cycles) / 5 min - 67°C.

The product of the first PCR reaction is diluted and used as a template for a
second PCR reaction according to the manufacturer's instructions using a pair of nested
primers which are located internally on the amplicon resulting from the first PCR
reaction. For example, 5 µl of the reaction product of the first PCR reaction mixture
25 may be diluted 180 times. Reactions are made in a 50 µl volume having a composition
identical to that of the first PCR reaction except the nested primers are used. The first
nested primer is specific for the adaptor, and is provided with the GenomeWalker™ kit.
The second nested primer is specific for the particular extended cDNA or 5' EST for
which the promoter is to be cloned and should have a melting temperature, length, and
30 location in the extended cDNA or 5' EST which is consistent with its use in PCR
reactions. The reaction parameters of the second PCR reaction are as follows: 1 min -

94°C / 2 sec - 94°C, 3 min - 72°C (6 cycles) / 2 sec - 94°C, 3 min - 67°C (25 cycles) / 5 min - 67°C. The product of the second PCR reaction is purified, cloned, and sequenced using standard techniques.

Alternatively, two or more human genomic DNA libraries can be constructed by using two or more restriction enzymes. The digested genomic DNA is cloned into vectors which can be converted into single stranded, circular, or linear DNA. A biotinylated oligonucleotide comprising at least 15 nucleotides from the extended cDNA or 5' EST sequence is hybridized to the single stranded DNA. Hybrids between the biotinylated oligonucleotide and the single stranded DNA containing the extended cDNA or EST sequence are isolated as described in Example 29 above. Thereafter, the single stranded DNA containing the extended cDNA or EST sequence is released from the beads and converted into double stranded DNA using a primer specific for the extended cDNA or 5' EST sequence or a primer corresponding to a sequence included in the cloning vector. The resulting double stranded DNA is transformed into bacteria. DNAs containing the 5' EST or extended cDNA sequences are identified by colony PCR or colony hybridization.

Once the upstream genomic sequences have been cloned and sequenced as described above, prospective promoters and transcription start sites within the upstream sequences may be identified by comparing the sequences upstream of the extended cDNAs or 5' ESTs with databases containing known transcription start sites, transcription factor binding sites, or promoter sequences.

In addition, promoters in the upstream sequences may be identified using promoter reporter vectors as described in Example .

25

EXAMPLE 59

Identification of Promoters in Cloned Upstream Sequences

The genomic sequences upstream of the extended cDNAs or 5' ESTs are cloned into a suitable promoter reporter vector, such as the pSEAP-Basic, pSEAP-Enhancer, p β gal-Basic, p β gal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline

30

phosphatase, β galactosidase, or green fluorescent protein. The sequences upstream of the extended cDNAs or 5' ESTs are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer for augmenting transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert indicates that a promoter sequence is present in the inserted upstream sequence.

Appropriate host cells for the promoter reporter vectors may be chosen based on the results of the above described determination of expression patterns of the extended cDNAs and ESTs. For example, if the expression pattern analysis indicates that the mRNA corresponding to a particular extended cDNA or 5' EST is expressed in fibroblasts, the promoter reporter vector may be introduced into a human fibroblast cell line.

Promoter sequences within the upstream genomic DNA may be further defined by constructing nested deletions in the upstream DNA using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory sites within the promoter may be identified using site directed mutagenesis or linker scanning to obliterate potential transcription factor binding sites within the promoter individually or in combination. The effects of these mutations on transcription levels may be determined by inserting the mutations into the cloning sites in the promoter reporter vectors.

EXAMPLE 60

Cloning and Identification of Promoters

Using the method described in Example 58 above with 5' ESTs, sequences upstream of several genes were obtained. Using the primer pairs GGG AAG ATG GAG ATA GTA TTG CCT G (SEQ ID NO:29) and CTG CCA TGT ACA TGA TAG AGA GAT TC (SEQ

ID NO:30), the promoter having the internal designation P13H2 (SEQ ID NO:31) was obtained.

Using the primer pairs GTA CCA GGGG ACT GTG ACC ATT GC (SEQ ID NO:32) and CTG TGA CCA TTG CTC CCA AGA GAG (SEQ ID NO:33), the promoter
5 having the internal designation P15B4 (SEQ ID NO:34) was obtained.

Using the primer pairs CTG GGA TGG AAG GCA CGG TA (SEQ ID NO:35) and GAG ACC ACA CAG CTA GAC AA (SEQ ID NO:36), the promoter having the internal designation P29B6 (SEQ ID NO:37) was obtained.

Figure 4 provides a schematic description of the promoters isolated and the way they
10 are assembled with the corresponding 5' tags. The upstream sequences were screened for the presence of motifs resembling transcription factor binding sites or known transcription start sites using the computer program MatInspector release 2.0, August 1996.

Table VII describes the transcription factor binding sites present in each of these promoters. The columns labeled matrice provides the name of the MatInspector matrix used.
15 The column labeled position provides the 5' position of the promoter site. Numeration of the sequence starts from the transcription site as determined by matching the genomic sequence with the 5' EST sequence. The column labeled "orientation" indicates the DNA strand on which the site is found, with the + strand being the coding strand as determined by matching the genomic sequence with the sequence of the 5' EST. The column labeled "score" provides
20 the MatInspector score found for this site. The column labeled "length" provides the length of the site in nucleotides. The column labeled "sequence" provides the sequence of the site found.

Bacterial clones containing plasmids containing the promoter sequences described above described above are presently stored in the inventor's laboratories under the internal
25 identification numbers provided above. The inserts may be recovered from the deposited materials by growing an aliquot of the appropriate bacterial clone in the appropriate medium. The plasmid DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a
30 cesium chloride gradient, size exclusion chromatography, or anion exchange chromatography. The plasmid DNA obtained using these procedures may then be manipulated using standard

cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the EST insertion. The PCR product which corresponds to the 5' EST can then be manipulated using standard cloning techniques familiar to those skilled in the art.

5 The promoters and other regulatory sequences located upstream of the extended cDNAs or 5' ESTs may be used to design expression vectors capable of directing the expression of an inserted gene in a desired spatial, temporal, developmental, or quantitative manner. A promoter capable of directing the desired spatial, temporal, developmental, and quantitative patterns may be selected using the results of the expression analysis described in
10 Example 26 above. For example, if a promoter which confers a high level of expression in muscle is desired, the promoter sequence upstream of an extended cDNA or 5' EST derived from an mRNA which is expressed at a high level in muscle, as determined by the method of Example 26, may be used in the expression vector.

Preferably, the desired promoter is placed near multiple restriction sites to facilitate
15 the cloning of the desired insert downstream of the promoter, such that the promoter is able to drive expression of the inserted gene. The promoter may be inserted in conventional nucleic acid backbones designed for extrachromosomal replication, integration into the host chromosomes or transient expression. Suitable backbones for the present expression vectors include retroviral backbones, backbones from eukaryotic episomes such as SV40 or Bovine
20 Papilloma Virus, backbones from bacterial episomes, or artificial chromosomes.

Preferably, the expression vectors also include a polyA signal downstream of the multiple restriction sites for directing the polyadenylation of mRNA transcribed from the gene inserted into the expression vector.

Following the identification of promoter sequences using the procedures of Examples
25 58-60, proteins which interact with the promoter may be identified as described in Example 61 below.

EXAMPLE 61**Identification of Proteins Which Interact with Promoter Sequences, Upstream
Regulatory Sequences, or mRNA**

Sequences within the promoter region which are likely to bind transcription factors
5 may be identified by homology to known transcription factor binding sites or through
conventional mutagenesis or deletion analyses of reporter plasmids containing the promoter
sequence. For example, deletions may be made in a reporter plasmid containing the promoter
sequence of interest operably linked to an assayable reporter gene. The reporter plasmids
carrying various deletions within the promoter region are transfected into an appropriate host
10 cell and the effects of the deletions on expression levels is assessed. Transcription factor
binding sites within the regions in which deletions reduce expression levels may be further
localized using site directed mutagenesis, linker scanning analysis, or other techniques familiar
to those skilled in the art.

Nucleic acids encoding proteins which interact with sequences in the promoter
15 may be identified using one-hybrid systems such as those described in the manual
accompanying the Matchmaker One-Hybrid System kit available from Clontech (Catalog
No. K1603-1), the disclosure of which is incorporated herein by reference. Briefly, the
Matchmaker One-hybrid system is used as follows. The target sequence for which it is
desired to identify binding proteins is cloned upstream of a selectable reporter gene and
20 integrated into the yeast genome. Preferably, multiple copies of the target sequences are
inserted into the reporter plasmid in tandem. A library comprised of fusions between
cDNAs to be evaluated for the ability to bind to the promoter and the activation domain
of a yeast transcription factor, such as GAL4, is transformed into the yeast strain
containing the integrated reporter sequence. The yeast are plated on selective media to
25 select cells expressing the selectable marker linked to the promoter sequence. The
colonies which grow on the selective media contain genes encoding proteins which bind
the target sequence. The inserts in the genes encoding the fusion proteins are further
characterized by sequencing. In addition, the inserts may be inserted into expression
vectors or *in vitro* transcription vectors. Binding of the polypeptides encoded by the
30 inserts to the promoter DNA may be confirmed by techniques familiar to those skilled in
the art, such as gel shift analysis or DNase protection analysis.

VII. Use of 5' ESTs (or cDNAs or Genomic DNAs Obtainable Therefrom) in Gene Therapy

The present invention also comprises the use of 5'ESTs (or cDNA or genomic DNA obtainable therefrom) in gene therapy strategies, including antisense and triple helix strategies as described in Examples 62 and 63 below. In antisense approaches, nucleic acid sequences complementary to an mRNA are hybridized to the mRNA intracellularly, thereby blocking the expression of the protein encoded by the mRNA. The antisense sequences may prevent gene expression through a variety of mechanisms. For example, the antisense sequences may inhibit the ability of ribosomes to translate the mRNA. Alternatively, the antisense sequences may block transport of the mRNA from the nucleus to the cytoplasm, thereby limiting the amount of mRNA available for translation. Another mechanism through which antisense sequences may inhibit gene expression is by interfering with mRNA splicing. In yet another strategy, the antisense nucleic acid may be incorporated in a ribozyme capable of specifically cleaving the target mRNA.

EXAMPLE 62

Preparation and Use of Antisense Oligonucleotides

The antisense nucleic acid molecules to be used in gene therapy may be either DNA or RNA sequences. They may comprise a sequence complementary to the sequence of the 5'EST (or cDNA or genomic DNA obtainable therefrom). The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex with sufficient stability to inhibit the expression of the mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green *et al.*, *Ann. Rev. Biochem.* 55:569-597, 1986; and Izant and Weintraub, *Cell* 36:1007-1015, 1984, which are hereby incorporated by reference.

In some strategies, antisense molecules are obtained from a nucleotide sequence encoding a protein by reversing the orientation of the coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules may be transcribed using *in vitro* transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach

involves transcription of the antisense nucleic acids *in vivo* by operably linking DNA containing the antisense sequence to a promoter in an expression vector.

Alternatively, oligonucleotides which are complementary to the strand normally transcribed in the cell may be synthesized *in vitro*. Thus, the antisense nucleic acids are complementary to the corresponding mRNA and are capable of hybridizing to the mRNA to create a duplex. In some embodiments, the antisense sequences may contain modified sugar phosphate backbones to increase stability and make them less sensitive to RNase activity. Examples of modifications suitable for use in antisense strategies are described by Rossi *et al.*, *Pharmacol. Ther.* 50(2):245-254, 1991, which is hereby incorporated by reference.

Various types of antisense oligonucleotides complementary to the sequence of the 5'EST (or cDNA or genomic DNA obtainable therefrom) may be used. In one preferred embodiment, stable and semi-stable antisense oligonucleotides described in International Application No. PCT WO94/23026, hereby incorporated by reference, are used. In these molecules, the 3' end or both the 3' and 5' ends are engaged in intramolecular hydrogen bonding between complementary base pairs. These molecules are better able to withstand exonuclease attacks and exhibit increased stability compared to conventional antisense oligonucleotides.

In another preferred embodiment, the antisense oligodeoxynucleotides against herpes simplex virus types 1 and 2 described in International Application No. WO 95/04141, hereby incorporated by reference, are used.

In yet another preferred embodiment, the covalently cross-linked antisense oligonucleotides described in International Application No. WO 96/31523, hereby incorporated by reference, are used. These double- or single-stranded oligonucleotides comprise one or more, respectively, inter- or intra-oligonucleotide covalent cross-linkages, wherein the linkage consists of an amide bond between a primary amine group of one strand and a carboxyl group of the other strand or of the same strand, respectively, the primary amine group being directly substituted in the 2' position of the strand nucleotide monosaccharide ring, and the carboxyl group being carried by an aliphatic spacer group substituted on a nucleotide or nucleotide analog of the other strand or the same strand, respectively.

The antisense oligodeoxynucleotides and oligonucleotides disclosed in International Application No. WO 92/18522, incorporated by reference, may also be used. These molecules are stable to degradation and contain at least one transcription control recognition sequence which binds to control proteins and are effective as decoys therefore. These
5 molecules may contain "hairpin" structures, "dumbbell" structures, "modified dumbbell" structures, "cross-linked" decoy structures and "loop" structures.

In another preferred embodiment, the cyclic double-stranded oligonucleotides described in European Patent Application No. 0 572 287 A2, hereby incorporated by reference are used. These ligated oligonucleotide "dumbbells" contain the binding site for a
10 transcription factor and inhibit expression of the gene under control of the transcription factor by sequestering the factor.

Use of the closed antisense oligonucleotides disclosed in International Application No. WO 92/19732, hereby incorporated by reference, is also contemplated. Because these molecules have no free ends, they are more resistant to degradation by exonucleases than are
15 conventional oligonucleotides. These oligonucleotides may be multifunctional, interacting with several regions which are not adjacent to the target mRNA.

The appropriate level of antisense nucleic acids required to inhibit gene expression may be determined using *in vitro* expression analysis. The antisense molecule may be introduced into the cells by diffusion, injection, infection, transfection or h-region-mediated
20 import using procedures known in the art. For example, the antisense nucleic acids can be introduced into the body as a bare or naked oligonucleotide, oligonucleotide encapsulated in lipid, oligonucleotide sequence encapsidated by viral protein, or as an oligonucleotide operably linked to a promoter contained in an expression vector. The expression vector may be any of a variety of expression vectors known in the art, including retroviral or viral vectors,
25 vectors capable of extrachromosomal replication, or integrating vectors. The vectors may be DNA or RNA.

The antisense molecules are introduced onto cell samples at a number of different concentrations preferably between 1×10^{-10} M to 1×10^{-4} M. Once the minimum concentration that can adequately control gene expression is identified, the optimized dose is translated into
30 a dosage suitable for use *in vivo*. For example, an inhibiting concentration in culture of 1×10^{-7} translates into a dose of approximately 0.6 mg/kg bodyweight. Levels of oligonucleotide

approaching 100 mg/kg bodyweight or higher may be possible after testing the toxicity of the oligonucleotide in laboratory animals. It is additionally contemplated that cells from the vertebrate are removed, treated with the antisense oligonucleotide, and reintroduced into the vertebrate.

5 It is further contemplated that the antisense oligonucleotide sequence is incorporated into a ribozyme sequence to enable the antisense to specifically bind and cleave its target mRNA. For technical applications of ribozyme and antisense oligonucleotides see Rossi *et al.*, *supra*.

10 In a preferred application of this invention, the polypeptide encoded by the gene is first identified, so that the effectiveness of antisense inhibition on translation can be monitored using techniques that include but are not limited to antibody-mediated tests such as RIAs and ELISA, functional assays, or radiolabeling.

15 The 5' ESTs of the present invention (or cDNAs or genomic DNAs obtainable therefrom) may also be used in gene therapy approaches based on intracellular triple helix formation. Triple helix oligonucleotides are used to inhibit transcription from a genome. They are particularly useful for studying alterations in cell activity as it is associated with a particular gene. The 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) of the present invention or, more preferably, a portion of those sequences, can be used to inhibit gene expression in individuals having diseases associated with expression of a particular gene. Similarly, a portion of 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) can be used to study the effect of inhibiting transcription of a particular gene within a cell. Traditionally, homopurine sequences were considered the most useful for triple helix strategies. However, homopyrimidine sequences can also inhibit gene expression. Such homopyrimidine oligonucleotides bind to the major groove at homopurine:homopyrimidine sequences. Thus, both types of sequences from the 5'EST or from the gene corresponding to the 5'EST are contemplated within the scope of this invention.

25

EXAMPLE 63Preparation and Use of Triple Helix Probes

The sequences of the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) are scanned to identify 10-mer to 20-mer homopyrimidine or homopurine stretches which could be used in triple-helix based strategies for inhibiting gene expression. Following identification of candidate homopyrimidine or homopurine stretches, their efficiency in inhibiting gene expression is assessed by introducing varying amounts of oligonucleotides containing the candidate sequences into tissue culture cells which normally express the target gene. The oligonucleotides may be prepared on an oligonucleotide synthesizer or they may be purchased commercially from a company specializing in custom oligonucleotide synthesis, such as GENSET, Paris, France.

The oligonucleotides may be introduced into the cells using a variety of methods known to those skilled in the art, including but not limited to calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection or native uptake.

Treated cells are monitored for altered cell function or reduced gene expression using techniques such as Northern blotting, RNase protection assays, or PCR based strategies to monitor the transcription levels of the target gene in cells which have been treated with the oligonucleotide. The cell functions to be monitored are predicted based upon the homologies of the target gene corresponding to the extended cDNA from which the oligonucleotide was derived with known gene sequences that have been associated with a particular function. The cell functions can also be predicted based on the presence of abnormal physiologies within cells derived from individuals with a particular inherited disease, particularly when the extended cDNA is associated with the disease using techniques described in Example 56.

The oligonucleotides which are effective in inhibiting gene expression in tissue culture cells may then be introduced *in vivo* using the techniques described above and in Example 62 at a dosage calculated based on the *in vitro* results, as described in Example 62.

In some embodiments, the natural (beta) anomers of the oligonucleotide units can be replaced with alpha anomers to render the oligonucleotide more resistant to nucleases. Further, an intercalating agent such as ethidium bromide, or the like, can be attached to the 3' end of the alpha oligonucleotide to stabilize the triple helix. For information on the

generation of oligonucleotides suitable for triple helix formation see Griffin *et al.*, *Science* 245:967-971, 1989, which is hereby incorporated by this reference.

EXAMPLE 64

5 Use of cDNAs Obtained Using the 5' ESTs to Express an Encoded Protein in a Host Organism

The cDNAs obtained as described above using the 5' ESTs of the present invention may also be used to express an encoded protein in a host organism to produce a beneficial effect. In such procedures, the encoded protein may be transiently expressed in the host organism or stably expressed in the host organism. The encoded protein may have any of the activities described above. The encoded protein may be a protein which the host organism lacks or, alternatively, the encoded protein may augment the existing levels of the protein in the host organism.

15 A full length extended cDNA encoding the signal peptide and the mature protein, or an extended cDNA encoding only the mature protein is introduced into the host organism. The extended cDNA may be introduced into the host organism using a variety of techniques known to those of skill in the art. For example, the extended cDNA may be injected into the host organism as naked DNA such that the encoded protein is expressed in the host organism, thereby producing a beneficial effect.

Alternatively, the extended cDNA may be cloned into an expression vector downstream of a promoter which is active in the host organism. The expression vector may be any of the expression vectors designed for use in gene therapy, including viral or retroviral vectors. The expression vector may be directly introduced into the host organism such that the encoded protein is expressed in the host organism to produce a beneficial effect. In another approach, the expression vector may be introduced into cells *in vitro*. Cells containing the expression vector are thereafter selected and introduced into the host organism, where they express the encoded protein to produce a beneficial effect.

EXAMPLE 65

Use of Signal Peptides Encoded by 5' ESTs or Sequences obtained Therefrom
to Import Proteins Into Cells

The short core hydrophobic region (h) of signal peptides encoded by the 5'ESTS or
5 extended cDNAs derived from SEQ ID NOs: 38-195 may also be used as a carrier to import
a peptide or a protein of interest, so-called cargo, into tissue culture cells (Lin *et al.*, *J. Biol.*
Chem., 270: 14225-14258, 1995; Du *et al.*, *J. Peptide Res.*, 51: 235-243, 1998; Rojas *et al.*,
Nature Biotech., 16: 370-375, 1998).

When cell permeable peptides of limited size (approximately up to 25 amino acids)
10 are to be translocated across cell membrane, chemical synthesis may be used in order to add
the h region to either the C-terminus or the N-terminus to the cargo peptide of interest.
Alternatively, when longer peptides or proteins are to be imported into cells, nucleic acids can
be genetically engineered, using techniques familiar to those skilled in the art, in order to link
the extended cDNA sequence encoding the h region to the 5' or the 3' end of a DNA
15 sequence coding for a cargo polypeptide. Such genetically engineered nucleic acids are then
translated either *in vitro* or *in vivo* after transfection into appropriate cells, using conventional
techniques to produce the resulting cell permeable polypeptide. Suitable hosts cells are then
simply incubated with the cell permeable polypeptide which is then translocated across the
membrane.

20 This method may be applied to study diverse intracellular functions and cellular
processes. For instance, it has been used to probe functionally relevant domains of
intracellular proteins and to examine protein-protein interactions involved in signal
transduction pathways (Lin *et al.*, *supra*; Lin *et al.*, *J. Biol. Chem.*, 271: 5305-5308, 1996;
Rojas *et al.*, *J. Biol. Chem.*, 271: 27456-27461, 1996; Liu *et al.*, *Proc. Natl. Acad. Sci. USA*,
25 93: 11819-11824, 1996; Rojas *et al.*, *Bioch. Biophys. Res. Commun.*, 234: 675-680, 1997).

Such techniques may be used in cellular therapy to import proteins producing
therapeutic effects. For instance, cells isolated from a patient may be treated with imported
therapeutic proteins and then re-introduced into the host organism.

Alternatively, the h region of signal peptides of the present invention could be used in
30 combination with a nuclear localization signal to deliver nucleic acids into cell nucleus. Such
oligonucleotides may be antisense oligonucleotides or oligonucleotides designed to form

triple helixes, as described in examples 62 and 63 respectively, in order to inhibit processing and/or maturation of a target cellular RNA.

As discussed above, the cDNAs or portions thereof obtained using the 5' ESTs of the present invention can be used for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination for expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris *et al.*, *Cell* 75:791-803, 1993, the disclosure of which is hereby incorporated by reference) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins or polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins

involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

5 Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation *Molecular Cloning; A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, Fritsch and Maniatis eds., 1989, and *Methods in Enzymology; Guide to Molecular Cloning Techniques*, Academic Press, Berger and Kimmel eds., 1987.

10 Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid
15 preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Although this invention has been described in terms of certain preferred embodiments, other embodiments which will be apparent to those of ordinary skill in the art in view of the
20 disclosure herein are also within the scope of this invention. Accordingly, the scope of the invention is intended to be defined only by reference to the appended claims. All documents cited herein are incorporated herein by reference in their entirety.

Step	Search characteristic		Selection Characteristics		
	Program	Strand	Parameters	Identity (%)	Length (bp)
Miscellaneous	blastn	both	S=61 X=16	90	17
tRNA	fasta	both	-	80	60
rRNA	blastn	both	S=108	80	40
mtRNA	blastn	both	S=108	80	40
Prokaryotic	blastn	both	S=144	90	40
Fungal	blastn	both	S=144	90	40
Alu	fasta*	both	-	70	40
L1	blastn	both	S=72	70	40
Repeats	blastn	both	S=72	70	40
Promoters	blastn	top	S=54 X=16	90	15†
Vertebrate	fasta*	both	S=108	90	30
ESTs	blastn	both	S=108 X=16	90	30
Proteins	blastx [□]	top	E = 0.001	-	-

Table 1: Parameters used for each step of EST analysis

- * use "Quick Fast" Database scanner
- † alignment further constrained to begin closer than 10bp to EST's end
- using BLOSUM62 substitution matrix

TABLE II

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELINE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID38	new	14.3	Substantia nigra	47-39-4-A10-PU
ID39	new	11.1	Fetal brain	57-9-4-C5-PU
ID40	new	10.6	Fetal brain	57-19-1-B11-PU
ID41	new	9.1	Fetal brain	57-7-1-G12-PU
ID42	new	8.8	Substantia nigra	47-22-3-D2-PU
ID43	new	8.7	Fetal brain	57-21-2-H11-PU
ID44	new	8.4	Substantia nigra	47-37-3-F6-PU
ID45	new	8.2	Substantia nigra	47-54-1-A8-PU
ID46	new	8.2	Substantia nigra	47-15-1-E5-PU
ID47	new	8	Substantia nigra	47-24-1-A6-PU
ID48	new	7.8	Fetal brain	57-10-3-H10-PU
ID49	new	7.7	Substantia nigra	47-17-1-D7-PU
ID50	new	7.6	Cerebellum	55-9-4-A4-PU
ID51	new	7.5	Substantia nigra	47-18-3-C2-PU
ID52	new	7.4	Fetal brain	57-19-1-C8-PU
ID53	new	7.4	Substantia nigra	47-1-4-C5-PU
ID54	new	7.2	Substantia nigra	47-24-1-B5-PU
ID55	new	7.2	Cerebellum	55-6-2-A9-PU
ID56	new	7.1	Substantia nigra	47-30-4-A8-PU
ID57	new	6.9	Fetal brain	57-21-1-D5-PU
ID58	new	6.9	Substantia nigra	47-4-2-C7-PU
ID59	new	6.8	Substantia nigra	47-12-3-A8-PU
ID60	new	6.8	Substantia nigra	47-7-4-E6-PU
ID61	new	6.8	Substantia nigra	47-15-3-G3-PU
ID62	new	6.7	Fetal brain	57-20-2-B5-PU
ID63	new	6.6	Substantia nigra	47-2-2-E6-PU
ID64	new	6.5	Fetal brain	57-25-1-G3-PU
ID65	new	6.5	Fetal brain	57-7-4-B12-PU
ID66	new	6.5	Substantia nigra	47-21-1-D9-PU
ID67	new	6.4	Substantia nigra	47-31-2-H9-PU
ID68	new	5.7	Cerebellum	55-1-3-D11-PU
ID69	new	5.7	Cerebellum	55-7-2-A1-PU
ID70	new	5.7	Fetal brain	57-28-3-C1-PU
ID71	new	5.6	Fetal brain	57-9-4-D11-PU
ID72	new	5.5	Substantia nigra	47-7-4-C10-PU
ID73	new	5.5	Fetal brain	57-22-1-E11-PU
ID74	new	5.4	Fetal brain	57-20-2-D9-PU
ID75	new	5.4	Substantia nigra	47-39-3-E7-PU
ID76	new	5.4	Surrenals	62-3-1-G5-PU
ID77	new	5.4	Fetal brain	57-18-4-H5-PU
ID78	new	5.4	Fetal brain	57-22-2-H8-PU
ID79	new	5.3	Fetal brain	57-22-2-E12-PU
ID80	new	5.3	Fetal brain	57-23-3-B8-PU
ID81	new	5.2	Fetal brain	57-6-3-C5-PU
ID82	new	5.2	Substantia nigra	47-7-1-D2-PU
ID83	new	5.1	Fetal brain	57-7-2-G9-PU
ID84	new	5.1	Fetal brain	57-10-3-D3-PU
ID85	new	5.1	Substantia nigra	47-4-4-F2-PU
ID86	new	5.1	Fetal brain	57-4-4-H6-PU

<u>SEQ. ID</u> <u>NO.</u>	<u>CATEGORY</u>	<u>VON HEIJNE</u> <u>SCORE</u>	<u>TISSUE</u> <u>SOURCE</u>	<u>INTERNAL</u> <u>DESIGNATION</u>
ID87	new	5	Substantia nigra	47-10-2-G12-PU
ID88	new	5	Cerebellum	55-10-3-E12-PU
ID89	new	4.9	Substantia nigra	47-8-2-D1-PU
ID90	new	4.9	Fetal brain	57-3-4-C9-PU
ID91	new	4.9	Substantia nigra	47-14-1-C3-PU
ID92	new	4.8	Substantia nigra	47-3-4-C8-PU
ID93	new	4.8	Substantia nigra	47-15-1-B10-PU
ID94	new	4.8	Fetal brain	57-26-3-A12-PU
ID95	new	4.7	Substantia nigra	47-26-3-B10-PU
ID96	new	4.7	Substantia nigra	47-26-1-B6-PU
ID97	new	4.7	Surrenals	62-5-1-B8-PU
ID98	new	4.6	Substantia nigra	47-15-4-H9-PU
ID99	new	4.6	Cerebellum	55-2-4-D3-PU
ID100	new	4.5	Fetal brain	57-6-1-B1-PU
ID101	new	4.5	Fetal brain	57-26-4-E4-PU
ID102	new	4.5	Substantia nigra	47-2-2-A7-PU
ID103	new	4.5	Substantia nigra	47-55-2-B3-PU
ID104	new	4.5	Substantia nigra	47-54-1-C9-PU
ID105	new	4.4	Cerebellum	55-8-2-A2-PU
ID106	new	4.4	Substantia nigra	47-4-2-H4-PU
ID107	new	4.3	Fetal brain	57-27-3-B11-PU
ID108	new	4.2	Fetal brain	57-22-4-D2-PU
ID109	new	4.2	Substantia nigra	47-20-4-E2-PU
ID110	new	4.2	Substantia nigra	47-2-3-H2-PU
ID111	new	4.1	Substantia nigra	47-22-3-G5-PU
ID112	new	4.1	Fetal brain	57-18-3-A5-PU
ID113	new	4.1	Fetal brain	57-9-3-H7-PU
ID114	new	4.1	Surrenals	62-5-2-B6-PU
ID115	new	4	Cerebellum	55-12-1-E12-PU
ID116	new	4	Fetal brain	57-20-1-A5-PU
ID117	new	3.9	Substantia nigra	47-22-4-F6-PU
ID118	new	3.8	Fetal brain	57-19-3-E1-PU
ID119	new	3.8	Substantia nigra	47-18-3-G5-PU
ID120	new	3.8	Substantia nigra	47-20-1-G3-PU
ID121	new	3.8	Fetal brain	57-6-4-A1-PU
ID122	new	3.8	Fetal brain	57-27-3-G10-PU
ID123	new	3.7	Substantia nigra	47-2-4-C7-PU
ID124	new	3.7	Cerebellum	55-6-1-E6-PU
ID125	new	3.6	Fetal brain	57-4-4-F7-PU
ID126	new	3.6	Substantia nigra	47-30-2-B1-PU
ID127	new	3.6	Substantia nigra	47-29-1-F11-PU
ID128	new	3.6	Substantia nigra	47-39-3-D4-PU
ID129	new	3.5	Substantia nigra	47-15-2-G3-PU
ID130	new	3.5	Fetal brain	57-18-3-E6-PU
ID131	new	3.5	Substantia nigra	47-40-2-G6-PU
ID132	new	3.5	Fetal brain	57-6-4-D7-PU
ID133	new	3.5	Substantia nigra	47-55-4-A8-PU
ID134	ext-est-not-vrt	9.8	Substantia nigra	47-39-4-B9-PU
ID135	ext-est-not-vrt	9.2	Cerebellum	55-11-1-H5-PU
ID136	ext-est-not-vrt	9	Substantia nigra	47-4-4-G1-PU
ID137	ext-est-not-vrt	7.2	Substantia nigra	47-2-3-G9-PU

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HEIJNE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID138	ext-est-not-vrt	7.2	Cerebellum	55-10-3-F5-PU
ID139	ext-est-not-vrt	5.6	Fetal brain	57-4-4-G6-PU
ID140	ext-est-not-vrt	4.2	Cerebellum	55-7-1-D11-PU
ID141	ext-est-not-vrt	3.7	Substantia nigra	47-19-2-F7-PU
ID142	ext-est-not-vrt	3.7	Substantia nigra	47-1-4-D2-PU
ID143	ext-est-not-vrt	3.6	Cerebellum	55-5-4-A6-PU
ID144	ext-est-not-vrt	3.6	Cerebellum	55-4-4-H3-PU
ID145	ext-est-not-vrt	3.5	Cerebellum	55-3-1-G6-PU
ID146	ext-est-not-vrt	3.5	Substantia nigra	47-55-2-H2-PU
ID147	est-not-ext	12.4	Substantia nigra	47-39-4-H8-PU
ID148	est-not-ext	11.4	Fetal brain	57-26-4-A4-PU
ID149	est-not-ext	11.1	Substantia nigra	47-2-3-D1-PU
ID150	est-not-ext	9.2	Substantia nigra	47-4-1-E4-PU
ID151	est-not-ext	9	Substantia nigra	47-40-4-G9-PU
ID152	est-not-ext	8.8	Fetal brain	57-5-4-G3-PU
ID153	est-not-ext	7.5	Substantia nigra	47-13-4-C1-PU
ID154	est-not-ext	7.4	Fetal brain	57-20-4-E2-PU
ID155	est-not-ext	7	Substantia nigra	47-24-4-H4-PU
ID156	est-not-ext	6.9	Substantia nigra	47-26-2-B2-PU
ID157	est-not-ext	6.8	Substantia nigra	47-11-1-A2-PU
ID158	est-not-ext	6.4	Fetal brain	57-19-2-G8-PU
ID159	est-not-ext	6.4	Fetal brain	57-19-4-H8-PU
ID160	est-not-ext	6.3	Substantia nigra	47-39-2-A11-PU
ID161	est-not-ext	6.2	Fetal brain	57-24-2-B4-PU
ID162	est-not-ext	5.9	Surrenals	62-1-1-G3-PU
ID163	est-not-ext	5.7	Fetal brain	57-2-4-H4-PU
ID164	est-not-ext	5.6	Fetal brain	57-8-2-D3-PU
ID165	est-not-ext	5.5	Cerebellum	55-11-4-G2-PU
ID166	est-not-ext	5.4	Substantia nigra	47-24-1-B6-PU
ID167	est-not-ext	5.4	Substantia nigra	47-55-3-B10-PU
ID168	est-not-ext	5.4	Surrenals	62-8-1-B12-PU
ID169	est-not-ext	5.3	Substantia nigra	47-39-1-C9-PU
ID170	est-not-ext	5.3	Fetal brain	57-20-2-F1-PU
ID171	est-not-ext	5.2	Fetal brain	57-25-1-F10-PU
ID172	est-not-ext	5.2	Fetal brain	57-28-4-B12-PU
ID173	est-not-ext	5.1	Substantia nigra	47-15-2-D12-PU
ID174	est-not-ext	5.1	Substantia nigra	47-2-3-G3-PU
ID175	est-not-ext	4.9	Substantia nigra	47-40-3-D8-PU
ID176	est-not-ext	4.9	Substantia nigra	47-40-3-G11-PU
ID177	est-not-ext	4.9	Substantia nigra	47-14-3-D2-PU
ID178	est-not-ext	4.8	Substantia nigra	47-19-1-B7-PU
ID179	est-not-ext	4.8	Substantia nigra	47-19-1-A3-PU
ID180	est-not-ext	4.7	Substantia nigra	47-55-3-G2-PU
ID181	est-not-ext	4.6	Substantia nigra	47-3-4-G7-PU
ID182	est-not-ext	4.5	Substantia nigra	47-29-1-B7-PU
ID183	est-not-ext	4.4	Fetal brain	57-21-4-G6-PU
ID184	est-not-ext	4.3	Substantia nigra	47-2-1-E12-PU
ID185	est-not-ext	4.3	Substantia nigra	47-9-4-D2-PU
ID186	est-not-ext	4.3	Fetal brain	57-2-4-F8-PU
ID187	est-not-ext	4.3	Fetal brain	57-18-1-D5-PU
ID188	est-not-ext	4.2	Substantia nigra	47-8-4-D2-PU

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELINE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID189	est-not-ext	4.1	Substantia nigra	47-17-3-H11-PU
ID190	est-not-ext	3.9	Fetal brain	57-28-2-G6-PU
ID191	est-not-ext	3.7	Fetal brain	57-27-3-G1-PU
ID192	est-not-ext	3.7	Substantia nigra	47-37-4-G11-PU
ID193	est-not-ext	3.7	Surrenals	62-11-3-A2-PU
ID194	ext-vrt-not-genomic	10.9	Surrenals	62-10-2-E4-PU
ID195	ext-vrt-not-genomic	8.9	Substantia nigra	47-14-3-H7-PU

TABLE III

<u>SEQ. ID NO.</u>	<u>SIGNAL PEPTIDE</u>
ID38	MRVRIGLTLLLCVLLSLASA
ID39	MGLHLRPYRVGLLPDGLLFLLLMLLA
ID40	MATLSFVFLLLGAVSWPPASA
ID41	MFLFLSPATPVLPPSLDSRDLLPHLFWGRAGSSSSPALSPVLCRLGLVSLAFQ
ID42	MWAMESGHLLWALLFMQSLWP
ID43	MTHYRNILGLLCCVLATMA
ID44	MKLLLLLASLIERSS
ID45	MARNQALVCLPSFQNAFIPVEDLPTSFXLFLALCASFS
ID46	MPNESWQIPCGKQEAETLNFQSLLLLFYSFYVLA
ID47	MQTTTFIDVTVDQHVAKSNDHLSVLVLLICLVSSYLP
ID48	MALGEEKAEAEASXDTKAQSYGRGSCRERELDIPGPMSGEQPPRLEAEGGLISPVWGAEX YLP LLAGLGTLA
ID49	MTSLYLKHLLCISPFVPFTSG
ID50	MTDSPNAHGLALTTKWMMPAVSLNLTYYLPSWYLCLATLTLFHTSFS
ID51	MASSHWNETTTSVYQYLGQVQKIYPFDNWNTACFVILLFIFTVVS
ID52	MSLLFVFCLECSIFLLNMWVACLLS
ID53	MFVVTVLLLLPLVAFITL
ID54	MNRSCRNTGIYALQFLFVFA
ID55	MTQTTWGAPTRASNHPLPAWLTL SLLAWVTLTHL
ID56	MLKXXAVLCVCAAAWC
ID57	MISAHCNLHLLGSSISPASA
ID58	MXXKACRTLAWLPXPFLPFLSLPLDQT
ID59	MAVKRLGLLL VFLPHQRG
ID60	MQAVDNLTSAPGNTSLCTRDKITQVLFPLLYTVLFFVGLITNGLAMRIFFQIRSKSNFI IFLKNTVISDMLMLTFPFKLS
ID61	MWTLPSLSASFQFLGSLRPSHILWFFLPSLXCPEC
ID62	MSLTDVPM SLLFQPSHSATG
ID63	MDWSLAFLLVSLYWSHM
ID64	MYLLILFFMVGRIP
ID65	MNKPPWEESWGQNQLSGEPATWSLCISPLPGREPSLLVVSCLLFHQA
ID66	MLILELTMMLSFLLSIDSLSVSG
ID67	MKLQRSRAFRIECSAILRRAERLVWNDVCSESQSQRSDCLLGAAWASRLRT
ID68	MVIFTLCVFTLPFLCA
ID69	MWGALPVLVVGTVSSQGQA
ID70	MTRLVCGFLQISLSLA
ID71	MNLLPLLLHHLTFH
ID72	MLSARDRRDRHP EEGVVAELQGFAVDKAFLTSHKGILLETALTLIIFICTASISA
ID73	MLTMSVTLSPRS
ID74	MFXPVALIFPISVSDPTIHPITQAQNLESXLQSFLLISSVRPISQ
ID75	MLLFFPFGETVSLHHPWCVA VLRSWLAASS
ID76	MPLKNLFSVGLWDPYNLLKKHVLVVCYLSWRVSS
ID77	MAMAQKLSHLLPSLRQVIQ
ID78	MIAPTLKGT PSSSAPLALVALAPHSVQK
ID79	MCLFPVSPCPAYSFSSEXGAVLLL VESLCLVFNLLS
ID80	MKIAVLCFLLLIIF
ID81	MCSPRSPLNLSLVPVGAVLLSSLPISP
ID82	MGLHISLIKFLLANGPHIPSHQRPFEKGEKSCRIEVVTLPLTSHCLA
ID83	MKTTYVIFMQSKALLTLYV FVASSMQ
ID84	MNALVFLIFLRFINI
ID85	MQLGPLHTVSTPFFFCWGFLLTGHSLSHS
ID86	MGRGWERTVCSLGWRGGPDPLSWATCWSGARSRHTRVSSIVNGYVGSVCCCVGPLRG

SEQ. ID NO.	SIGNAL PEPTIDE
ID87	MPEAVEQSAHLFVTWSSQRALS
ID88	MPGHTFTFKSCWLIASVPLVFW
ID89	MLLLTFKWFLFCLIGLDLLCQV
ID90	MATTGRRQAEPPVVRPAHSRPPRPVPGSSSLGLAGLMSPVPNLHLLPLTTP
ID91	MVPFTYLQAHFTLCSG
ID92	MFFSFLLTINLVSL
ID93	MWPGRECKNWGLLCFASECTT
ID94	MLTPFSLEEKLLECHYVLAKLAGACLLTLRQPPTH
ID95	MKRIKSMMGKVEHKIKGEKQRSRHKIVFVGLIFLKSSA
ID96	MQSALCLFCKICPFTHG
ID97	MMHNIIVKELIVTFFLGITVVQMLISVTGLKGVEAQNGSESEVFVGKYETLVFYWPSLLC LAFLGRFL
ID98	MSNCLQNFLKITSTRLLCSRLCQQLRS
ID99	MSGXGLFLRTTAAARA
ID100	MNPLHKHCAAGPLTWLHLLSHLKS
ID101	MPKDKRGARHNSPHFSFAVLRVLHLPALT
ID102	MTIHVLRKCCQMGRNNEWLPGLVIPLCVSRQLLTGART
ID103	MQAASFGRGRNGLDNWGIAALLGLLQLRFK
ID104	MSPSLGDRCSSWLHLVSHLESISGPLLNIPENLLCCHRCTNC
ID105	MSGAEPTTFIRYFLLPCLINLAIG
ID106	MVYDYFISQQLFSFLLSTIPT
ID107	MLFLCSCSLSLNQL
ID108	MFFLMVLLFRSNKWT
ID109	MLPLQGLCTCYFLHLEFLSHVTTSLASS
ID110	MYFYGLTFHFLLNTILLFG
ID111	MRWNLFFFCILRNQTKLWASQGSQDAQS
ID112	MFIAALFTMAKTWN
ID113	MPGXKHFLRVFRXSAXRSVGYYXKPGTSRASLWVXLPPXXVIA
ID114	MRLESPDENFAVVQEHAIHHIDGPLRRFLLLEVHEPVALGPLFVTGHFA
ID115	MAGSPDREVLLPTVLRGSYC
ID116	MHVSMLEGFDENLDVQELILQDAFQVWDPKSLIRKGRERHLFLFEISLVFS
ID117	MNVGTAHSEVNXTNRMKXRGIWLSYVLAIGLLHIVLLS
ID118	MLSFXAIXXYIPTNS
ID119	MDEYSWWCHVLEVVKQMFTFINITLWLGSLC
ID120	MRRKGQGHAFIFLIQWKTCLS
ID121	MFLISGHVHLIYNILFLAVSSFMP
ID122	MTPRILSEVQFSAFCPYWTIARILERVGSACFRLELCAAIVGYFVLDVRTFLFVVCVIC VTLN
ID123	MCSLLSGWGQLLRC
ID124	MLFSFCFPVHFVNPSSLFPPSSVSLIPFNFSASGLCA
ID125	MTWILRILFVIGSXL
ID126	MSSTYCGNSSAKMSVNEVSASFSLSEQKTGFAFVGILCIFLGLLIIRC
ID127	MTDIWLTMLSMIVGATCY
ID128	MXXCWYAFISLGYLG
ID129	MFIRTLKTTVLPFMRTAPQLALS WVPPXCRV
ID130	MRTGAEMRTNSSVLIFFCLLPYTYH
ID131	MIVIPSWLENEGLELGFSHRTFA
ID132	MLKKEIAHHSPLVSCPVTCTKYRTLRLRLVISVFLSFLPSYP
ID133	MTXPSRAQTVDXGIAKHCAYSLPGVALTLG
ID134	MPFRLLIPLGLLCALLPQHHG
ID135	MXLVLVFLCSLLAPMVA
ID136	MALRRPPRLRLCARLPDFFLLLLFRGCLIG
ID137	MGGNGSTCKPDTERQGTLSAAPTTSAPCLSNHHNKKHLILAFCAGVLLTLLLI AFIFL

<u>SEQ. ID</u> <u>NO.</u>	<u>SIGNAL PEPTIDE</u>
ID138	MGTADSDMAPEAPQHTHIDVHIHQESALAKLLLTCCSA
ID139	MSLLSFLFARVNLG
ID140	MARSPLRRRGRPTWSLSTPRPGSPTSSSRSWWCCPARLTLTSG
ID141	MKITTTLTLLACHLQLEVG
ID142	MLMPVVGRGNGIPQTVSEWLRLLPFLGVLALLGYLAV
ID143	MDRLGSFSNDPSDKPPCRGCSSYLMEFYKCAECGPPFFLCLQCFTRG
ID144	MSDVNVSAIPKKNSGHIYNKNISQKDCDCLHVVEPMPVRGPDVEAYCLRCECKYEERSS VTIKVTIHYLSILGLLLLYMVYTLV
ID145	MAXCRRCRSQRRSHCCQDRRLRRPRLTLWRHHTALSLSL SMAPPNP
ID146	MTRLGGKGGQFPFGQKIISKDILALTALSVXRKXS
ID147	MKGWGWLALLGALLGTAWA
ID148	MHRLLCLLLLFGGGDP
ID149	MLWRQLIYWQLLALFFLPFCLC
ID150	MRLLLLLLVAASAMVRS
ID151	MSSGXELLWPGAALLVLLGVAASLC
ID152	MTKEIFFFTVELVCENKELCSSPRWRNAIQSNFSKVTSFFMSCHHFKGLAPLPHVYTQG NCRPISCLGLTLMPPASS
ID153	MTDIGCLYFRALCLPRGAWG
ID154	MVPSLVIPDLTCLFLFLNLRWS
ID155	MALRLKLAAATSASA
ID156	MWGNKFGVLLFLYSVLLTKG
ID157	MYTFRKLSPYLNKIVFVCSVLGQSWG
ID158	MESRVLLRTFCLIFGLGAVWG
ID159	MLVLKKHSVNIAAQTCFKFNIFRILIFLGFFLGLFH
ID160	MDVKCPGCYKITTVFSHAQTVVLCVGCSTVLC
ID161	MCILSVLHALPAGIA
ID162	MLVVEASSSVRLASSEVTWSILVTPSASTPIISLSAGPLRTPSHSKTWLLGALEPASE
ID163	MYSFPTTVVEEILSLSLQIAFPTVSC
ID164	MLMLPLRSLALVRE
ID165	MVPLVAVVSGPRAQLFACLLRLGTQ
ID166	MDNRFATAFVIACVLSLIST
ID167	MPEYCGNEVTPTAAQAPEVTYEAEEGSLWTLTSLDGHLL
ID168	MNRVLCAPAAGAVRA
ID169	MAFTFAAFCYMLALLLTAALIFF
ID170	MXXXXEXLLAFHHDCEA
ID171	MGPYNVAVPSDVSHARFYFLFHRPLRLNLLILIEG
ID172	MMNFRQRMGWIGVGLYLLASAAA
ID173	MLFASGGFXVKLYDIEQQIRNALENIRKEMKLLAQAGSLKGSLSVEEQLSLISGCPNIQ EAVEGAMHIQECVPEDLELKKKIFAQLDSIIDESSLKRFXFLSHAFXVVCWLGPCEA
ID174	MQCFLGGLGLCSLPLSPSAVCP
ID175	MSSFLLSFSQSLS
ID176	MLTASLAFQLVDG
ID177	MYXRRELSILCILSAFNFLVCLSLG
ID178	MGLSAMDTSIIVFGVSWVMLVYS
ID179	MYFWRDVAVSLDTLWALPRQQPGLGNRVLGLLSGTNKDYKGQKLAEQMFQGILFSAIV GFIYG
ID180	MHWGKRWXLXXGGLLICXLXIGTATP
ID181	MAXRYNRLTVLAGAXLALGLXTCLSVLFG
ID182	MGFTGFFTATCFISKVFMTCLCRPPISS
ID183	MIMYLFVICVIFEIRNYAFSILIVLLPVLFSLK
ID184	MSTVGLXHFPLTRICPAPWGLRLWEKLTLLSPGIA
ID185	MLALAXHLSTVES
ID186	MLLSIGMLMLXATQVYTILTVQLFAFLNLLPVEA

<u>SEQ. ID</u> <u>NO.</u>	<u>SIGNAL PEPTIDE</u>
ID187	MSTVGLFHFTPLTRICPAPWGLRLWEKLTLLSPGIA
ID188	MELTIFILRLAIYILTFPLYLLNFLGLWSWICK
ID189	MSLLHGKNCVITIRPTGQPLNGDLLLYLCCMINIHH
ID190	MSFNISYFIAFPNLSQA
ID191	MKLKXNVLTILLPVHLLIT
ID192	MAALVTVLFTGVRR
ID193	MASVGECPAPVPVKDKKLEVKLGELPSWILMRDFSPSGIFG
ID194	MLALLVLVTVALASA
ID195	MRIISRQIVLLFSGFWGLAMG

Minimum signal peptide score	false positive rate	false negative rate	proba(0.1)	proba(0.2)
3.5	0.121	0.036	0.467	0.664
4	0.096	0.06	0.519	0.708
4.5	0.078	0.079	0.565	0.745
5	0.062	0.098	0.615	0.782
5.5	0.05	0.127	0.659	0.813
6	0.04	0.163	0.694	0.836
6.5	0.033	0.202	0.725	0.855
7	0.025	0.248	0.763	0.878
7.5	0.021	0.304	0.78	0.889
8	0.015	0.368	0.816	0.909
8.5	0.012	0.418	0.836	0.92
9	0.009	0.512	0.856	0.93
9.5	0.007	0.581	0.863	0.934
10	0.006	0.679	0.835	0.919

TABLE IV

126/2

Minimum signal peptide score	All ESTs	New ESTs	ESTs matching public EST closer than 40 bp from beginning	ESTs extending known mRNA more than 40 bp	ESTs extending public EST more than 40 bp
3.5	2674	947	599	23	150
4	2278	784	499	23	126
4.5	1943	647	425	22	112
5	1657	523	353	21	96
5.5	1417	419	307	19	80
6	1190	340	238	18	68
6.5	1035	280	186	18	60
7	893	219	161	15	48
7.5	753	173	132	12	36
8	636	133	101	11	29
8.5	543	104	83	8	26
9	456	81	63	6	24
9.5	364	57	48	6	18
10	303	47	35	6	15

TABLE V

Tissue	All ESTs	New ESTs	ESTs matching public EST closer than 40 bp from beginning	ESTs extending known mRNA more than 40 bp	ESTs extending public EST more than 40 bp
Brain	329	131	75	3	24
Cancerous prostate	134	40	37	1	6
Cerebellum	17	9	1	0	6
Colon	21	11	4	0	0
Dystrophic muscle	41	18	8	0	1
Fetal brain	70	37	18	0	1
Fetal kidney	227	116	46	1	19
Fetal liver	13	7	2	0	0
Heart	30	15	7	0	1
Hypertrophic prostate	86	23	22	2	2
Kidney	10	7	3	0	0
Large intestine	21	8	4	0	1
Liver	23	9	6	0	0
Lung	24	12	4	0	1
Lung (cells)	57	38	6	0	4
Lymph ganglia	163	60	23	2	12
Lymphocytes	23	6	4	0	2
Muscle	33	16	6	0	4
Normal prostate	181	61	45	7	11
Ovary	90	57	12	1	2
Pancreas	48	11	6	0	1
Placenta	24	5	1	0	0
Prostate	34	16	4	0	2
Spleen	56	28	10	0	1
Substantia nigra	108	47	27	1	6
Surrenals	15	3	3	1	0
Testis	131	68	25	1	8
Thyroid	17	8	2	0	2
Umbilical cord	55	17	12	1	3
Uterus	28	15	3	0	2
Non tissue-specific	568	48	177	2	28
Total	2677	947	601	23	150

TABLE VI

Description of Transcription Factor Binding Sites present on promoters isolated from SignalTag sequences

Promoter sequence P13H2 (546 bp):

Matrix	Position	Orientation	Score	Length	Sequence
CMYB_01	-502	+	0.983	9	TGTCAGTTG
MYOD_Q6	-501	-	0.981	10	CCCAACTGAC
S8_01	-444	-	0.960	11	AATAGAATTAG
S8_01	-425	+	0.968	11	AACTAAATTAG
DELTAEF1_01	-390	-	0.960	11	GCACACCTCAG
GATA_C	-384	-	0.964	11	AGATAAATCCA
CMYB_01	-349	+	0.958	9	CTTCAGTTG
GATA1_02	-343	+	0.959	14	TTGTAGATAGGACA
GATA_C	-339	+	0.953	11	AGATAGGACAT
TAL1ALPHA47_01	-235	+	0.973	18	CATAACAGATGGTAAG
TAL1BETA47_01	-235	+	0.983	18	CATAACAGATGGTAAG
TAL1BETA1F2_01	-235	+	0.978	16	CATAACAGATGGTAAG
MYOD_Q6	-232	-	0.954	10	ACCATCTGTT
GATA1_04	-217	-	0.953	13	TCAAGATAAAGTA
IK1_01	-128	+	0.983	13	AGTTGGGAATTCC
IK2_01	-128	+	0.985	12	AGTTGGGAATTCC
CREL_01	-123	+	0.962	10	TGGGAATTCC
GATA1_02	-96	+	0.950	14	TCAGTGATATGGCA
SRY_02	-41	-	0.951	12	TAAACAAAACA
E2F_02	-33	+	0.957	8	TTTAGCGC
MZF1_01	-5	-	0.975	8	TGAGGGGA

Promoter sequence P15B4 (861bp) :

Matrix	Position	Orientation	Score	Length	Sequence
NFY_Q6	-748	-	0.956	11	GGACCAATCAT
MZF1_01	-738	+	0.962	8	CCTGGGGA
CMYB_01	-684	+	0.994	9	TGACCGTTG
VMYB_02	-682	-	0.985	9	TCCAACGGT
STAT_01	-673	+	0.968	9	TTCTTGGA
STAT_01	-673	-	0.951	9	TTCCAGGAA
MZF1_01	-556	-	0.956	8	TTGGGGGA
IK2_01	-451	+	0.965	12	GAATGGGATTTT
MZF1_01	-424	+	0.986	8	AGAGGGGA
SRY_02	-398	-	0.955	12	GAAAACAAAACA
MZF1_01	-216	+	0.960	8	GAAGGGGA
MYOD_Q6	-190	+	0.981	10	AGCATCTGCC
DELTAEF1_01	-178	+	0.958	11	TCCCACCTTCC
S8_01	5	-	0.992	11	GAGGCAATTAT
MZF1_01	16	-	0.986	8	AGAGGGGA

Promoter sequence P29B6 (555 bp) :

Matrix	Position	Orientation	Score	Length	Sequence
ARNT_01	-311	+	0.964	16	GGA CTCACGTGCTGCT
NMYC_01	-309	+	0.965	12	ACTCACGTGCTG
USF_01	-309	+	0.985	12	ACTCACGTGCTG
USF_01	-309	-	0.985	12	CAGCACGTGAGT
NMYC_01	-309	-	0.958	12	CAGCACGTGAGT
MYCMAX_02	-309	-	0.972	12	CAGCACGTGAGT
USF_C	-307	+	0.997	8	TCACGTGC
USF_C	-307	-	0.991	8	GCACGTGA
MZF1_01	-292	-	0.968	8	CATGGGGA
ELK1_02	-105	+	0.963	14	CTCTCCGGAAGCCT
CETS1P54_01	-102	+	0.974	10	TCCGGAAGCC
AP1_Q4	-42	-	0.963	11	AGTGACTGAAC
AP1FJ_Q2	-42	-	0.961	11	AGTGACTGAAC
PADS_C	45	+	1.000	9	TGTGGTCTC

TABLE VII

CLAIMS

1. A purified or isolated nucleic acid comprising the sequence of one of SEQ ID NOs: 38-195 or comprising a sequence complementary thereto.
- 5 2. The nucleic acid of Claim 1, wherein said nucleic acid is recombinant.
3. A purified or isolated nucleic acid comprising at least 10 consecutive bases of the sequence of one of SEQ ID NOs: 38-195 or one of the sequences complementary thereto.
4. A purified or isolated nucleic acid comprising at least 15 consecutive bases of one of the sequences of SEQ ID NOs: 38-195 or one of the sequences complementary thereto.
- 10 5. The nucleic acid of Claim 4, wherein said nucleic acid is recombinant.
6. A purified or isolated nucleic acid of at least 15 bases capable of hybridizing under stringent conditions to the sequence of one of SEQ ID NOs: 38-195 or one of the sequences complementary to the sequences of SEQ ID NOs: 38-195.
- 15 7. The nucleic acid of Claim 6, wherein said nucleic acid is recombinant.
8. A purified or isolated nucleic acid encoding a human gene product, said human gene product having a sequence partially encoded by one of the sequences of SEQ ID NO: 38-195.
- 20 9. A purified or isolated nucleic acid having the sequence of one of SEQ ID NOs: 38-195 or having a sequence complementary thereto.
10. A purified or isolated nucleic acid comprising the nucleotides of one of SEQ ID NOs: 38-195 which encode a signal peptide.
11. A purified or isolated polypeptides comprising a signal peptide encoded by one of the sequences of SEQ ID NOs: 38-195.
- 25 12. A vector encoding a fusion protein comprising a polypeptide and a signal peptide, said vector comprising a first nucleic acid encoding a signal peptide encoded by one of the sequences of SEQ ID NOs: 38-195 operably linked to a second nucleic acid encoding a polypeptide.
- 30 13. A method of directing the extracellular secretion of a polypeptide or the insertion of a polypeptide into the membrane comprising the steps of:

obtaining a vector according to Claim 12; and

introducing said vector into a host cell such that said fusion protein is secreted into the extracellular environment of said host cell or inserted into the membrane of said host cell.

14. A method of importing a polypeptide into a cell comprising contacting said
5 cell with a fusion protein comprising a signal peptide encoded by one of the sequences of SEQ ID NOs: 38-195 operably linked to said polypeptide.

15. A method of making a cDNA encoding a human secretory protein that is partially encoded by one of SEQ ID NOs 38-195, comprising the steps of:

obtaining a cDNA comprising one of the sequences of SEQ ID NOs: 38-195;

10 contacting said cDNA with a detectable probe comprising at least 15 consecutive nucleotides of said sequence of SEQ ID NO: 38-195 or a sequence complementary thereto under conditions which permit said probe to hybridize to said cDNA;

identifying a cDNA which hybridizes to said detectable probe; and

isolating said cDNA which hybridizes to said probe.

16. An isolated or purified cDNA encoding a human secretory protein, said
15 human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 15.

17. The cDNA of Claim 16 wherein said cDNA comprises the full protein coding
20 sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

18. A method of making a cDNA comprising one of the sequences of SEQ ID NOs: 38-195, comprising the steps of:

contacting a collection of mRNA molecules from human cells with a first primer capable of hybridizing to the polyA tail of said mRNA;

25 hybridizing said first primer to said polyA tail;

reverse transcribing said mRNA to make a first cDNA strand;

making a second cDNA strand complementary to said first cDNA strand using at least one primer comprising at least 15 nucleotides of one of the sequences of SEQ ID NOs 38-195; and

30 isolating the resulting cDNA comprising said first cDNA strand and said second cDNA strand.

19. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 18.

5 20. The cDNA of Claim 19 wherein said cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

21. The method of Claim 18, wherein the second cDNA strand is made by:
contacting said first cDNA strand with a first pair of primers, said first pair of primers comprising a second primer comprising at least 15 consecutive nucleotides of one of the
10 sequences of SEQ ID NOs 38-195 and a third primer having a sequence therein which is included within the sequence of said first primer;

performing a first polymerase chain reaction with said first pair of nested primers to generate a first PCR product;

contacting said first PCR product with a second pair of primers, said second pair of
15 primers comprising a fourth primer, said fourth primer comprising at least 15 consecutive nucleotides of said sequence of one of SEQ ID NO:s 38-195 , and a fifth primer, said fourth and fifth primers being capable of hybridizing to sequences within said first PCR product; and
performing a second polymerase chain reaction, thereby generating a second PCR product.

20 22. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-195, or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 21.

23. The cDNA of Claim 22 wherein said cDNA comprises the full protein coding
25 sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

24. The method of Claim 18 wherein the second cDNA strand is made by:
contacting said first cDNA strand with a second primer comprising at least 15
consecutive nucleotides of the sequences of SEQ ID NOs: 38-195;

hybridizing said second primer to said first strand cDNA; and
30 extending said hybridized second primer to generate said second cDNA strand.

25. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein partially encoded by one of SEQ ID NOs 38-195 or comprising a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 24.

5 26. The cDNA of Claim 25, wherein said cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-195.

27. A method of making a protein comprising one of the sequences of SEQ ID NO: 196-353, comprising the steps of:

obtaining a cDNA encoding the full protein sequence partially included in one of the sequences of sequence of SEQ ID NO: 38-195;

10 inserting said cDNA in an expression vector such that said cDNA is operably linked to a promoter;

introducing said expression vector into a host cell whereby said host cell produces the protein encoded by said cDNA; and

15 isolating said protein.

28. An isolated protein obtainable by the method of Claim 27.

29. A method of obtaining a promoter DNA comprising the steps of:

obtaining DNAs located upstream of the nucleic acids of SEQ ID NO: 38-195 or the sequences complementary thereto;

20 screening said upstream DNAs to identify a promoter capable of directing transcription initiation; and

isolating said DNA comprising said identified promoter.

30. The method of Claim 29, wherein said obtaining step comprises chromosome walking from said nucleic acids of SEQ ID NO: 38-195 or sequences complementary thereto.

25 31. The method of Claim 30, wherein said screening step comprises inserting said upstream sequences into a promoter reporter vector.

32. The method of Claim 30, wherein said screening step comprises identifying motifs in said upstream DNAs which are transcription factor binding sites or transcription start sites.

30 33. An isolated promoter obtainable by the method of Claim 32.

34. An isolated or purified protein comprising one of the sequences of SEQ ID NO: 196-353.

35. In an array of discrete EST's or fragments thereof of at least 15 nucleotides in length, the improvement comprising inclusion in said array of at least one of the sequences of
5 SEQ ID NOs: 38-195, or one of the sequences complementary to the sequences of SEQ ID NOs: 38-195, or a fragment thereof of at least 15 consecutive nucleotides.

36. The array of Claim 35 including therein at least two of the sequences of SEQ ID NOs: 38-195, the sequences complementary to the sequences of SEQ ID NOs: 38-195, or fragments thereof of at least 15 consecutive nucleotides.

10 37. The array of Claim 35 including therein at least five of the sequences of SEQ ID NOs: 38-195, the sequences complementary to the sequences of SEQ ID NOs: 38-195, or fragments thereof of at least 15 consecutive nucleotides.

1/4




Figure 1




Figure 2





Figure 3

Promoter P13H2



Promoter P15B4



Promoter P29B6




Figure 4

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME : GENSET SA
- (B) STREET :24, RUE ROYALE
- (C) CITY: PARIS
- (E) COUNTRY : FRANCE
- (F) POSTAL CODE (ZIP) : 75008

(ii) TITLE OF INVENTION: 5' ESTs FOR SECRETED PROTEINS
IDENTIFIED FROM BRAIN TISSUE

(iii) NUMBER OF SEQUENCES: 353

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy Disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: Win95
- (D) SOFTWARE: Word

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(ix) FEATURE:

- (A) NAME/KEY: Cap
- (B) LOCATION: 1
- (D) OTHER INFORMATION: m7Gppp added to 1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGCAUCCUAC UCCCAUCCAA UUCCACCCUA ACUCCUCCCA UCUCCAC

47

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GCAUCCUACU CCAUCCAAU UCCACCCUAA CUCCUCCCAU CUCCAC

46

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATCAAGAATT CGCAGGAGAC CATTA

25

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

TAATGGTCTC GTGCGAATTC TTGAT

25

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CCGACAAGAC CAACGTCAAG GCCGC

25

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID

- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TCACCAGCAG GCAGTGGCTT AGGAG

25

(2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

AGTGATTTCCT GCTACTTTGG ATGGC

25

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCTTGGTCTT GTTCTGGAGT TTAGA

25

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TCCAGAATGG GAGACAAGCC AATTT

25

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

AGGGAGGAGG AACAGCGTG AGTCC

25

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ATGGGAAAGG AAAAGACTCA TATCA

25

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

AGCAGCAACA ATCAGGACAG CACAG

25

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

ATCAAGAATT CGCACGAGAC CATTA

25

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 67 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

ATCGTTGAGA CTCGTACCAG CAGAGTCACG AGAGAGACTA CACGGTACTG GTTTTTTTTT

60

TTTTTVN

67

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CCAGCAGAGT CACGAGAGAG ACTACACGG

29

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: SINGLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

CACGAGAGAG ACTACACGGT ACTGG

25

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 526 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Lymph ganglia

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(261..376)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 166..281
id N70479
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(380..486)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 54..160
id N70479
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(110..145)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 403..438
id N70479
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(196..229)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 315..348
id N70479
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 90..140
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.2
seq LLLITAILAVAVG/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

```

AATATRARAC AGCTACAATA TTCCAGGGCC ARTCACTTGC CATTCTCAT AACAGCGTCA      60
GAGAGAAAGA ACTGACTGAR ACGTTTGAG ATG AAG AAA GTT CTC CTC CTG ATC      113
                               Met Lys Lys Val Leu Leu Leu Ile
                               -15                               -10
ACA GCC ATC TTG GCA GTG GCT GTW GGT TTC CCA GTC TCT CAA GAC CAG      161
Thr Ala Ile Leu Ala Val Ala Val Gly Phe Pro Val Ser Gln Asp Gln
                               -5                               1                               5
GAA CGA GAA AAA AGA AGT ATC AGT GAC AGC GAT GAA TTA GCT TCA GGR      209
Glu Arg Glu Lys Arg Ser Ile Ser Asp Ser Asp Glu Leu Ala Ser Gly
                               10                               15                               20
WTT TTT GTG TTC CCT TAC CCA TAT CCA TTT CGC CCA CTT CCA CCA ATT      257
Xaa Phe Val Phe Pro Tyr Pro Tyr Pro Phe Arg Pro Leu Pro Pro Ile
                               25                               30                               35
CCA TTT CCA AGA TTT CCA TGG TTT AGA CGT AAN TTT CCT ATT CCA ATA      305
Pro Phe Pro Arg Phe Pro Trp Phe Arg Arg Xaa Phe Pro Ile Pro Ile
40                               45                               50                               55
CCT GAA TCT GCC CCT ACA ACT CCC CTT CCT AGC GAA AAG TAAACAARAA      354
Pro Glu Ser Ala Pro Thr Thr Pro Leu Pro Ser Glu Lys
                               60                               65
GGAAAAGTCA CRATAACCT GGTCACCTGA AATTGAAATT GAGCCACTTC CTTGAARAAT      414
CAAAATTCCT GTTAATAAAA RAAAAACAAA TGTAATTGAA ATAGCACACA GCATTCTCTA      474
GTCAATATCT TTAGTGATCT TCTTTAATAA ACATGAAAGC AAAAAAAAAA AA      526

```

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..17
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.2
seq LLLITAILAVAVG/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

```

Met Lys Lys Val Leu Leu Leu Ile Thr Ala Ile Leu Ala Val Ala Val
  1                               5                               10                               15

```

Gly

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 822 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: kidney

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 260..464
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 153..357
id H57434
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 118..184
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 98..164
id H57434
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 56..113
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 35..92
id H57434
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 454..485
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 348..379
id H57434
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 118..545
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..428
id N27248
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 65..369
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 41..345
id H94779
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 61..399
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 6..344
id H09880
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 408..458
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 355..405
id H09880
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 60..399
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 56..395
id H29351
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 393..432
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 391..430
id H29351
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 346..408
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.5
seq SFLPSALVIWTS/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

ACTCCTTTTA GCATAGGGGC TTCGGCGCCA GCGGCCAGCG CTAGTCGGTC TGGTAAGTGC	60
CTGATGCCGA GTTCCGTCTC TCGCGTCTTT TCCTGGTCCC AGGCAAAGCG GASGNAGATC	120
CTCAAACGGC CTAGTGCTTC GCGCTTCCGG AGAAAATCAG CGGTCTAATT AATTCCTCTG	180

```

GTTTGTGAA GCAGTTACCA AGAATCTTCA ACCCTTTCCC ACAAAAGCTA ATTGAGTACA      240
CGTTCCTGTT GAGTACACGT TCCTGTTGAT TTACAAAAGG TGCAGGTATG AGCAGGTCTG      300
AAGACTAACA TTTTGTGAAG TTGTAAAACA GAAAACCTGT TAGAA ATG TGG TGG TTT      357
                                   Met Trp Trp Phe
                                   -20

CAG CAA GGC CTC AGT TTC CTT CCT TCA GCC CTT GTA ATT TGG ACA TCT      405
Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu Val Ile Trp Thr Ser
      -15                      -10                      -5

GCT GCT TTC ATA TTT TCA TAC ATT ACT GCA GTA ACA CTC CAC CAT ATA      453
Ala Ala Phe Ile Phe Ser Tyr Ile Thr Ala Val Thr Leu His His Ile
      1                      5                      10                      15

GAC CCG GCT TTA CCT TAT ATC AGT GAC ACT GGT ACA GTA GCT CCA RAA      501
Asp Pro Ala Leu Pro Tyr Ile Ser Asp Thr Gly Thr Val Ala Pro Xaa
      20                      25                      30

AAA TGC TTA TTT GGG GCA ATG CTA AAT ATT GCG GCA GTT TTA TGT CAA      549
Lys Cys Leu Phe Gly Ala Met Leu Asn Ile Ala Ala Val Leu Cys Gln
      35                      40                      45

AAA TAGAAATCAG GAARATAATT CAACTTAAAG AAKTTCATTT CATGACCAAA      602
Lys

CTCTTCARAA ACATGTCTTT ACAAGCATAT CTCTTGTATT GCTTTCTACA CTGTTGAATT      662
GTCTGGCAAT ATTTCTGCAG TGGAAAATTT GATTARMTA GTTCTTGACT GATAAATATG      722
GTAAGGTGGG CTTTCCCCC TGTGTAATTG GCTACTATGT CTTACTGAGC CAAGTTGTAW      782
TTTGAAATAA AATGATATGA GAGTGACACA AAAAAAAAAA      822

```

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..21
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5
seq SFLPSALVIWTS/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Met Trp Trp Phe Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu Val
 1 5 10 15
 Ile Trp Thr Ser Ala
 20

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 405 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(103..398)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
 region 1..296
 id AA442893
 est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 185..295
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9
 seq LSYASSALSPCLT/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

ATCACCTTCT TCTCCATCCT TSTCTGGGCC AGTCCCCARC CCAGTCCCTC TCCTGACCTG 60
 CCCAGCCCAA GTCAGCCTTC AGCACGCGCT TTTCTGCACA CAGATATTCC AGGCCTACCT 120
 GGCATTCCAG GACCTCCGMA ATGATGCTCC AGTCCCTTAC AAGCGCTTCC TGGATGAGGG 180
 TGGC ATG GTG CTG ACC ACC CTC CCC TTG CCC TCT GCC AAC AGC CCT GTG 229
 Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val
 -35 -30 -25
 AAC ATG CCC ACC ACT GGC CCC AAC AGC CTG AGT TAT GCT AGC TCT GCC 277
 Asn Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala
 -20 -15 -10
 CTG TCC CCC TGT CTG ACC GCT CCA AAK TCC CCC CGG CTT GCT ATG ATG 325
 Leu Ser Pro Cys Leu Thr Ala Pro Xaa Ser Pro Arg Leu Ala Met Met
 -5 1 5 10
 CCT GAC AAC TAAATATCCT TATCCAAATC AATAAARWRA RAATCCTCCC TCCARAAGGG 384
 Pro Asp Asn

TTTCTAAAAA CAAAAAAA A

405

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..37
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9
seq LSYASSALSPCLT/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val Asn
1 5 10 15
Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala Leu
20 25 30
Ser Pro Cys Leu Thr
35

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cancerous prostate

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..331
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..183
id AA397994
est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 328..485
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 179..336
 id AA397994
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: complement(182..496)
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 14..328
 id AA399680
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 196..240
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.5
 seq ILSTVTALTFXA/LD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

AAAAAATTGG TCCCAGTTTT CACCCTGCCG CAGGGCTGGC TGGGGAGGGC AGCGGTTTAG	60
ATTAGCCGTG GCCTAGGCCG TTTAACGGGG TGACACGAGC NTGCAGGGCC GAGTCCAAGG	120
CCCGGAGATA GGACCAACCG TCAGGAATGC GAGGAATGTT TTTCTTCGGA CTCTATCGAG	180
GCACACAGAC AGACC ATG GGG ATT CTG TCT ACA GTG ACA GCC TTA ACA TTT	231
Met Gly Ile Leu Ser Thr Val Thr Ala Leu Thr Phe	
-15 -10 -5	
GCC ARA GCC CTG GAC GGC TGC AGA AAT GGC ATT GCC CAC CCT GCA AGT	279
Ala Xaa Ala Leu Asp Gly Cys Arg Asn Gly Ile Ala His Pro Ala Ser	
1 5 10	
GAG AAG CAC AGA CTC GAG AAA TGT AGG GAA CTC GAG ASC ASC CAC TCG	327
Glu Lys His Arg Leu Glu Lys Cys Arg Glu Leu Glu Xaa Xaa His Ser	
15 20 25	
GCC CCA GGA TCA ACC CAS CAC CGA AGA AAA ACA ACC AGA AGA AAT TAT	375
Ala Pro Gly Ser Thr Xaa His Arg Arg Lys Thr Thr Arg Arg Asn Tyr	
30 35 40 45	
TCT TCA GCC TGAAATGAAK CCGGGATCAA ATGGTTGCTG ATCARAGCCC ATATTTAAAT	434
Ser Ser Ala	
TGGAAAAGTC AAATTGASCA TTATTAAATA AAGCTTGTTT AATATGTCTC AAACAAAAAA	494
AA	496

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..15
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5
seq ILSTVTALTFXA/LD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

```

Met Gly Ile Leu Ser Thr Val Thr Ala Leu Thr Phe Ala Xaa Ala
 1             5             10             15

```

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 623 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 49..96
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.1
seq LVLTLCTLPLAVA/SA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

```

AAAGATCCCT GCAGCCCGGC AGGAGAGAAG GCTGAGCCTT CTGGCGTC ATG GAG AGG      57
                                     Met Glu Arg
                                     -15

CTC GTC CTA ACC CTG TGC ACC CTC CCG CTG GCT GTG GCG TCT GCT GGC      105
Leu Val Leu Thr Leu Cys Thr Leu Pro Leu Ala Val Ala Ser Ala Gly
      -10             -5             1

TGC GCC ACG ACG CCA GCT CGC AAC CTG AGC TGC TAC CAG TGC TTC AAG      153
Cys Ala Thr Thr Pro Ala Arg Asn Leu Ser Cys Tyr Gln Cys Phe Lys

```

5	10	15	
GTC AGC AGC TGG ACG GAG TGC CCG CCC ACC TGG TGC AGC CCG CTG GAC Val Ser Ser Trp Thr Glu Cys Pro Pro Thr Trp Cys Ser Pro Leu Asp 20 25 30 35			201
CAA GTC TGC ATC TCC AAC GAG GTG GTC GTC TCT TTT AAA TGG AGT GTA Gln Val Cys Ile Ser Asn Glu Val Val Val Ser Phe Lys Trp Ser Val 40 45 50			249
CGC GTC CTG CTC AGC AAA CGC TGT GCT CCC AGA TGT CCC AAC GAC AAC Arg Val Leu Leu Ser Lys Arg Cys Ala Pro Arg Cys Pro Asn Asp Asn 55 60 65			297
ATG AAK TTC GAA TGG TCG CCG GCC CCC ATG GTG CAA GGC GTG ATC ACC Met Xaa Phe Glu Trp Ser Pro Ala Pro Met Val Gln Gly Val Ile Thr 70 75 80			345
AGG CGC TGC TGT TCC TGG GCT CTC TGC AAC AGG GCA CTG ACC CCA CAG Arg Arg Cys Cys Ser Trp Ala Leu Cys Asn Arg Ala Leu Thr Pro Gln 85 90 95			393
GAG GGG CGC TGG GCC CTG CRA GGG GGG CTC CTG CTC CAG GAC CCT TCG Glu Gly Arg Trp Ala Leu Xaa Gly Gly Leu Leu Leu Gln Asp Pro Ser 100 105 110 115			441
AGG GGC ARA AAA ACC TGG GTG CGG CCA CAG CTG GGG CTC CCA CTC TGC Arg Gly Xaa Lys Thr Trp Val Arg Pro Gln Leu Gly Leu Pro Leu Cys 120 125 130			489
CTT CCC AWT TCC AAC CCC CTC TGC CCA RGG GAA ACC CAG GAA GGA Leu Pro Xaa Ser Asn Pro Leu Cys Pro Xaa Glu Thr Gln Glu Gly 135 140 145			534
TAACACTGTG GGTGCCCCCA CCTGTGCATT GGGACCACRA CTTACCCCTC TTGGARACAA			594
TAAACTCTCA TGCCCCCAAA AAAAAAAAAA			623

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..16
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.1
seq LVLTLCTLPLAVA/SA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met Glu Arg Leu Val Leu Thr Leu Cys Thr Leu Pro Leu Ala Val Ala
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 848 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: kidney

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 32..73
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.7
seq LWLLFFLVTAIHA/EL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

```

AACTTTGCCT TGTGTTTTCC ACCCTGAAAG A ATG TTG TGG CTG CTC TTT TTT CTG    55
                               Met Leu Trp Leu Leu Phe Phe Leu
                               -10

GTG ACT GCC ATT CAT GCT GAA CTC TGT CAA CCA GGT GCA GAA AAT GCT    103
Val Thr Ala Ile His Ala Glu Leu Cys Gln Pro Gly Ala Glu Asn Ala
-5 1 5 10

TTT AAA GTG AGA CTT AGT ATC AGA ACA GCT CTG GGA GAT AAA GCA TAT    151
Phe Lys Val Arg Leu Ser Ile Arg Thr Ala Leu Gly Asp Lys Ala Tyr
15 20 25

GCC TGG GAT ACC AAT GAA GAA TAC CTC TTC AAA GCG ATG GTA GCT TTC    199
Ala Trp Asp Thr Asn Glu Glu Tyr Leu Phe Lys Ala Met Val Ala Phe
30 35 40

TCC ATG AGA AAA GTT CCC AAC AGA GAA GCA ACA GAA ATT TCC CAT GTC    247
Ser Met Arg Lys Val Pro Asn Arg Glu Ala Thr Glu Ile Ser His Val
45 50 55

CTA CTT TGC AAT GTA ACC CAG AGG GTA TCA TTC TGG TTT GTG GTT ACA    295
Leu Leu Cys Asn Val Thr Gln Arg Val Ser Phe Trp Phe Val Val Thr
60 65 70

GAC CCT TCA AAA AAT CAC ACC CTT CCT GCT GTT GAG GTG CAA TCA GCC    343
Asp Pro Ser Lys Asn His Thr Leu Pro Ala Val Glu Val Gln Ser Ala
75 80 85 90

ATA AGA ATG AAC AAG AAC CGG ATC AAC AAT GCC TTC TTT CTA AAT GAC    391
Ile Arg Met Asn Lys Asn Arg Ile Asn Asn Ala Phe Phe Leu Asn Asp

```

95	100	105	
CAA ACT CTG GAA TTT TTA AAA ATC CCT TCC ACA CTT GCA CCA CCC ATG			439
Gln Thr Leu Glu Phe Leu Lys Ile Pro Ser Thr Leu Ala Pro Pro Met			
110	115	120	
GAC CCA TCT GTG CCC ATC TGG ATT ATT ATA TTT GGT GTG ATA TTT TGC			487
Asp Pro Ser Val Pro Ile Trp Ile Ile Phe Gly Val Ile Phe Cys			
125	130	135	
ATC ATC ATA GTT GCA ATT GCA CTA CTG ATT TTA TCA GGG ATC TGG CAA			535
Ile Ile Ile Val Ala Ile Ala Leu Leu Ile Leu Ser Gly Ile Trp Gln			
140	145	150	
CGT ADA ARA AAG AAC AAA GAA CCA TCT GAA GTG GAT GAC GCT GAA RAT			583
Arg Xaa Xaa Lys Asn Lys Glu Pro Ser Glu Val Asp Asp Ala Glu Xaa			
155	160	165	170
AAK TGT GAA AAC ATG ATC ACA ATT GAA AAT GGC ATC CCC TCT GAT CCC			631
Xaa Cys Glu Asn Met Ile Thr Ile Glu Asn Gly Ile Pro Ser Asp Pro			
175	180	185	
CTG GAC ATG AAG GGA GGG CAT ATT AAT GAT GCC TTC ATG ACA GAG GAT			679
Leu Asp Met Lys Gly Gly His Ile Asn Asp Ala Phe Met Thr Glu Asp			
190	195	200	
GAG AGG CTC ACC CCT CTC TGAAGGGCTG TTGTTCTGCT TCCTCAARAA			727
Glu Arg Leu Thr Pro Leu			
205			
ATTAAACATT TGTTTCTGTG TGA CTGCTGA GCATCCTGAA ATACCAAGAG CAGATCATAT			787
TTTTTGTTC ACCATTCTTC TTTTGTAATA AATTTGAAT GTGCTTGAAA AAAAAAAAAA			847
C			848

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 1..14
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.7
seq LWLLFFLVTAIHA/EL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Met Leu Trp Leu Leu Phe Phe Leu Val Thr Ala Ile His Ala

1

5

10

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

GGGAAGATGG AGATAGTATT GCCTG

25

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CTGCCATGTA CATGATAGAG AGATTC

26

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 546 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Genomic DNA

(ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..517

(ix) FEATURE:

- (A) NAME/KEY: transcription start site
- (B) LOCATION: 518

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 17..25
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name CMYB_01

score 0.983
sequence TGTCAGTTG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(18..27)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MYOD_Q6
score 0.961
sequence CCCAACTGAC

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(75..85)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name S8_01
score 0.960
sequence AATAGAATTAG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 94..104
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name S8_01
score 0.966
sequence AACTAAATTAG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(129..139)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name DELTAEF1_01
score 0.960
sequence GCACACCTCAG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(155..165)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name GATA_C
score 0.964
sequence AGATAAATCCA

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 170..178
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name CMYB_01
score 0.958
sequence CTTCAGTTG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 176..189
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name GATA1_02
score 0.959
sequence TTGTAGATAGGACA

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site

(B) LOCATION: 180..190
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name GATA_C
score 0.953
sequence AGATAGGACAT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 284..299
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name TAL1ALPHA47_01
score 0.973
sequence CATAACAGATGGTAAG

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 284..299
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name TAL1BETA47_01
score 0.983
sequence CATAACAGATGGTAAG

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 284..299
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name TAL1BETA1TF2_01
score 0.978
sequence CATAACAGATGGTAAG

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(287..296)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name MYOD_Q6
score 0.954
sequence ACCATCTGTT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(302..314)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name GATA1_04
score 0.953
sequence TCAAGATAAAGTA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 393..405
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name IK1_01
score 0.963
sequence AGTTGGGAATTCC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 393..404
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name IK2_01
score 0.985
sequence AGTTGGGAATTC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
 (B) LOCATION: 396..405
 (C) IDENTIFICATION METHOD: matinspector prediction
 (D) OTHER INFORMATION: name CREL_01
 score 0.962
 sequence TGGGAATTCC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
 (B) LOCATION: 423..436
 (C) IDENTIFICATION METHOD: matinspector prediction
 (D) OTHER INFORMATION: name GATA1_02
 score 0.950
 sequence TCAGTGATATGGCA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
 (B) LOCATION: complement(478..489)
 (C) IDENTIFICATION METHOD: matinspector prediction
 (D) OTHER INFORMATION: name SRY_02
 score 0.951
 sequence TAAACAAAACA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
 (B) LOCATION: 486..493
 (C) IDENTIFICATION METHOD: matinspector prediction
 (D) OTHER INFORMATION: name E2F_02
 score 0.957
 sequence TTTAGCGC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
 (B) LOCATION: complement(514..521)
 (C) IDENTIFICATION METHOD: matinspector prediction
 (D) OTHER INFORMATION: name MZF1_01
 score 0.975
 sequence TGAGGGGA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

```

TGAGTGCAGT GTTACATGTC AGTTGGGTTA AGTTTGTTAA TGTCATTCAA ATCTTCTATG   60
TCTTGATTG CTGCTAATT CTATTATTTT TGGAACTAAA TTAGTTTGAT GGTCTATTA   120
GTTATTGACT GAGGTGTGCT AATCTCCCAT TATGTGGATT TATCTATTTT TTCAGTTGTA   180
GATAGGACAT TGATAGATAC ATAAGTACCA GGACAAAAGC AGGGAGATCT TTTTCCAAA   240
ATCAGGAGAA AAAAATGACA TCTGGAAAAC CTATAGGGAA AGGCATAACA GATGGTAAGG   300
ATACTTTATC TTGAGTAGGA GAGCCTTCCT GTGGCAACGT GGAGAAGGGA AGAGGTCGTA   360
GAATTGAGGA GTCAGCTCAG TTAGAAGCAG GGAGTTGGGA ATTCCGTTCA TGTGATTTAG   420
CATCAGTGAT ATGGCAAATG TGGGACTAAG GGTAGTGATC AGAGGGTTAA AATTGTGTGT   480
TTTGTTTTAG CGCTGCTGGG GCATCGCCTT GGGTCCCCTC AACAGATTG CCATGAATCT   540

```

CTTCAT

546

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GTACCAGGGA CTGTGACCAT TGC

23

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

CTGTGACCAT TGCTCCAAG AGAG

24

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 861 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Genomic DNA

(ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..806

(ix) FEATURE:

- (A) NAME/KEY: transcription start site
- (B) LOCATION: 807

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(60..70)
- (C) IDENTIFICATION METHOD: matinspector prediction

(D) OTHER INFORMATION: name NFY_Q6
score 0.956
sequence GGACCAATCAT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 70..77
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name MZF1_01
score 0.962
sequence CCTGGGGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 124..132
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name CMYB_01
score 0.994
sequence TGACCGTTG

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(126..134)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name VMYB_02
score 0.985
sequence TCCAACGGT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 135..143
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name STAT_01
score 0.968
sequence TTCCTGGAA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(135..143)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name STAT_01
score 0.951
sequence TTCCAGGAA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(252..259)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name MZF1_01
score 0.956
sequence TTGGGGGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 357..368
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name IK2_01
score 0.965
sequence GAATGGGATTC

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 384..391
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MZF1_01
score 0.986
sequence AGAGGGGA

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(410..421)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name SRY_02
score 0.955
sequence GAAACAAAACA

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 592..599
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MZF1_01
score 0.960
sequence GAAGGGGA

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 618..627
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MYOD_Q6
score 0.981
sequence AGCATCTGCC

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 632..642
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name DELTAEF1_01
score 0.958
sequence TCCCACCTTC

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(813..823)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name S8_01
score 0.992
sequence GAGGCAATTAT

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(824..831)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MZF1_01
score 0.986
sequence AGAGGGGA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

TACTATAGGG CACGCGTGGT CGACGGCCGG GCTGTTCTGG AGCAGAGGGC ATGTCAGTAA 60

TGATTGGTCC CTGGGAAGG TCTGGCTGGC TCCAGCACAG TGAGGCATTT AGGTATCTCT 120

CGGTGACCGT TGGATTCTCTG GAAGCAGTAG CTGTTCTGTT TGGATCTGGT AGGGACAGGG 180
CTCAGAGGGC TAGGCACGAG GGAAGGTCAG AGGAGAAGGS AGGSARGGCC CAGTGAGARG 240
GGAGCATGCC TTCCCCAAC CCTGGCTTSC YCTTGYMAM AGGGCGKTTY TGGGMACTTR 300
AAYTCAGGGC CCAASCAGAA SCACAGGCCC AKTCNTGGCT SMAAGCACAA TAGCCTGAAT 360
GGGATTTTCTAG GTTAGNCAGG GTGAGAGGGG AGGCTCTCTG GCTTAGTTTT GTTTTGTGTTT 420
CCAAATCAAG GTAACCTGCT CCCTTCTGCT ACGGGCCTTG GTCTTGCTT GTCCTCACCC 480
AGTCGGAACCT CCCTACCACT TTCAGGAGAG TGGTTTTAGG CCCGTGGGGC TGTCTCTGTTT 540
CAAGCAGTGT GAGAACATGG CTGGTAGAGG CTCTAGCTGT GTGCGGGGCC TGAAGGGGAG 600
TGGGTTCTCG CCCAAAGAGC ATCTGCCCCTT TTCCCACCTT CCCTTCTCCC ACCAGAAGCT 660
TGCTTGAGCT GTTTGGACAA AAATCCAAAC CCCACTTGGC TACTCTGGCC TGGCTTCAGC 720
TTGGAACCCA ATACCTAGGC TTACAGGCCA TCCTGAGCCA GGGGCCTCTG GAAATTCTCT 780
TCCTGATGGT CCTTTAGGTT TGGGCACAAA ATATAATTGC CTCTCCCCTC TCCCATTTTC 840
TCTCTTGGGA GCAATGGTCA C 861

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

CTGGGATGGA AGGCACGGTA

20

(2) INFORMATION FOR SEQ ID NO: 36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

GAGACCACAC AGCTAGACAA

20

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 555 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Genomic DNA

(ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..500

(ix) FEATURE:

- (A) NAME/KEY: transcription start site
- (B) LOCATION: 501

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 191..206
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name ARNT_01
score 0.964
sequence GGACTCACGTGCTGCT

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 193..204
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name NMYC_01
score 0.965
sequence ACTCACGTGCTG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 193..204
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name USF_01
score 0.985
sequence ACTCACGTGCTG

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(193..204)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name USF_01
score 0.985
sequence CAGCACGTGAGT

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(193..204)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name NMYC_01
score 0.956
sequence CAGCACGTGAGT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(193..204)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name MYCMAX_02
score 0.972
sequence CAGCACGTGAGT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 195..202
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name USF_C
score 0.997
sequence TCACGTGC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(195..202)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name USF_C
score 0.991
sequence GCACGTGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(210..217)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name MZF1_01
score 0.968
sequence CATGGGGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 397..410
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name ELK1_02
score 0.963
sequence CTCTCCGGAAGCCT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: 400..409
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name CETS1P54_01
score 0.974
sequence TCCGGAAGCC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(460..470)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name AP1_Q4
score 0.963
sequence AGTGACTGAAC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site
(B) LOCATION: complement(460..470)
(C) IDENTIFICATION METHOD: matinspector prediction
(D) OTHER INFORMATION: name AP1FJ_Q2
score 0.961

sequence AGTGACTGAAC

(ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 547..555
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name PADS_C
score 1.000
sequence TGTGGTCTC

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

```
CTATAGGGCA CGCKTGGTCG ACGGCCCCGGG CTGGTCTGGT CTGKTGTGGA GTCGGGTTGA    60
AGGACAGCAT TTGKACATC TGGTCTACTG CACCTTCCCT CTGCCGTGCA CTTGGCCTTT    120
KAWAAGCTCA GCACCGGTGC CCATCACAGG GCCGGCAGCA CACACATCCC ATTACTCAGA    180
AGGAACTGAC GGACTCACGT GCTGCTCCGT CCCCATGAGC TCAGTGGACC TGTCTATGTA    240
GAGCAGTCAG ACAGTGCCTG GGATAGAGTG AGAGTTCAGC CAGTAAATCC AAGTGATTGT    300
CATTCCTGTC TGCATTAGTA ACTCCCAACC TAGATGTGAA AACTTAGTTC TTTCTCATAG    360
GTTGCTCTGC CCATGGTCCC ACTGCAGACC CAGGCACTCT CCGGAAGCCT GGAAATCACC    420
CGTGTCTTCT GCCTGCTCCC GCTCACATCC CACACTTGTG TTCAGTCACT GAGTTACAGA    480
TTTTGCCTCC TCAATTTCTC TTGTCTTAGT CCCATCCTCT GTTCCCCTGG CCAGTTTGTC    540
TAGCTGTGTG GTCTC                                         555
```

(2) INFORMATION FOR SEQ ID NO: 38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 195 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 109..171
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 14.3
seq LLLCAVLLSLASA/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

```
AGAGACTAGA GCAGGAAGAG CAGCGGCGAG GCGGCGGTGG TGGCTGAGTC CGTGGTGGA    60
```

GAGGCGAAGG CGACAGCTCT AGGGGTTGGC ACCGGCCCCG AGAGGAGG ATG CGG GTC 117
 Met Arg Val
 -20

CGG ATA GGG CTG ACG CTG CTG CTG TGT GCG GTG CTG CTG AGC TTG GCC 165
 Arg Ile Gly Leu Thr Leu Leu Leu Cys Ala Val Leu Leu Ser Leu Ala
 -15 -10 -5

TCG GCG TCC TCG GAT GAA GAA GGC AAT GGG 195
 Ser Ala Ser Ser Asp Glu Glu Gly Asn Gly
 1 5

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 55..138
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 11.1
seq GLLFLLLLLLMLLA/DP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

AAGAGCCCAG AGAGCTGAAC CTGCATCCCG GACCTGCGGC GACCGTCGTA CACC ATG 57
 Met

GGC CTC CAC CTC CGC CCC TAC CGT GTG GGG CTG CTC CCG GAT GGC CTC 105
 Gly Leu His Leu Arg Pro Tyr Arg Val Gly Leu Leu Pro Asp Gly Leu
 -25 -20 -15

CTG TTC CTC TTG CTG CTG CTA ATG CTG CTC GCG GAC CCA GCG CTC CCG 153
 Leu Phe Leu Leu Leu Leu Leu Met Leu Leu Ala Asp Pro Ala Leu Pro
 -10 -5 1 5

GCC GCT AGG 162
 Ala Ala Arg

(2) INFORMATION FOR SEQ ID NO: 40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 421 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 20..82
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 10.6
seq LLLGAVSWPPASA/SG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

```

AGCGTGGCCG CCCGGGGCC ATG GCG ACA CTC AGC TTC GTC TTC CTG CTG CTG      52
      Met Ala Thr Leu Ser Phe Val Phe Leu Leu Leu
      -20                               -15

GGG GCA GTG TCC TGG CCT CCG GCT TCT GCC TCC GGC CAG GAG TTC TGG      100
Gly Ala Val Ser Trp Pro Pro Ala Ser Ala Ser Gly Gln Glu Phe Trp
-10              -5              1              5

CCC GGA CAA TCG GCG GCC GAT ATT CTG TCG GGG GCG GCT TCC CGC AGA      148
Pro Gly Gln Ser Ala Ala Asp Ile Leu Ser Gly Ala Ala Ser Arg Arg
      10              15              20

CGG TAT CTT CTG TAT GAC GTC AAC CCC CCG GAA GGC TTC AAC CTG CGC      196
Arg Tyr Leu Leu Tyr Asp Val Asn Pro Pro Glu Gly Phe Asn Leu Arg
      25              30              35

AGG GAT GTC TAT ATC CGA ATC GCC TCT CTC CTG AAG ACT CTG CTG AAG      244
Arg Asp Val Tyr Ile Arg Ile Ala Ser Leu Leu Lys Thr Leu Leu Lys
      40              45              50

ACG GAG GAG TGG GTG CTT RTC CTG CCT CCA TGG GGC CGC CTC TNN RAC      292
Thr Glu Glu Trp Val Leu Xaa Leu Pro Pro Trp Gly Arg Leu Xaa Xaa
      55              60              65              70

TGG CAG AGT SST GAC ATC CAC CAG GTC CGG ATT CCC TGG TCT GAG TTT      340
Trp Gln Ser Xaa Asp Ile His Gln Val Arg Ile Pro Trp Ser Glu Phe
      75              80              85

TTT GAT CTT CCA AGT CTC AAT AAA AAC ATC CCC GTC ATC GAG TAT GAG      388
Phe Asp Leu Pro Ser Leu Asn Lys Asn Ile Pro Val Ile Glu Tyr Glu
      90              95              100

CAG TTC ATC GCA GAA TCT GGT GGG CCC TTT ATT      421
Gln Phe Ile Ala Glu Ser Gly Gly Pro Phe Ile
      105              110

```

(2) INFORMATION FOR SEQ ID NO: 41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 182 base pairs

- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 6..167
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1
seq VLCLRGLVSLAFQ/GP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

```

AACAA ATG TTT TTA TTT TTG TCA CCA GCC ACC CCT GTC CTG CCG CCT TCT    50
  Met Phe Leu Phe Leu Ser Pro Ala Thr Pro Val Leu Pro Pro Ser
                -50                -45                -40

CTC GAC TCC AGA GAC CTG TTG CCT CAT CTC TTT TGG GGA AGA GCC GGC    98
  Leu Asp Ser Arg Asp Leu Leu Pro His Leu Phe Trp Gly Arg Ala Gly
                -35                -30                -25

AGC TCC TCC TCA TCC CCT GCC TTA AGT CCA GTT CTT TGC CTC AGG GGT    146
  Ser Ser Ser Ser Pro Ala Leu Ser Pro Val Leu Cys Leu Arg Gly
                -20                -15                -10

CTC GTT TCC TTG GCC TTC CAG GGT CCC CAC CCC GAG    182
  Leu Val Ser Leu Ala Phe Gln Gly Pro His Pro Glu
                -5                1                5

```

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 272 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 63..125
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.8
seq LLWALLFMQSLWP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

```

AGCAGTGCAG CATTAAATGGG CCGCTGACAT GAATATGGAG TAGTTTCTC TAGCAAAGAG      60
TA ATG TGG GCC ATG GAG TCA GGC CAC CTC CTC TGG GCT CTG CTG TTC      107
  Met Trp Ala Met Glu Ser Gly His Leu Leu Trp Ala Leu Leu Phe
    -20                -15                -10

ATG CAG TCC TTG TGG CCT CAA CTG ACT GAT GGA GCC ACT CGA GTC TAC      155
Met Gln Ser Leu Trp Pro Gln Leu Thr Asp Gly Ala Thr Arg Val Tyr
   -5                1                5                10

TAC CTG GGC ATC CGG GAT GTG CAG TGG AAC TAT GCT CCC AAG GGA AGA      203
Tyr Leu Gly Ile Arg Asp Val Gln Trp Asn Tyr Ala Pro Lys Gly Arg
    15                20                25

AAT GTC ATC ACG AAC CAG CCT CTG GAC AGT GAC ATA GTG GCT TCC AGC      251
Asn Val Ile Thr Asn Gln Pro Leu Asp Ser Asp Ile Val Ala Ser Ser
    30                35                40

TTC TTA AAG TCT GAC GAT GGG      272
Phe Leu Lys Ser Asp Asp Gly
    45

```

(2) INFORMATION FOR SEQ ID NO: 43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 195 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 130..186
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.7
seq ILGLCCVLATMA/NP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

```

AAACTAAAT TTCATGCAAA GTTGATGCCA TTTCAACATG AATATGATTC TATATAGTAA      60
GTTGAATGTG TAGAAATTCA GATATCTGGA AGTTATGAAA CAAGGAATCA ATTAGGTATA      120
CAAAAGCCG ATG ACT CAC TAT AGA AAT ATT CTT GGT CTC CTG TGC TGT GTA      171
  Met Thr His Tyr Arg Asn Ile Leu Gly Leu Leu Cys Cys Val
    -15                -10

TTA GCA ACC ATG GCC AAC CCC GGG      195
Leu Ala Thr Met Ala Asn Pro Gly

```

-5

1

(2) INFORMATION FOR SEQ ID NO: 44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 333 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 253..297
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.4
seq LLLLLASLIERSS/KT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

```

AANTCATTCT TTTGCCTGGA GTTTTGTGAG GTACCCGCTT GCTTTATGGG AAAAGGCTGC   60
TCCGGAACTG CCCTACTTTA GACTTTTTCA TGGTTATCAA TCKGKACAMA GAATCACCAA   120
ACTGATAAAG CAGGAACNAG AGGGCAAATC ACGCTGCCAA GACAACTGTG TAATTCGCTC   180
GAAAAAGAAA CGAAGACAAT GTATATAAAA ATATGCAAGA ATCACAGGAA ACCCACATAT   240
CCAACCACCT AG ATG AAG TTG TTG CTG CTG TTA GCA TCA CTC ATA GAA AGA   291
          Met Lys Leu Leu Leu Leu Leu Ala Ser Leu Ile Glu Arg
          -15                      -10                      -5

AGT TCC AAA ACA AGC TGC TWN NGA CAG CAC TAT TCC AGC CAG   333
Ser Ser Lys Thr Ser Cys Xaa Xaa Gln His Tyr Ser Ser Gln
      1              5              10

```

(2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 219 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 64..177
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 8.2
 seq SFXLFLALCASFS/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

```

ATATTTAATT TATTATCTTC CTTAATTTGT AGATGAGAAA CTGAGGCATA GTAATGGACA   60

GGA ATG GCT AGG AAT CAA GCC TTG GTC TGC CTT CCC TCT TTT CAG AAT   108
  Met Ala Arg Asn Gln Ala Leu Val Cys Leu Pro Ser Phe Gln Asn
                -35                      -30                      -25

GCC TTT ATT CCT GTT GAA GAT CTT CCC ACA TCA TTT TKG CTC TTT CTC   156
Ala Phe Ile Pro Val Glu Asp Leu Pro Thr Ser Phe Xaa Leu Phe Leu
                -20                      -15                      -10

GCC CTC TGT GCT TCC TTC TCC TTT TTT TTA TKT CTT TCT CTT TCC CTC   204
Ala Leu Cys Ala Ser Phe Ser Phe Phe Leu Xaa Leu Ser Leu Ser Leu
                -5                      1                      5

CCT TCC TTC TTT TTT   219
Pro Ser Phe Phe Phe
  10
  
```

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 286 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 164..268
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 8.2
 seq SLLLLFYSFYVLA/VK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

```

ACATTATTTG TCAAAACCTC ATTAAGGTTT TGACAACAAA TGATAATTGG AAACAGTTTT   60

CTGGCTCCTT TAGTAGTAGC ATGTCATTTT ATCTGCTTCT TGGCCTTAGT ATGTTTGAAG   120

TGAAACTGTT WRABVWAGAT GTTTGTATTT GGTTCAGCT TAA ATG CCA AAT GAA   175
                Met Pro Asn Glu
                -35
  
```

TCT TGG CAA ATT CCA TGT GGC AAG CAA GAA GCT GAA ACT CTA TTC AAC	223
Ser Trp Gln Ile Pro Cys Gly Lys Gln Glu Ala Glu Thr Leu Phe Asn	
-30 -25 -20	
TTC CAA AGC TTG TTG TTG TTA TTT TAT TCC TTC TAT GTT CTA GCT GTT	271
Phe Gln Ser Leu Leu Leu Leu Phe Tyr Ser Phe Tyr Val Leu Ala Val	
-15 -10 -5 1	
AAA AGA GGG CCA GGG	286
Lys Arg Gly Pro Gly	
5	

(2) INFORMATION FOR SEQ ID NO: 47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 209 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 93..200
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8
seq LVLLICLVSSYLP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

AATTACTTCA ACTGTTGACT ATATTTTGTT TATRTTCTC ATACTGTTGT TTCTTGAATC	60
CATTCCAGTC AGGTGTGTGC CCCCATCACT TC ATG CAA ACT ACT TTT ATC GAT	113
Met Gln Thr Thr Phe Ile Asp	
-35 -30	
GTC ACT GTT GAC CAA CAT GTT GCT AAA TCT AAT GAT CAC TTG TCA GTC	161
Val Thr Val Asp Gln His Val Ala Lys Ser Asn Asp His Leu Ser Val	
-25 -20 -15	
CTA GTC TTA CTT ATT TGC TTA GTC TCA TCT TAT TTG CCA CAG CTC CCG	209
Leu Val Leu Ile Cys Leu Val Ser Ser Tyr Leu Pro Gln Leu Pro	
-10 -5 1	

(2) INFORMATION FOR SEQ ID NO: 48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 363 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 124..342
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.8
seq YLPLLAGLGLTLA/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

```

AAGGGTCCGC CAGCTCGGGG CCAGCGCATG GGGCTGCTGA GACCGCTCCG GACGTGCGAA   60
SGTTTCGCGGT GCGGTAGGAA GCTGAAGTCT TCAGCAGAAC CCTGACCTAG GTAGGGAGTC   120
TCA ATG GCC CTG GGT GAA GAA AAG GCA GAA GCG GAA GCA TCT GRR GAC   168
  Met Ala Leu Gly Glu Glu Lys Ala Glu Ala Glu Ala Ser Xaa Asp
                    -70                      -65                      -60
ACA AAG GCC CAG TCC TAT GGG AGA GGG AGC TGC AGG GAG CGG GAG CTG   216
  Thr Lys Ala Gln Ser Tyr Gly Arg Gly Ser Cys Arg Glu Arg Glu Leu
                    -55                      -50                      -45
GAC ATC CCA GGG CCC ATG AGT GGG GAG CAG CCC CCA CGC CTG GAA GCT   264
  Asp Ile Pro Gly Pro Met Ser Gly Glu Gln Pro Pro Arg Leu Glu Ala
                    -40                      -35                      -30
GAG GGA GGG CTC ATC TCC CCT GTA TGG GGG GCA GAA NGA TAC CTG CCC   312
  Glu Gly Gly Leu Ile Ser Pro Val Trp Gly Ala Glu Xaa Tyr Leu Pro
                    -25                      -20                      -15
CTA CTT GCT GGA TTG GGA CTG ACC CTG GCG GCC CCT CTA GAG CCC ACA   360
  Leu Leu Ala Gly Leu Gly Leu Thr Leu Ala Ala Pro Leu Glu Pro Thr
                    -5                      1                      5
ACG
Thr
363

```

(2) INFORMATION FOR SEQ ID NO: 49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 271 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 149..211
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.7
seq LLCISPFVPEFTSG/NK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

```

AAAACTCTAT GAGTGTCTTT TTGAGACCAT AAAGCAGACT TTAGTAACTT TCTATTTCTG   60
TAAGTACTAA ATGTCTGGCA TTTTAACTT TTGTAGAATA CATAATGTTG RAACTGGAA   120
TAATACTATT KATTTTCACC TGTGAAAA ATG ACT TCA TTG TAC TTG AAA CAC   172
                               Met Thr Ser Leu Tyr Leu Lys His
                               -20                               -15

CTC CTT TGC ATT TCT CCA TTT GTG CCA TTC ACT AGT GGA AAT AAA TTG   220
Leu Leu Cys Ile Ser Pro Phe Val Pro Phe Thr Ser Gly Asn Lys Leu
-10                               -5                               1

TAT TAT ACC ATG ATC TAC TGG CTT TTT AAA ACT GTA TTA AAT ATG CAC   268
Tyr Tyr Thr Met Ile Tyr Trp Leu Phe Lys Thr Val Leu Asn Met His
  5                               10                               15

GGG
Gly
20

```

271

(2) INFORMATION FOR SEQ ID NO: 50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 378 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 205..345
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.6
seq CLATLTLFHTSFS/FQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

```

AGGGCTCTCC ATGAAGCGGG TGTGTGCAAC TTA CTGGAA CCAGATATTG AGCACTTATT   60
TTATATCAGG CAGAATACAG AGCCACTTAY ATGRKWTCAA TKAATTCTTA CAACTCCCTT   120
GGGAARGAGG TGTTTGTCTT CTTTCATTT AACAGATGAG AAGACTGAGA CTTGATGAGT   180

```

```

TTAGTCAATT GTTCTGAGTA GTAA ATG ACA GAC TCT CCC AAT GCT CAT GGC      231
                Met Thr Asp Ser Pro Asn Ala His Gly
                -45                      -40

TTA GCT CTC ACC ACC AAG TGG ATG ATG CCT GCT GTC TCT TTG AAC TTG      279
Leu Ala Leu Thr Thr Lys Trp Met Met Pro Ala Val Ser Leu Asn Leu
                -35                      -30                      -25

ACC TAT TAC TTG CCA TCT TGG TAC CTT TGT TTG GCC ACT CTT ACT TTA      327
Thr Tyr Tyr Leu Pro Ser Trp Tyr Leu Cys Leu Ala Thr Leu Thr Leu
                -20                      -15                      -10

TTC CAC ACC TCT TTC TCC TTC CAA GCT TCT GAG TCT GTC AAA GCC ATC      375
Phe His Thr Ser Phe Ser Phe Gln Ala Ser Glu Ser Val Lys Ala Ile
                -5                      1                      5                      10

ACG                                                                378
Thr

```

(2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 376 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 143..286
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5
seq FVILLLFIFTVVS/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

```

AATCGCTTCA GCAGCATCCT CTCAGACAAG AGCCACTATT TCTGATTCAG ATCACCTGTC      60

ATCGAAGTTT AAAGAAGGGG AAACAGGAGA CAGAAATACA CTGAACCAA AAGATTCAA      120

AGAGCAAGTG GAATCTCTAA GA ATG GCT TCC AGC CAC TGG AAT GAA ACC ACT      172
                Met Ala Ser Ser His Trp Asn Glu Thr Thr
                -45                      -40

ACC TCT GTT TAT CAG TAC CTT GGT TTT CAA GTT CAA AAA ATT TAC CCT      220
Thr Ser Val Tyr Gln Tyr Leu Gly Phe Gln Val Gln Lys Ile Tyr Pro
                -35                      -30                      -25

TTC CAT GAC AAC TGG AAC ACT GCC TGC TTT GTC ATC CTG CTT TTA TTT      268
Phe His Asp Asn Trp Asn Thr Ala Cys Phe Val Ile Leu Leu Leu Phe
                -20                      -15                      -10

```

[illegible]

(2) INFORMATION FOR SEO ID NO: 52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 429 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 298..372
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 7.4
seq IFLNMMWVACLLS/GE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

ACTGAGATCC	ACCTGGGCTC	CCAAGACAGA	TAGTGGGGTC	TTGTCCAGTT	CATATATTAT	60
CTACAGTTGC	CAATGTGATA	TAAAATATGA	TTTAGGTCAT	ATCACCTACT	TGAAAGTCTG	120
CACTGGTGCT	GTCTTGGTAA	CAGAAAGAGA	AAGACTGAAT	TCCCTTATTT	TGCCCCATCAG	180
GCTCTCCTTG	ATATAGTCTC	TGCCCCACCA	CCTCTCCAAC	CTCACATCAT	CTCTCTTCTG	240
CCTCATACGC	TATGCTCCGG	CACGTATAGG	TTCCTATACA	ATTTTGTTTC	ATACTTG	297
ATG TCT TTA CTT TTT GTC TTT TGC CTG GAA TGC AGT ATT TTT CTA TTG	345					
Met Ser Leu Leu Phe Val Phe Cys Leu Glu Cys Ser Ile Phe Leu Leu						
-25 -20 -15 -10						
AAT ATG TGG GTT GCT TGC CTT CTG AGT GGT GAG ATT CCC CAT TCC TCA	393					
Asn Met Trp Val Ala Cys Leu Leu Ser Gly Glu Ile Pro His Ser Ser						
-5 1 5						
TGG AAN MTA AAG TTA ATT GGC ACC TTG CCC ACT TCT	429					
Trp Xaa Xaa Lys Leu Ile Gly Thr Leu Pro Thr Ser						
10 15						

(2) INFORMATION FOR SEQ ID NO: 53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 244 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 149..202
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.4
seq VLLLLPLVAFITL/KF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

```

AAAATTTCAA AACGTAGGAT AAAAGGAATC AAATGTATTA ATGGAWAAKC AACTGAACTT   60
AATTTTCGATT CTTTCTATC ATTTTTCCT AGGCTAKAGA TAGACTAAAT TCATATCTGA   120
AAATTCTCAA TTTTGTAGAA AAGACAAA ATG TTT GTC GTT ACA GTT TTG TTG   172
                               Met Phe Val Val Thr Val Leu Leu
                               -15

TTG CTG CCC TTA GTT GCT TTC ATT ACC CTC AAA TTC TGT AAC TTG ATT   220
Leu Leu Pro Leu Val Ala Phe Ile Thr Leu Lys Phe Cys Asn Leu Ile
-10                -5                1                5

AAT TTT CCA ACT CWK AGA CAC GGG   244
Asn Phe Pro Thr Xaa Arg His Gly
                10

```

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 212 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide

(B) LOCATION: 12..77
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 7.2
 seq ILYALQFLFLVFA/PS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

```

AATTCAATCA A ATG AAC CGA TCT TGT AGA AAC ACT GGA ATC ATT TAT GCG      50
      Met Asn Arg Ser Cys Arg Asn Thr Gly Ile Ile Tyr Ala
                -20                      -15                      -10

TTG CAG TTT CTC TTT CTT GTT TTC GCT CCT TCT TCC CTG GGA TAT TTT      98
Leu Gln Phe Leu Phe Leu Val Phe Ala Pro Ser Ser Leu Gly Tyr Phe
                -5                      1                      5

GAG TGG ATT GTG GCT ATT AAT CAA GAT CTC GTG CTA TTC GTG TTT TGC      146
Glu Trp Ile Val Ala Ile Asn Gln Asp Leu Val Leu Phe Val Phe Cys
                10                      15                      20

TTG TCG TTT TCG CTC AGG ATT AGC ATC ATT CAA GGC AAA CGC AAA GCT      194
Leu Ser Phe Ser Leu Arg Ile Ser Ile Ile Gln Gly Lys Arg Lys Ala
                25                      30                      35

GCT TTT CCC ACC CCC CCC
Ala Phe Pro Thr Pro Pro
212
40                      45
  
```

(2) INFORMATION FOR SEQ ID NO: 55:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 221 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 87..191
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 7.2
 seq LSLLLAWVTILTHL/LS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

```

ACCTTCGACC CCCACCCTGG GAATACCAAT CCAGTGATTC CACCATCTAC TCACTGTCCC      60

TCATCGTTGG TGTCTTCTCT AACTTC ATG ACC CAA ACC ACA TGG GGA GCC CCC      113
      Met Thr Gln Thr Thr Trp Gly Ala Pro
                -35                      -30

ACC AGG GCC AGC AAT CAC CCT CTC CCT GCA TGG CTC ACC CTC AGC CTC      161
  
```

Thr Arg Ala Ser Asn His Pro Leu Pro Ala Trp Leu Thr Leu Ser Leu
 -25 -20 -15

CTC CTG GCC TGG GTG ACC CTT ACA CAC CTT CTC TCT GTG CTC ACA CAT 209
 Leu Leu Ala Trp Val Thr Leu Thr His Leu Leu Ser Val Leu Thr His
 -10 -5 1 5

CCA ACC CTC CTG 221
 Pro Thr Leu Leu
 10

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 328 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 74..121
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1
seq XXAVLCVCAAAWC/SQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

AGAAASNCCG AGTTGGCAGA GCAGGGCTGC ATTTCCAGCA GGAGCTGCGA GCACAGTGCT 60

GGCTCACAAAC AAG ATG CTC AAG GDV NNA GCC GTA CTG TGT GTG TGT GCA 109
 Met Leu Lys Xaa Xaa Ala Val Leu Cys Val Cys Ala
 -15 -10 -5

GCC GCT TGG TGC AGT CAG TCT CTC GCA GCT GCC GCG GCG GTG GCT GCA 157
 Ala Ala Trp Cys Ser Gln Ser Leu Ala Ala Ala Ala Val Ala Ala
 1 5 10

DCC GGG GGG CGG TCG GAC GGC GGT AAT TTT CTG GAT GAT AAA CAA TGG 205
 Xaa Gly Gly Arg Ser Asp Gly Gly Asn Phe Leu Asp Asp Lys Gln Trp
 15 20 25

CTC ACC ACA ATC TCT CAG TAT GAC AAG GAA GTC GGA CAG TGG AAC AAA 253
 Leu Thr Thr Ile Ser Gln Tyr Asp Lys Glu Val Gly Gln Trp Asn Lys
 30 35 40

TTC CGA GAC GAT GAT TAT TTC CGC ACT TGG AGT CCA GGA AAA CCC TTC 301
 Phe Arg Asp Asp Asp Tyr Phe Arg Thr Trp Ser Pro Gly Lys Pro Phe
 45 50 55 60

GAT CAG GCT TTA GAT CSA GCT AAC GGG 328
 Asp Gln Ala Leu Asp Xaa Ala Asn Gly

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 233 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 168..227
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9
seq LHLGSSISPASA/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

```

AATGAAAAGC AGTTTGTTAT TATGCAGGAA AATCAGTTTC ATCATTTTAG TTACACTAAA   60
CACTTTTGGC AGCTTAATAT GACCTTTTTA AATTTTCTKT TATTTTTTTT ATTTTATTT   120
CTTTAAGATG GAGTCTTGCT CTGTTGCCCG GGCTGGAGTA CAATGGC ATG ATC TCA   176
                                     Met Ile Ser
                                     -20
GCT CAC TGC AAC CTC CAC CTC CTG GGT TCA AGC ATT TCT CCT GCC TCA   224
Ala His Cys Asn Leu His Leu Leu Gly Ser Ser Ile Ser Pro Ala Ser
   -15                -10                -5
GCC TCC CTG
Ala Ser Leu
   1

```

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 170 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 6..89
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9
seq FLPFLLSLPLDQT/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

```

AAGAT ATG TGC SCB AAA GCC TGC AGA ACC CTC GCT TGG TTG CCT GAS CCG      50
Met. Xaa Xaa Lys Ala Cys Arg Thr Leu Ala Trp Leu Pro Xaa Pro
      -25                -20                -15

TTC TTA CCC TTT CTC CTC AGT CTT CCC TTG GAC CAG ACG CTT CCT CGC      98
Phe Leu Pro Phe Leu Leu Ser Leu Pro Leu Asp Gln Thr Leu Pro Arg
      -10                -5                1

CAG GGT CCT GGC CAA TCC CTG TCC TTC CCA GAA AAC TAC CAG ACT CTT      146
Gln Gly Pro Gly Gln Ser Leu Ser Phe Pro Glu Asn Tyr Gln Thr Leu
      5                10                15

CCC AAG AGC ACC CGA CAC CCT GGG      170
Pro Lys Ser Thr Arg His Pro Gly
      20                25

```

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 19..75
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8
seq GLLLVFLPHPQRG/GQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

```

ACATCGCCCG AGCAGGGG ATG GCG GTG AAG CGG CTA GGG CTG CTG TTG GTG      51
Met Ala Val Lys Arg Leu Gly Leu Leu Val
      -15                -10

TTC CTG CCT CAT CCG CAG CGG GGA GGA CAG GAG AGG TCT GCC CAC ACC      99
Phe Leu Pro His Pro Gln Arg Gly Gly Gln Glu Arg Ser Ala His Thr
      -5                1                5

CCG AGG CAG CAC CCA GCT CGC CCC ACT TCC CTC TCG CAG GGG GAG AGA      147

```

Pro Arg Gln His Pro Ala Arg Pro Thr Ser Leu Ser Gln Gly Glu Arg
 10 15 20
 CCA GGA CGC GGT GGG GGG TGG GGG AAT GGC CGT GAC GCC CAC CAG 192
 Pro Gly Arg Gly Gly Gly Trp Gly Asn Gly Arg Asp Ala His Gln
 25 30 35

(2) INFORMATION FOR SEQ ID NO: 60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 433 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 77..325
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8
seq LLMILTFPFKILS/DA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

AAGGCTGCAA TAACTACTAC TTACTGGATA CATTCAAACC CTCCAGAATC AACAGTTATC 60
 AGGTAACCAA CAAGAA ATG CAA GCC GTC GAC AAC CTC ACC TCT GCG CCT GGG 112
 Met Gln Ala Val Asp Asn Leu Thr Ser Ala Pro Gly
 -80 -75
 AAC ACC AGT CTG TGC ACC AGA GAC TAC AAA ATC ACC CAG GTC CTC TTC 160
 Asn Thr Ser Leu Cys Thr Arg Asp Tyr Lys Ile Thr Gln Val Leu Phe
 -70 -65 -60
 CCA CTG CTC TAC ACT GTC CTG TTT TTT GTT GGA CTT ATC ACA AAT GGC 208
 Pro Leu Leu Tyr Thr Val Leu Phe Phe Val Gly Leu Ile Thr Asn Gly
 -55 -50 -45 -40
 CTG GCG ATG AGG ATT TTC TTT CAA ATC CGG AGT AAA TCA AAC TTT ATT 256
 Leu Ala Met Arg Ile Phe Phe Gln Ile Arg Ser Lys Ser Asn Phe Ile
 -35 -30 -25
 ATT TTT CTT AAG AAC ACA GTC ATT TCT GAT CTT CTC ATG ATT CTG ACT 304
 Ile Phe Leu Lys Asn Thr Val Ile Ser Asp Leu Leu Met Ile Leu Thr
 -20 -15 -10
 TTT CCA TTC AAA ATT CTT AGT GAT GCC AAA CTG GGA ACA GGA CCA CTG 352
 Phe Pro Phe Lys Ile Leu Ser Asp Ala Lys Leu Gly Thr Gly Pro Leu
 -5 1 5
 AGA ACT TTT GTG TGT CAA GTT ACC TCC GTC ATA TTT TAT KKC RSA ATG 400
 Arg Thr Phe Val Cys Gln Val Thr Ser Val Ile Phe Tyr Xaa Xaa Met

10

15

20

25

TAT ATC AGT ATT TCA TTC CTG GGA CTG ATA ACT
 Tyr Ile Ser Ile Ser Phe Leu Gly Leu Ile Thr
 30 35

433

(2) INFORMATION FOR SEQ ID NO: 61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 176 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 12..119
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8
seq LWFFLPSLXCPEC/CP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

ATCTGGTCTG C ATG TGG ACC CTC CCC AGC CTC AGT GCA TCG TTT CAG CCT 50
 Met Trp Thr Leu Pro Ser Leu Ser Ala Ser Phe Gln Pro
 -35 -30 -25

TTT CTG GGC AGC CTT CGC CCT TCT CAC ATC CTG TGG TTC TTC CTG CCC 98
 Phe Leu Gly Ser Leu Arg Pro Ser His Ile Leu Trp Phe Phe Leu Pro
 -20 -15 -10

TCC CTC CMC TGC CCA GAA TGC TGC CCT CCT GAT CCA GGA TCT CCA GCC 146
 Ser Leu Xaa Cys Pro Glu Cys Cys Pro Pro Asp Pro Gly Ser Pro Ala
 -5 1 5

TCC AGG GAC CCT AAC GTG GCC TGC GAA CGG 176
 Ser Arg Asp Pro Asn Val Ala Cys Glu Arg
 10 15

(2) INFORMATION FOR SEQ ID NO: 62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 287 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 36..101
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.7
seq LLLFQPSSHSATG/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

```

ATGAAACAAA ATATGTTCTA CCAGATAAAG TATTG ATG TCC CTA ACT GAT GTC      53
                               Met Ser Leu Thr Asp Val
                               -20

CCC ATG TCC CTT CTG CTT TTT CAA CCC AGT TCC CAC AGT GCT ACT GGA      101
Pro Met Ser Leu Leu Leu Phe Gln Pro Ser Ser His Ser Ala Thr Gly
-15                               -10                               -5

TCA TCT ATC AAA ATT ATA ATA CTT AAT TAC ATC ATA CTC CAG TTC AAA      149
Ser Ser Ile Lys Ile Ile Ile Leu Asn Tyr Ile Ile Leu Gln Phe Lys
 1                               5                               10                               15

ACC CTT CAA ACA CTT CCT AAT GCT TTG AGG ATA CAC ATC AAA GTC TTT      197
Thr Leu Gln Thr Leu Pro Asn Ala Leu Arg Ile His Ile Lys Val Phe
      20                               25                               30

CAC ATT TAC TGT TCA TTT GTT TCC AGG TTT CAC TAT TAT AAA AAT ACT      245
His Ile Tyr Cys Ser Phe Val Ser Arg Phe His Tyr Tyr Lys Asn Thr
      35                               40                               45

GCC ACA GTA TTT TTC AGG TCA GTA TTA AAA AGG AGA ATG GGG      287
Ala Thr Val Phe Phe Arg Ser Val Leu Lys Arg Arg Met Gly
      50                               55                               60

```

(2) INFORMATION FOR SEQ ID NO: 63:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 453 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 238..288
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.6
seq LAFLLVSLYWSHM/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

```

AATAGGGCTG AACCGAGGAC TGAAAAAGGG AGGAGGCAGA CCACTCGGAG AGGAGCTGGG      60
AAGCAGTGCA GAGAGGAGAG CGGASSNAGC TGCCGCTGAG CAAAGTGCGA CTGTATCTGG      120
TTTGGCCTGC TCTTCCTCAC CTTCTCCTT TCCCTGAGCT GGCTGTACAT CGGGCTCGTC      180
CTTCTCAATG ACCTGCACAA CTTCAATGAA TTCCTCTTCC GCCGCTGGGG AACTG      237
ATG GAC TGG TCC CTG GCA TTC CTG CTG GTA TCT CTC TAC TGG TCA CAT      285
Met Asp Trp Ser Leu Ala Phe Leu Leu Val Ser Leu Tyr Trp Ser His
   -15                      -10                      -5

ATG CAT CCT TGC TAT TGG TCC TGG CCC TGC TCC TGC GGC TTT GTA GAC      333
Met His Pro Cys Tyr Trp Ser Trp Pro Cys Ser Cys Gly Phe Val Asp
   1                      5                      10                      15

AGC CCC TGC ATC TGC ACA GCC TCC ACA AGG TGC TGC TGC TCC TCA TTA      381
Ser Pro Cys Ile Cys Thr Ala Ser Thr Arg Cys Cys Cys Ser Ser Leu
           20                      25                      30

TGC TGC TTG TGG CGG CTG GCC TTG TGG GAC TGG ACA TCC AAT GGC AGC      429
Cys Cys Leu Trp Arg Leu Ala Leu Trp Asp Trp Thr Ser Asn Gly Ser
           35                      40                      45

AGG AGT GGC ATA GCT TGC GTG TGT      453
Arg Ser Gly Ile Ala Cys Val Cys
   50                      55

```

(2) INFORMATION FOR SEQ ID NO: 64:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 103..147
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5
seq LLILFFMVGRIIP/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

```

ATATTATATT TTTTAGTCAG ACTCACAGAA TTGAGTTGAT TTTATTCCTC ATTGGGTGGC      60
ACACTATTAT CGTTCTTCCC AAMCTTGTTT AGTATTTGTW TT ATG TAT TTA CTA      114

```

Met Tyr Leu Leu
-15

ATA CTT TTC TTT ATG GTA GGC AGA ATT ATC CCT TCC CCC CAC CGG 159
Ile Leu Phe Phe Met Val Gly Arg Ile Ile Pro Ser Pro His Arg
-10 -5 1

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 245 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 78..221
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5
seq LLVVSCLLFHQ/IH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

AATGGATTAT GTGCTATGTG CACTCCACAG CCTAAGAACT ACAATCTTGG TTGTTGTTAA 60
TTTTTGATAC CTAAAG ATG AAT AAG CCT CCC TGG GAA GAA TCC TGG GGG 110
Met Asn Lys Pro Pro Trp Glu Glu Ser Trp Gly
-45 -40
CAA AAT CAA CTC TCA GGT GAG CCC GCT ACT TGG AGT CTT TGC ATT TCT 158
Gln Asn Gln Leu Ser Gly Glu Pro Ala Thr Trp Ser Leu Cys Ile Ser
-35 -30 -25
CCT TTG CCT GGA AGA GAG CCG AGT TTA CTT GTT GTG TCT TGT TGC CTC 206
Pro Leu Pro Gly Arg Glu Pro Ser Leu Leu Val Val Ser Cys Cys Leu
-20 -15 -10
TTG TTT CAT CAG GCT ATC CAT AAT AAA TTA CTG TGG CGG 245
Leu Phe His Gln Ala Ile His Asn Lys Leu Leu Trp Arg
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 299 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 177..248
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5
seq FLILLSIDSLVSG/FL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

```

AATTTTTTGC GACTTATTGA GGTGTCTTT TCTGTCATTC AGGAAAGTTT TATGGTTTTC   60
TCTGGAAAGG TCTTGCTTGT TTCTTGTTAG GATTCTTGAT ACTTTTGAGT CTGTTTCCTC  120
CTTTGGTGTG TCATAATGGT TATTGATGGT GTATTGGGAA GTTATAGATA ATTTGG ATG   179
                                     Met
TTG ATC TTA GAG CTA ACA ATG ATG CTG AGC TTT CTA ATT CTA TTG TCA   227
Leu Ile Leu Glu Leu Thr Met Met Leu Ser Phe Leu Ile Leu Leu Ser
      -20                      -15                      -10

ATT GAT TCT CTT GTA TCG GGT TTT TTA AGT AAG CGA AAA GGT CTG CGC   275
Ile Asp Ser Leu Val Ser Gly Phe Leu Ser Lys Arg Lys Gly Leu Arg
      -5                      1                      5

GTC TGT GAT GGA AGC CGG TCC GGG   299
Val Cys Asp Gly Ser Arg Ser Gly
10                      15

```

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 183..338
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4
seq CLLGAAWASRLRT/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

```

AATAGACGAG TAGACCTTAG GCGAGGGAAG GCATGTACTT TCCCTAAGAA GCGGGAGGAG    60
AGAAATAAAT AGAGGGCAGT GGGATGAGAA GAACCTCGCG GGATGGGAAG CCGCGGAGGG    120
AAGGGCCGTC TTTGGTTACC TGGAGAGCGG GAGCAGCTGC GGATCCCTTA TGAAGTGCCC    180
GG ATG AAG CTC CAG CGC TCA CGC GCT TTC CGC ATT GAG TGC AGC GCC    227
  Met Lys Leu Gln Arg Ser Arg Ala Phe Arg Ile Glu Cys Ser Ala
      -50                -45                -40
ATC TTG AGA AGG GCG GAG CGT CTT GTK TGG AAT GAC GTC TGT TCA GAG    275
Ile Leu Arg Arg Ala Glu Arg Leu Val Trp Asn Asp Val Cys Ser Glu
      -35                -30                -25
AGC CAA TCC CAG TCT CGC GAC TCC TGC TTG CTG GGC GCG GCT TGG GCC    323
Ser Gln Ser Gln Ser Arg Asp Ser Cys Leu Gly Ala Ala Trp Ala
      -20                -15                -10
TCC AGG CTG CGC ACG CAG CCG CAT CCG    350
Ser Arg Leu Arg Thr Gln Pro His Pro
      -5                1

```

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 295 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 236..283
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq FTLCVFTLPFLCA/CL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

```

ACAAAAGGGA ATGTATTAGT GTCCCAGGAC TACTGTGACA AGGCACAAA AACTGGGTGC    60
TTTACAACAA GAAACTGTA CAGTGCTGGA GGCTAGAAGT CAAAATCAAG TGATTGGTTG    120
GACCATGCTC TCTCTGACAG TGACAGGGGA GAACCCTCCC TCGCCTCTCC TGGCTTCTGG    180
TATGCACCAG CAATTCCTGG CGTTCCTTGG CTCCTAGAAG CATCACTCCT ATCAC ATG    238
                                     Met
GTC ATC TTC ACC CTG TGT GTC TTC ACA CTA CCC TTT CTC TGT GCA TGT    286
Val Ile Phe Thr Leu Cys Val Phe Thr Leu Pro Phe Leu Cys Ala Cys
-15                -10                -5                1

```


CTG CCC AGG
Leu Pro Arg

295

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 285 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 184..240
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq VLVVGTWSSQGQA/NS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

```

AAAACCTCTAG AAATCTGTGT TTCCGGAAAG CAGGGGTGGG AGCCGCCAG TGCCTTCCCC    60
AAAACCCAAC ACTGAGATTT CAATGRTATG KTTTGKGAGT CTTTTTAAAA ATTCTCTCTG   120
GTTGCTGGGC TCACTGACCA CGGTCAGGTT TGGAAGAGCA CCAGGCTGTG CGGGCCGAGG   180
CGG ATG TGG GGA GCA CTT CCT GTC CTC GTG GTG GGA ACC TGG TCC AGC       228
  Met Trp Gly Ala Leu Pro Val Leu Val Val Gly Thr Trp Ser Ser
           -15                      -10                      -5
CAA GGG CAG GCG AAC AGC TGT GCG GGG CGG GGG ATG GGG CCA GAC GTG       276
  Gln Gly Gln Ala Asn Ser Cys Ala Gly Arg Gly Met Gly Pro Asp Val
           1                      5                      10

TGT GGA GCG                               285
Cys Gly Ala
           15

```

(2) INFORMATION FOR SEQ ID NO: 70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 247 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 143..190
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq LVCGLQISLSLA/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

```
AGTGACTGTA CACTGGGTTT CTTTGGAAC TCGTCTTTT CTCAATAATG GCAGGATCCC 60
CGATTTTAGA AAGTGGGCAG TGCTTTGTGT TACAGGGTTT GGGCTAAAGA CTGTTTGATA 120
ACCAATATTT TACAAGAATT AA ATG ACC AGA TTG GTC TGT GGC TTT CTC CAA 172
                Met Thr Arg Leu Val Cys Gly Phe Leu Gln
                -15                               -10
ATT TCC TTA TCC CTA GCT TCA TTG TTT CTG ACA ATT CCT CTA ATG TGG 220
Ile Ser Leu Ser Leu Ala Ser Leu Phe Leu Thr Ile Pro Leu Met Trp
-5              1              5              10
TAC ATG CAA TCA AAA TGG TGG CGT GGG 247
Tyr Met Gln Ser Lys Trp Trp Arg Gly
                15
```

(2) INFORMATION FOR SEQ ID NO: 71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 114 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 55..99
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.6
seq FLLPLLLHHLTFH/GR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

```
AATCTATTTG TCCTCTAGAA ACTCTTCATA ATGCCCTATC TTAAAGCAA GTGG ATG 57
                Met
                -15
```

AAT TTC TTG CTT CCA TTG CTT CTC CAT CAT CTG ACG TTC CAC GGA AGA 105
Asn Phe Leu Leu Pro Leu Leu Leu His His Leu Thr Phe His Gly Arg
-10 -5 1

CCG CTG AAG 114
Pro Leu Lys
5

(2) INFORMATION FOR SEQ ID NO: 72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 367 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 125..298
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.5
seq LIIFICXTASISA/YM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

CTGCTTCACT	TTCAGGTTTC	TCGAAGTGCC	TTCTTGCTCC	TGTCTGTTTC	CCCATCCTGC		60
CAGATTCTCG	TTTCTCTTGC	TGGGCTTTTTG	GCAGTAGGGG	GCTGTGTTGG	TGGGCCCTAC		120
GAAG ATG CTC AGT GCT CGA GAT CGC CGG GAC CGG CAC CCT GAG GAG GGG							169
Met Leu Ser Ala Arg Asp Arg Arg Asp Arg His Pro Glu Glu Gly							
-55			-50			-45	
GTA GTT GCA GAG CTC CAG GGC TTC GCG GTG GAC AAG GCC TTC CTC ACC							217
Val Val Ala Glu Leu Gln Gly Phe Ala Val Asp Lys Ala Phe Leu Thr							
-40			-35			-30	
TCC CAC AAG GGC ATC CTG CTG GAA ACC GAG CTG GCC CTG ACC CTC ATC							265
Ser His Lys Gly Ile Leu Leu Glu Thr Glu Leu Ala Leu Thr Leu Ile							
-25			-20			-15	
ATC TTC ATC TGC DTC ACG GCC TCC ATC TCT GCC TAC ATG GCC GCG GCG							313
Ile Phe Ile Cys Xaa Thr Ala Ser Ile Ser Ala Tyr Met Ala Ala Ala							
-10			-5			1	5
CTA CTG GAG TTC TTC ATC ACA CTT GCC TTC CTC TTC CTC TAT GCC ACC							361
Leu Leu Glu Phe Phe Ile Thr Leu Ala Phe Leu Phe Leu Tyr Ala Thr							
10			15			20	
CCA GCG							367
Pro Ala							

(2) INFORMATION FOR SEQ ID NO: 73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 338 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 225..263
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5
seq MLTMSVTLSPLRS/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

```

AAGATGTGCT GTAAACATCA AAAGAAGACG GTGGGATCAG GAGATGTTTG GACAGCTCTT   60
TATTCAGAAC ATCAAGGACT GTTGACAGTT GAATAAHNAG GGCGGAGGCT TATGGGATTG  120
CTAATGAGAT ACAAAGCCAC CTTGGAATAA AAATAAAHTT CTCTCTGTTG GCTCCTCCGG  180
CCATGGAGAG CTGTTTTVGA AAGAAGTGAG GTTTAGACTT CTCC ATG TTA ACC ATG   236
                                   Met Leu Thr Met
                                   -10

AGC GTG ACA CTT TCC CCC CTG AGG TCA CAG GAC CTG GAT CCC ATG GCT   284
Ser Val Thr Leu Ser Pro Leu Arg Ser Gln Asp Leu Asp Pro Met Ala
          -5                      1                      5

ACT GAT GCT TCA CCC ATG GCC ATC AAC ATG ACA CCC ACT GTG GAG CAG   332
Thr Asp Ala Ser Pro Met Ala Ile Asn Met Thr Pro Thr Val Glu Gln
        10                      15                      20

GGA CTG                                     338
Gly Leu
    25

```

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 279 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 100..237
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq FFLLISSVRPISQ/TF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

```

ATTTAAATTT TGTGATCAAA GATGCCAAAT GGACACAACA CCATACCCTA GTTATTTCCC      60
TACTGTGTTT TCCTACAAGC TTACCTGCAT TTAGTAMCC ATG TTT ATR CCC GTA      114
                               Met Phe Xaa Pro Val
                               -45
GCA CTG ATC TTC CCC ATC TCA GTA AGT GAC CCT ACC ATT CAC CCT ATT      162
Ala Leu Ile Phe Pro Ile Ser Val Ser Asp Pro Thr Ile His Pro Ile
-40                               -35                               -30
ACT CAA GCC CAG AAC CTA GAA AGC NTC CTA CAG TCC TTC TTT CTT CTA      210
Thr Gln Ala Gln Asn Leu Glu Ser Xaa Leu Gln Ser Phe Phe Leu Leu
-25                               -20                               -15                               -10
ATA TCA TCT GTA AGA CCC ATT AGT CAA ACC TTC AAA ATA GAT CTT TCT      258
Ile Ser Ser Val Arg Pro Ile Ser Gln Thr Phe Lys Ile Asp Leu Ser
                               -5                               1                               5
CCA TCT GTG CGG GCH ACC GGG      279
Pro Ser Val Arg Ala Thr Gly
10

```

(2) INFORMATION FOR SEQ ID NO: 75:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 91..183
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq WCAVLRSWLAASS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

```

ATGCAAAAAA GCACTGTGCC TTCTGTCATT GAATCCTAGA ACACTGAGAT GAGAAAAACA. 60
CTGTAGATGT ATAGGTTAGA TATATAGGGA ATG TTA TTA TTT TTT CCT TTT TTT 114
                               Met Leu Leu Phe Phe Pro Phe Phe
                               -30                               -25
GGA GAG ACG GTC TCG CTC CAT CAC CCA TGC TGG TGT GCA GTG CTG CGA 162
Gly Glu Thr Val Ser Leu His His Pro Cys Trp Cys Ala Val Leu Arg
          -20                               -15                               -10
TCG TGG CTC GCT GCA TCC TCC GCC CCT CGG 192
Ser Trp Leu Ala Ala Ser Ser Ala Pro Arg
          -5                               1

```

(2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 199 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 32..136
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq LVVVCYLSWRVSS/RS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

```

ATACTCAAAA TCTACATTTT AGTTAGTTGA T ATG CCT CTT AAG AAT CTG TTC 52
                               Met Pro Leu Lys Asn Leu Phe
                               -35                               -30
TCT GTT GGT CTG TGG GAT CCT TAC AAT TTA CTG AAG AAA CAT GTT TTG 100
Ser Val Gly Leu Trp Asp Pro Tyr Asn Leu Leu Lys Lys His Val Leu
          -25                               -20                               -15
GTT GTT GTC TGC TAT TTA TCC TGG AGA GTG TCT TCC AGA AGT TGG ACT 148
Val Val Val Cys Tyr Leu Ser Trp Arg Val Ser Ser Arg Ser Trp Thr
          -10                               -5                               1
TTG CTG ATT ACA CCT GTA ACA CTT CAT GCT TCT CTG TCC ACC CAG GCC 196
Leu Leu Ile Thr Pro Val Thr Leu His Ala Ser Leu Ser Thr Gln Ala
          5                               10                               15                               20
CGG 199
Arg

```

(2) INFORMATION FOR SEQ ID NO: 77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 418 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 209..265
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq LSHLLPSLRQVIQ/EP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

```

ATGTTTGATT GGTGTGCCTG AAAGTGAAGG GGAGAATGAA AACATCTGCC AGCTCCGCCG    60
ACCCGGCCCC CGCGGCCCTC CCAGCTCGGC TCCGGCTCAG TGGACAGGAA CCACTGAAGT    120
TTGCCTGACA CCATCAACCA GGCCCTAGTC ACCTGGCTTT GCCTTTGCCC TGCTGTGTGA    180
TCTTAGCTCC CTGCCCAGGC CCACAGCC ATG GCC ATG GCC CAG AAA CTC AGC      232
                               Met Ala Met Ala Gln Lys Leu Ser
                               -15

CAC CTC CTG CCG AGT CTG CGG CAG GTC ATC CAG GAG CCT CAG CTA TCT      280
His Leu Leu Pro Ser Leu Arg Gln Val Ile Gln Glu Pro Gln Leu Ser
-10                               -5                               1                               5

CTG CAG CCA GAG MYG GTC TTC ACG GTG GAT CGA GCT GAG GTG CCG CCG      328
Leu Gln Pro Glu Xaa Val Phe Thr Val Asp Arg Ala Glu Val Pro Pro
                               10                               15                               20

CTC TTC TGG AAG CCG TAC ATC TAT GCG GGC TAM CGG CCG CTG CAT CAG      376
Leu Phe Trp Lys Pro Tyr Ile Tyr Ala Gly Xaa Arg Pro Leu His Gln
                               25                               30                               35

ACC TGG CGC TTC TAT TTC CGC ACG CTG TTC CAG CAG CAC AAC              418
Thr Trp Arg Phe Tyr Phe Arg Thr Leu Phe Gln Gln His Asn
                               40                               45                               50

```

(2) INFORMATION FOR SEQ ID NO: 78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 370 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 149..232
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.4
seq LALVALAPHSVQK/SX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

```

ATGCCTCGAC CCCTTTAGAA GTTCCTTTTC CCCAGTCATC CTCTGGAACA GCCCTACCTT    60
TGGAAGCTGC CCCTGAAGCC CCAAHCTTC CTAMCCARAC CTAATAGGGC CTCCCATCTC    120
CCCAGCTGCC TTAGCTCTAG CCTCTCCC ATG ATA GCT CCA ACT CTG AAA GGG      172
                               Met Ile Ala Pro Thr Leu Lys Gly
                               -25

ACC CCT TCC TCT TCA GCT CCC TTA GCT CTG GTT GCC CTG GCT CCC CAC      220
Thr Pro Ser Ser Ser Ala Pro Leu Ala Leu Val Ala Leu Ala Pro His
-20                      -15                      -10                      -5

TCA GTT CAG AAG AGT TYN VCT TTT CCA CCT AAC CTT CTT ACT TCA CCT      268
Ser Val Gln Lys Ser Xaa Xaa Phe Pro Pro Asn Leu Leu Thr Ser Pro
                      1                      5                      10

CCT TCA GTG GCT GHA GCT GAG TCA GGG TCA GTG ATA ACT CTG TCA GCT      316
Pro Ser Val Ala Xaa Ala Glu Ser Gly Ser Val Ile Thr Leu Ser Ala
                      15                      20                      25

SCC ATT GCT CCC TCA GAA CCA AAG ACT AAT CTT AAT AAA GTT CCC TCT      364
Xaa Ile Ala Pro Ser Glu Pro Lys Thr Asn Leu Asn Lys Val Pro Ser
                      30                      35                      40

GAG GTA                                                                370
Glu Val
45

```

(2) INFORMATION FOR SEQ ID NO: 79:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 208 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 86..196
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.3
seq LVESLCLVFNNLLS/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

AGATATGTCA	ACACCTAAGA	ATCAGAACTA	GTATCTTGAA	AAGTTAGGAG	CCCTGGGGTT	60										
TTGTTTTTGC	TTTTGCTTTT	AAAGG	ATG	TGT	TTG	TTC	CCT	GTA	TCA	CCG	TGC	112				
			Met	Cys	Leu	Phe	Pro	Val	Ser	Pro	Cys					
					-35					-30						
CCA	GCT	TAC	TCC	TTT	TCT	TCG	GAA	ASR	STT	GGT	GCC	GTA	TTG	TTA	CTG	160
Pro	Ala	Tyr	Ser	Phe	Ser	Ser	Glu	Xaa	Xaa	Gly	Ala	Val	Leu	Leu	Leu	
			-25				-20					-15				
GTT	GAA	TCT	CTG	TGT	TTG	GTT	TTT	AAT	TTG	TTA	TCT	TTG	CCC	CCA	AGG	208
Val	Glu	Ser	Leu	Cys	Leu	Val	Phe	Asn	Leu	Leu	Ser	Leu	Pro	Pro	Arg	
		-10					-5					1				

(2) INFORMATION FOR SEQ ID NO: 80:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 412 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 140..184
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.3
seq IAVLFCFLLLIIF/QT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

AAACATTCCC	TTCTGTCCTT	TCTTTGTTTT	TAAAGAAAGC	TCTGATTTTG	TTTCATTTTC	60
AGCTGGAGAC	TTAAATGACA	CCAAGCAAAG	CCTACTTAGT	TTAGATCTCC	AGAAATTGGC	120
TGGTGGA	AVN	RAATCAAAC	ATG AAG ATT	GCA GTT TTG	TTT TGT TTT TTG CTG	172
			Met Lys Ile	Ala Val Leu	Phe Cys Phe Leu Leu	
			-15	-10	-5	

```

CTT ATC ATT TTT CAA ACT GAC TTT GGA AAA AAT GAA GAA ATT CCT AGG 220
Leu Ile Ile Phe Gln Thr Asp Phe Gly Lys Asn Glu Glu Ile Pro Arg
      1              5              10

AAG CAA AGG AGG AAG ATC TAC CAC AGA AGG TTG AGG AAA AGT TCA ACV 268
Lys Gln Arg Arg Lys Ile Tyr His Arg Arg Leu Arg Lys Ser Ser Thr
      15              20              25

TCA CAC AAG CAC AGA TCA AAC AGA CAG CTT GGA ATT CCG CAA ACA ACA 316
Ser His Lys His Arg Ser Asn Arg Gln Leu Gly Ile Pro Gln Thr Thr
      30              35              40

GTT TTT ACA CCA GTA GCA AGA CTT CCT ATT GTT AAC TTT GAT TAT AGC 364
Val Phe Thr Pro Val Ala Arg Leu Pro Ile Val Asn Phe Asp Tyr Ser
      45              50              55              60

ATG GAG GAA AAG TTT GAA TCC TTT TCA AGT TTT CCT GGA GTA GAA TCA 412
Met Glu Glu Lys Phe Glu Ser Phe Ser Ser Phe Pro Gly Val Glu Ser
      65              70              75

```

(2) INFORMATION FOR SEQ ID NO: 81:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 169 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 80..160
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq VGAVLLSSLPISP/QY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

```

ACCCTACCAT CCATACCCTC CATTCTTACT CCGTGTCTATG GAGAAACCTG CATAGTGCTG 60

GGGATTAGTG CCACAATTA ATG TGT AGT CCG AGG TCT CCC TTA AAT CTG TCT 112
      Met Cys Ser Pro Arg Ser Pro Leu Asn Leu Ser
      -25              -20

TTG GTC CCT GTC GGA GCA GTT CTG CTT AGC TCC CTC CCC ATT TCT CCA 160
Leu Val Pro Val Gly Ala Val Leu Leu Ser Ser Leu Pro Ile Ser Pro
      -15              -10              -5

CAG TAC GGG 169
Gln Tyr Gly
      1

```

(2) INFORMATION FOR SEQ ID NO: 82:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 230 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 24..167
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq EVVTLPLTSHCLA/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

```

ATGAATTCT AAAACTCTTT TTT ATG GGA CTT CAC ATT TCT CTG ATT AAA TTT    53
      Met Gly Leu His Ile Ser Leu Ile Lys Phe
                        -45                      -40

CTT CTT GCT AAT GGA CCC CAT ATT CCT AGT CAC CAA AGA CCT TTT GAA    101
Leu Leu Ala Asn Gly Pro His Ile Pro Ser His Gln Arg Pro Phe Glu
      -35                      -30                      -25

CCT AAA GGG GAA AAA AGC TGC AGA ATT GAA GTG GTG ACT CTG CCA CTT    149
Pro Lys Gly Glu Lys Ser Cys Arg Ile Glu Val Val Thr Leu Pro Leu
      -20                      -15                      -10

ACT AGC CAT TGT CTT GCC CAA GTT GCA AGT TCT GAC CTC ATC CAT AGG    197
Thr Ser His Cys Leu Ala Gln Val Ala Ser Ser Asp Leu Ile His Arg
      -5                      1                      5                      10

ATG AGA ACC ATA ACA GGT ACC TCG TCA CAC GGG    230
Met Arg Thr Ile Thr Gly Thr Ser Ser His Gly
      15                      20

```

(2) INFORMATION FOR SEQ ID NO: 83:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 349 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 158..235
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.1
seq LLTLYVFVASSMQ/IY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

```
ATACATGGAG GCATGTAATG AATACTGAAT GAATAAATTA TTAAACCTC AAGCTTATTC   60
TAGGCTTAAG CTCTGGTTCT CAATAAAATA CTACCAAATA AACATAAAAA TACGTACTTA  120
CTTTAGAGCA TTTTGAAAGT ACATAACTTA AGGAAAA ATG AAA ACT ACC TAT GTA   175
                               Met Lys Thr Thr Tyr Val
                               -25

ATA TTT ATG CAA AGC AAA GCA CTA TTA ACA TTG TAT GTA TTT GTA GCC   223
Ile Phe Met Gln Ser Lys Ala Leu Leu Thr Leu Tyr Val Phe Val Ala
-20                      -15                      -10                      -5

TCT TCT ATG CAA ATT TAT GTA TTA CAC ATT TCA AAT TAC CCA ACA GAT   271
Ser Ser Met Gln Ile Tyr Val Leu His Ile Ser Asn Tyr Pro Thr Asp
                      1                      5                      10

GAG CAT TTT CCT ATC ATT AAG CAT TTT TAT TTT ACT TTT AAA ATC CAC   319
Glu His Phe Pro Ile Ile Lys His Phe Tyr Phe Thr Phe Lys Ile His
          15                      20                      25

TTT AGT AAA ATT ATT TAT GTG CAG TAC AGT                               349
Phe Ser Lys Ile Ile Tyr Val Gln Tyr Ser
          30                      35
```

(2) INFORMATION FOR SEQ ID NO: 84:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 142 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 68..112
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.1
seq ALVFLIFLRFINI/SE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

AATTACTCCC TTCACTGGAA AGAATAAAAT ATTACCACAT AAATTCGTAT TATGATAATG 60

GCCCACT ATG AAT GCT TTG GTG TTC TTA ATT TTC TTA AGA TTT ATT AAT 109
 Met Asn Ala Leu Val Phe Leu Ile Phe Leu Arg Phe Ile Asn
 -15 -10 -5

ATT TCT GAA GTA ACT ACT AAA TGC CAA GCA GGG 142
 Ile Ser Glu Val Thr Thr Lys Cys Gln Ala Gly
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 190 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 86..172
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq WGFLLTGHSLSHS/SK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

AATTCCATCT CAGTTGCTGG TTTCACAGGC AGGACCACCC CTGGGCGCCT CTGTCCCCCG 60

GTCGGGGAGT CTGATCCTGC CTTCC ATG CAG CTG GGT CCC CTT CAC ACT GTG 112
 Met Gln Leu Gly Pro Leu His Thr Val
 -25

TCT ACA CCC TTT TTC TTC TGT TGG GGT TTT CTC CTC ACT GGC CAT TCT 160
 Ser Thr Pro Phe Phe Phe Cys Trp Gly Phe Leu Leu Thr Gly His Ser
 -20 -15 -10 -5

CTT TCY CAC TCC TCT AAG TCC TGC CAC CTG 190
 Leu Ser His Ser Ser Lys Ser Cys His Leu
 1 5

(2) INFORMATION FOR SEQ ID NO: 86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 226 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 41..211
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.1
seq VGSVCCCVGPLRG/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

```
CTCTAGGACT GGCTTGGTGG AGGTBRATGA AGGCCGTGAG ATG GGG CGG GGG TGG      55
                                         Met Gly Arg Gly Trp
                                         -55

GAG AGG ACA GTT TGC AGC CTT GGG TGG CGC GGG GGC CCA GAC CCT CTG      103
Glu Arg Thr Val Cys Ser Leu Gly Trp Arg Gly Gly Pro Asp Pro Leu
-50                      -45                      -40

TCT TGG GCC ACT TGT TGG TCG GGA GCC AGA TCT CGC CAC ACC CGC GTT      151
Ser Trp Ala Thr Cys Trp Ser Gly Ala Arg Ser Arg His Thr Arg Val
-35                      -30                      -25

TCC TCG ATT GTA AAC GGT TAT GTT GGG AGT GTT TGC TGC TGC GTG GGT      199
Ser Ser Ile Val Asn Gly Tyr Val Gly Ser Val Cys Cys Cys Val Gly
-20                      -15                      -10                      -5

CCG CTG AGA GGA TTA GTC TGG GCC CCT      226
Pro Leu Arg Gly Leu Val Trp Ala Pro
                      1                      5
```

(2) INFORMATION FOR SEQ ID NO: 87:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 195 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 115..180
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5
seq HLFVTWSSQRALS/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

```

AACCTGCCAG TKATGCAAAT GCCAAAATGT GGGTCATCAT ATAGTATATT TGAAACCTTT    60
CTGAACATGT ACACCACCCA ATGCTAGAGG CTGACTTGGA AACCGGTGGG TGCA ATG    117
                                         Met
CCC GAG GCT GTG GAA CAA TCA GCC CAT CTC TTT GTG ACC TGG AGC AGT    165
Pro Glu Ala Val Glu Gln Ser Ala His Leu Phe Val Thr Trp Ser Ser
   -20                      -15                      -10

CAG AGG GCC CTC AGT CAC CCC GCC CCA TTG                                195
Gln Arg Ala Leu Ser His Pro Ala Pro Leu
   -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 386 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 240..311
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5
seq CWLIALSVPLVFW/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

```

AATTTGTTTA GAAACATGCC TTTCATTAAT GTAATCCCT GCATTTACAG ATAGACAATC    60
TGAGCCAGGA CTAGCAGTCC AGGATTCCKG ATGTCTTACC TTGTCCTTAA CTCTGATACC    120
ATCCTACTGA CCAAAGTTNG CACTTCTTTG GACTTTTTTA GAGACTACCA AATGGGCTTT    180
GACACGTTCA TTTGGAATAC ATCAAGCCTT AATATTTATA TCTTATATCA TTGAACATA    239
ATG CCT GGC ACA CAC ACG TTT ACT TTT AAA AGC TGC TGG CTT ATA GCT    287
Met Pro Gly Thr His Thr Phe Thr Phe Lys Ser Cys Trp Leu Ile Ala
   -20                      -15                      -10

CTT TCT GTG CCT TTG GTC TTT TGG GTG ACA TTC TGG CCG TGT AAC TTT    335
Leu Ser Val Pro Leu Val Phe Trp Val Thr Phe Trp Pro Cys Asn Phe
   -5                      1                      5

TAT CCG TCC CTT GAC TTC TGC ATG TTA ACT AAG RCT AAG AGC ATA TTC    383
Tyr Pro Ser Leu Asp Phe Cys Met Leu Thr Lys Xaa Lys Ser Ile Phe

```

10

15

20

ATA
Ile
25

386

(2) INFORMATION FOR SEQ ID NO: 89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 197 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 69..134
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq LFCLIGLDLLCQV/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

```
ATAAATCTCA CAATTAGAAT TGCATAATTG ATTCACATCA ATAATAGACC TATATTTGAT    60
GTTTCAGCT ATG CTT TTA CTG ACT TTC AAA TGG TTT CTG TTT TGC TTA ATT    110
      Met Leu Leu Leu Thr Phe Lys Trp Phe Leu Phe Cys Leu Ile
                -20                -15                -10
GGT TTG GAT TTG CTC TGT CAG GTT TTT TCC CCT TAT TTC CTT AGT GAG    158
Gly Leu Asp Leu Leu Cys Gln Val Phe Ser Pro Tyr Phe Leu Ser Glu
                -5                1                5
AAA GTT GCA GAC CTG CTG TTT TAT ATG TCT CTG TTT TTT    197
Lys Val Ala Asp Leu Leu Phe Tyr Met Ser Leu Phe Phe
      10                15                20
```

(2) INFORMATION FOR SEQ ID NO: 90:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 169 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 5..160
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq VPNLHLLLPLTTP/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

```

ATTG ATG GCA ACY ACA GGC AGA CGG CAG GCT GAA CCT CCG CCC GTC CGG      49
Met Ala Thr Thr Gly Arg Arg Gln Ala Glu Pro Pro Pro Val Arg
      -50                      -45                      -40

CCC GCC CAT TCC CGA CCT CCA CCT AGG GTG CCT GGG AGC AGC AGT CTA      97
Pro Ala His Ser Arg Pro Pro Pro Arg Val Pro Gly Ser Ser Ser Leu
      -35                      -30                      -25

GGG CTG GCA GGA CTT ATG TCC CCC GTC CCC AAC CTT CAC CTA CTC CTC      145
Gly Leu Ala Gly Leu Met Ser Pro Val Pro Asn Leu His Leu Leu Leu
      -20                      -15                      -10

CCC CTT ACT ACT CCC CAA CCT CGG      169
Pro Leu Thr Thr Pro Gln Pro Arg
      -5                      1

```

(2) INFORMATION FOR SEQ ID NO: 91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 185 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 27..74
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq FIYLAQHFTLCSG/WS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

```

AATAGATAGG AGAGCAAGCC TCACCA ATG GTT CCC TTC ATC TAT CTG CAA GCC      53
Met Val Pro Phe Ile Tyr Leu Gln Ala
      -15                      -10

CAC TTT ACA CTC TGT TCT GGG TGG TCC AGC ACA TAC CGG GAC CTC CGG      101
His Phe Thr Leu Cys Ser Gly Trp Ser Ser Thr Tyr Arg Asp Leu Arg
      -5                      1                      5

```

AAG GGT GTG TAT GTG CCC TAC ACC CAG GGC AAG TGG GAA GGG GAG CTG 149
 Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu
 10 15 20 25

GGC ACC GAC CTG GTA AGC ATC CCC CAT GGC CCA AAG 185
 Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Lys
 30 35

(2) INFORMATION FOR SEQ ID NO: 92:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 239 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 150..191
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8
seq FFSFLLTINLVSL/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

ATAACATTAC AGTTTGGGCT CTTGCTGCCC AATGATTGAT TTATCTATAG TATAGATTTA 60
 TTTCTCACAG TACCTCTTGG AATGCTCATT TTTAACCCCA ATAGTTAAAT TTGCCTTGCT 120
 AAGCTACAAA AACAGGCACC AAAGCAGCA ATG TTT TTT AGT TTT CTG TTG ACC 173
 Met Phe Phe Ser Phe Leu Leu Thr
 -10
 ATA AAT CTC GTT TCT TTA CAA GTA GTA ATT CTA AAC AGA GTA TAC CTT 221
 Ile Asn Leu Val Ser Leu Gln Val Val Ile Leu Asn Arg Val Tyr Leu
 -5 1 5 10
 AAC CAG CCA GAT GCA CGG 239
 Asn Gln Pro Asp Ala Arg
 15

(2) INFORMATION FOR SEQ ID NO: 93:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 168 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 100..162
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.8
seq NWGLLCFASCTT/DR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

```
AGAGGATTGG AGAGGAAGAG TTGGATTCCA GAAATATGCA GGAGGTGGCC GCAGGAATTG    60
GGAGATAAGG CAGAGGGAAG AGGCCATGCC CAGTTTCTG ATG TGG CCT GGG CGT    114
                                     Met Trp Pro Gly Arg
                                     -20
GAA TGC AAA AAT TGG GGT CTT CTG TGC TTT GCA AGT GAG TGC ACC ACC    162
Glu Cys Lys Asn Trp Gly Leu Leu Cys Phe Ala Ser Glu Cys Thr Thr
-15                      -10                      -5
GAT CGG    168
Asp Arg
1
```

(2) INFORMATION FOR SEQ ID NO: 94:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 238 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 113..223
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.8
seq CLLTLRQPPTHG/GI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

```
AAGTTTATTT ATCAAGTGAA ATAGGGTTAG TACTACCAGT CTCATAAGGT TATTATGAGA    60
TACCAAATGG AAATCTAAAG AAGATCYTTG CACCATAGTA TGTGATCAGT GA ATG TTA    118
                                     Met Leu
```

ACT CCT TTC TCC TTG GAG GAG AAA CTT CTA GAA TGT CAT TAT GTA CTT 166
 Thr Pro Phe Ser Leu Glu Glu Lys Leu Leu Glu Cys His Tyr Val Leu
 -35 -30 -25 -20

GCA AAA CTA GCT GGG GCA TGT CTC CTG TTG ACT CTA AGG CAG CCA CCC 214
 Ala Lys Leu Ala Gly Ala Cys Leu Leu Leu Thr Leu Arg Gln Pro Pro
 -15 -10 -5

ACA CAT TCT GGA ATC CCA CAC GGG 238
 Thr His Ser Gly Ile Pro His Gly
 1 5

(2) INFORMATION FOR SEQ ID NO: 95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 204 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 79..198
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7
seq IVFVGLIFLKSSA/HR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

ATGATTGMGG GATTGTGTAG AGGTGATTTT GAAGATGGAA GACTTGTGCA CTGAAGAAAA 60

TGAGAAAAAT GAGAAGAA ATG AAA AGA ATA AAA TCA ATG ATG GGA AAA GTT 111
 Met Lys Arg Ile Lys Ser Met Met Gly Lys Val
 -40 -35 -30

GAA CAT ATA AAG ATT AAA GGA GAA AAA CAA AGA AGC CGT CAT GTA AAA 159
 Glu His Ile Lys Ile Lys Gly Glu Lys Gln Arg Ser Arg His Val Lys
 -25 -20 -15

ATA GTA TTT GTT GGG CTT ATT TTT CTA AAA AGC AGT GCA CAT AGG 204
 Ile Val Phe Val Gly Leu Ile Phe Leu Lys Ser Ser Ala His Arg
 -10 -5 1

(2) INFORMATION FOR SEQ ID NO: 96:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 136 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 29..79

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.7
seq LCLFCKICPPTHG/VA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

```
AACCAGAAAC TACTTTGCAT CCCATTGA ATG CAA TCT GCT CTG TGC CTC TTT      52
                               Met Gln Ser Ala Leu Cys Leu Phe
                               -15                      -10

TGT AAA ATC TGC CCA TTT ACA CAT GGT GTT GCC ACC CCA GCC TGG GAA      100
Cys Lys Ile Cys Pro Phe Thr His Gly Val Ala Thr Pro Ala Trp Glu
          -5                      1                      5

CTG AGC AGC AAG AGG AAA GCT TCC CAC CCG CCC CGG                      136
Leu Ser Ser Lys Arg Lys Ala Ser His Pro Pro Arg
      10                      15
```

(2) INFORMATION FOR SEQ ID NO: 97:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 387 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Surrenals

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 112..318

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.7
seq SLLCLAFLLGRFL/HM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

```
AGCTAGTAGC AGCTCTGGCA GAAGCAACGG TGGCTTCGAG GGATGGCGGC GGCTGCAACA      60

GGACCTGCAG CATCCCAGAG GAACTGACTA AGACTTTGGA ACAGAAACCA G ATG ATG      117
                               Met Met
```

CAC AAT ATT ATC GTC AAA GAG CTT ATT GTC ACA TTC TTC TTG GGA ATT	165
His Asn Ile Ile Val Lys Glu Leu Ile Val Thr Phe Phe Leu Gly Ile	
-65 -60 -55	
ACT GTG GTG CAG ATG CTA ATT TCA GTG ACT GGA TTA AAA GGT GTC GAA	213
Thr Val Val Gln Met Leu Ile Ser Val Thr Gly Leu Lys Gly Val Glu	
-50 -45 -40	
GCT CAG AAT GGC TCG GAA TCT GAG GTG TTT GTG GGG AAG TAT GAG ACC	261
Ala Gln Asn Gly Ser Glu Ser Glu Val Phe Val Gly Lys Tyr Glu Thr	
-35 -30 -25 -20	
CTC GTG TTT TAC TGG CCC TCG CTG CTG TGC CTT GCC TTC CTG CTG GGC	309
Leu Val Phe Tyr Trp Pro Ser Leu Leu Cys Leu Ala Phe Leu Leu Gly	
-15 -10 -5	
CGC TTC CTG CAT ATG TTT GTC AAG GCT CTG AGG GTG CAC CTC GGC TGG	357
Arg Phe Leu His Met Phe Val Lys Ala Leu Arg Val His Leu Gly Trp	
1 5 10	
GAG CTC CAG GTG GAA GAA AAA TCT GTC CTG	387
Glu Leu Gln Val Glu Glu Lys Ser Val Leu	
15 20	

(2) INFORMATION FOR SEQ ID NO: 98:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 324 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 19..99
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6
seq RLLCSRLCQQLRS/KR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

ATCATCTGAA TCTTAATC ATG TCC AAC TGC CTG CAA AAT TTC CTG AAA ATT	51
Met Ser Asn Cys Leu Gln Asn Phe Leu Lys Ile	
-25 -20	
ACA AGC ACT CGT CTT CTA TGT TCA AGA TTA TGC CAA CAG TTA AGA AGT	99
Thr Ser Thr Arg Leu Leu Cys Ser Arg Leu Cys Gln Gln Leu Arg Ser	
-15 -10 -5	
AAA AGG AAG TTT TTC GGA ACT GTG CCA ATA TCC AGA TTG CAT AGG CGA	147
Lys Arg Lys Phe Phe Gly Thr Val Pro Ile Ser Arg Leu His Arg Arg	
1 5 10 15	

```

GTT GTC ATT ACA GGC ATT GGC TTA GTG ACT CCT CTT GGT GTT GGA ACT   195
Val Val Ile Thr Gly Ile Gly Leu Val Thr Pro Leu Gly Val Gly Thr
      20              25              30

CAC CTG GTT TGG GAT CGT CTT ATC GGA GGA GAG AGT GGA ATT GTT TCA   243
His Leu Val Trp Asp Arg Leu Ile Gly Gly Glu Ser Gly Ile Val Ser
      35              40              45

CTG GTT GGT GAA GAG TAT AAG AGT ATC CCT TGC AGT GTT GCT GCT TAT   291
Leu Val Gly Glu Glu Tyr Lys Ser Ile Pro Cys Ser Val Ala Ala Tyr
      50              55              60

GTG CCA AGA GGT AGT GAT GAA GGT CAG TCA GGG   324
Val Pro Arg Gly Ser Asp Glu Gly Gln Ser Gly
      65              70              75

```

(2) INFORMATION FOR SEQ ID NO: 99:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 241 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 74..121
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6
seq XGLFLRTTAAARA/CR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

```

AATCAGACGT GTGTGTGTCC CTGCGGCGCT AAGAAGGGGA GACTGAGGCT GMGGCTGGGG   60

AACATCGGGC AGC ATG AGC GGC NCM GGG CTC TTC CTG CGC ACC ACG GCT   109
      Met Ser Gly Xaa Gly Leu Phe Leu Arg Thr Thr Ala
      -15              -10              -5

GCG GCT CGT GCC TGC CGG GGT CTG GTG GTC TCT ACC GCG AAC CGG CGG   157
Ala Ala Arg Ala Cys Arg Gly Leu Val Val Ser Thr Ala Asn Arg Arg
      1              5              10

CTA CTG CGC ACC AGC CCG CCT GTA CGA GCT TTC GCC AAA GAG CTT TTC   205
Leu Leu Arg Thr Ser Pro Pro Val Arg Ala Phe Ala Lys Glu Leu Phe
      15              20              25

CTA GGC AAA ATC RAG AAG GTA ACG CGA GCC CTG GGC   241
Leu Gly Lys Ile Xaa Lys Val Thr Arg Ala Leu Gly
      30              35              40

```

(2) INFORMATION FOR SEQ ID NO: 100:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 112..186
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq LTWLHLLLSHLKS/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

```

ACCTTCCCAC CACCCGCAGG AGCACCTGCC TCCATTACACA ACCGCCGACA CTACTGCAGC      60
CTCATCCTAA CTGACCTTGG CTTCTCTGGG AGACCCTTCC TGCACCCTGA A ATG AAT      117
                                     Met Asn
                                     -25

CCT CTG CAC AAG CAC TGT GCT GCA GGA CCC CTC ACC TGG CTC CAC CTC      165
Pro Leu His Lys His Cys Ala Ala Gly Pro Leu Thr Trp Leu His Leu
      -20                      -15                      -10

CTA CTT TCT CAC TTG AAG TCT TCT CTT GTA ATG CCT CCA AAG      207
Leu Leu Ser His Leu Lys Ser Ser Leu Val Met Pro Pro Lys
      -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 143 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 18..104

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.5
seq FAVLRVLHLPALT/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

```

AACACATCTT CCCTGAG ATG CCT AAG GAC AAA AGA GGA GCT AGA CAC AAC      50
              Met Pro Lys Asp Lys Arg Gly Ala Arg His Asn
              -25                      -20

TCT CCC CAT TTT TCC TTT GCT GTC TTA AGA GTG CTC CAT CTT CCA GCA      98
Ser Pro His Phe Ser Phe Ala Val Leu Arg Val Leu His Leu Pro Ala
      -15                      -10                      -5

CTG ACT GCC CCT CTG TGG CTG GCT CCT TTC TCT ACC CTC CCC AGG      143
Leu Thr Ala Pro Leu Trp Leu Ala Pro Phe Ser Thr Leu Pro Arg
      1                      5                      10

```

(2) INFORMATION FOR SEQ ID NO: 102:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 269 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 126..242
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq LCVSRQLLTGART/LF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:

```

AAGAACAAGC AGCTGACATG ATTGCTGTCG GTGTTACAG AGCCAAATAT GAAGATAATT      60

GTAAAGAAAG AGACGGTCGC ATCGGATAGA AGATGTGATC CTGCTCTCAC GTTTTTCCTT      120

CTGGC ATG ACC ATA CAC GTT TTG AGA AAA TGT TGC CAA ATG GGT AGA CTA      170
Met Thr Ile His Val Leu Arg Lys Cys Cys Gln Met Gly Arg Leu
      -35                      -30                      -25

AAC AAT GAA TGG CTG CCG GGT TTA GTC ATA CCT CTC TGT GTG AGC CGT      218
Asn Asn Glu Trp Leu Pro Gly Leu Val Ile Pro Leu Cys Val Ser Arg
      -20                      -15                      -10

CAA TTG CTG ACG GGA GCT AGG ACA TTA TTC CAG CTA CAA AAT GGG CCC      266
Gln Leu Leu Thr Gly Ala Arg Thr Leu Phe Gln Leu Gln Asn Gly Pro
      -5                      1                      5

```

GCG
Ala

269

(2) INFORMATION FOR SEQ ID NO: 103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 253..342
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq IAALLGLLQLRFK/AE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

```

GTCCAGATTG GGCCACTTCT TTCTCAGCTC TAATGACTTT CCTCAGTTCC GTGGGTTACT    60
CCTGCCAACT CGACGCCGGC CGCCATGACA CTCGCTCGGA AAGCGGCAGC GNATCATAGA    120
AAAGCGCCGC GGTGGCGTAG ACAGGCCCCG CGAASC GCCG GACGTGTCCT TGGCGCAAGG    180
GAGGCTGGGA TCGCGGAGGA CCGAGCGCGG GCTGGATTAA CCGCAGCCAG TGCCTAGCGC    240
AAGGTTAGGT GC ATG CAG GCG GCC AGC TTC GGC CGG GGA AGG AAT GGC CTG    291
          Met Gln Ala Ala Ser Phe Gly Arg Gly Arg Asn Gly Leu
          -30                -25                -20

GAT AAC TGG GGC ATC GCG GCG CTC CTC GGC CTC CTG CAG CTG CGT TTC    339
Asp Asn Trp Gly Ile Ala Ala Leu Leu Gly Leu Leu Gln Leu Arg Phe
          -15                -10                -5

AAA GCA GAG
Lys Ala Glu
      1

```

348

(2) INFORMATION FOR SEQ ID NO: 104:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 465 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 310..438
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.5
seq ENLLLCCHRCTNC/QR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

```

AAGGTTCTTT CTCATTTGTG TAGGTGGATG TTCCTTCAAT CTTTGAAGTT GCTGTCCCTT   60
GGATGGATTT TTGTTGCTTT CATCTTCTTT GATGCCCTTG GGGGCTTGAT TGTGGTATAA  120
GGTAGGTTCA GMHAACTCAT TTAAGKTCGC TTTTCCAGGA GTCACTGGGG ACCAGGAATG  180
AGTCCTGGTG CATGGTAATC CCATGCAGAG TTCCAGCCA TCCCTCTGTC AGCCAGCAC   240
CTGTGTCTTC CCACATCCAC TGTCAATGCC TTCCCTCTGA GATTGCTCA GAGTGC GCCT   300
GTCTTCGCA ATG TCC CCA TCT CTT GGT GAC AGA TGT TCC TCT TGG CTG CAT   351
      Met Ser Pro Ser Leu Gly Asp Arg Cys Ser Ser Trp Leu His
                -40                      -35                      -30

CTA GTC AGC CAT CTT GAA TCA ATC TCT GGA CCT CTA CTT AAT ATC CCT   399
Leu Val Ser His Leu Glu Ser Ile Ser Gly Pro Leu Leu Asn Ile Pro
                -25                      -20                      -15

GAG AAT CTC CTT CTC TGT TGT CAC CGG TGC ACC AAC TGC CAA AGA CAC   447
Glu Asn Leu Leu Leu Cys Cys His Arg Cys Thr Asn Cys Gln Arg His
                -10                      -5                      1

CAT TTT TGC AGT GTT TGG   465
His Phe Cys Ser Val Trp
      5

```

(2) INFORMATION FOR SEQ ID NO: 105:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 141 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 1..72
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.4

seq YFLLPCLINLAIG/VK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:

ATG TCA GGA GCT GAG CCA ACA ACC TTC ATC AGA TAT TTT CTT CTT CCA	48
Met Ser Gly Ala Glu Pro Thr Thr Phe Ile Arg Tyr Phe Leu Leu Pro	
-20 -15 -10	
TGT CTC ATA AAT TTG GCC ATT GGA GTT AAA TGG AAA ACA GCT TGG AAG	96
Cys Leu Ile Asn Leu Ala Ile Gly Val Lys Trp Lys Thr Ala Trp Lys	
-5 1 5	
AGG GGG GAA AGG CAG CTA AAC AAC ACT GTC TTT TTT TTT TTT TTT	141
Arg Gly Glu Arg Gln Leu Asn Asn Thr Val Phe Phe Phe Phe Phe	
10 15 20	

(2) INFORMATION FOR SEQ ID NO: 106:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 186 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 91..156
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4
seq QLLFSFLLSTIPT/SY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:

ATGTTACTAG GCAATAGGAG TTTTGAAC GCATTATAAT CTTATGAGAC CACCATCATA	60
TATACAGTTT GTTGTTGACC AAAACGTTAT ATG GTG TAT GAC TAT TTT ATT TCC	114
Met Val Tyr Asp Tyr Phe Ile Ser	
-20 -15	
CAA CAA CTG CTG TTC TCT TTT TTA CTC TCT ACT ATC CCC ACA TCT TAC	162
Gln Gln Leu Leu Phe Ser Phe Leu Leu Ser Thr Ile Pro Thr Ser Tyr	
-10 -5 1	
CAC CTT TCC CTT ACT TGC CAG CGG	186
His Leu Ser Leu Thr Cys Gln Arg	
5 10	

(2) INFORMATION FOR SEQ ID NO: 107:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 332 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 261..302
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.3
 seq LFLCSCSLSLNQL/LT
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

```

AAATCAGATT CACCTAAAC GGAGATCATT TATAGAAAGA AAAAAACACA AAGGTCTTCT   60
GTTGAACTGT TCAGTAAGTG TGTACTGATA CAATAATATG GCAGGATATT GGGTTTCCAA  120
CTCAGAGTTG ATAGTTCAGA ATTAAATTTT GGCCTATGTC TTGCTTTTAA GCTTTTAAAG  180
CCATTCTGAC GCTACCATAG ACATGCCCAT CTGGTAGATG AGAAATTCAG GTGTAGAAAG  240
ACTTGCAGAA ATCTTATGAG ATG CTC TTT CTT TGT TCC TGT TCT CTT TCT CTG  293
                Met Leu Phe Leu Cys Ser Cys Ser Leu Ser Leu
                -10                               -5

AAC CAG CTC CTA ACT TAC ATC TTT GTA GTC CCA CCC TGG                332
Asn Gln Leu Leu Thr Tyr Ile Phe Val Val Pro Pro Trp
                1                               5                               10

```

(2) INFORMATION FOR SEQ ID NO: 108:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 187 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 122..166
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.2

seq FLMVLLFRSNKWT/GK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

```

AAAGAAAATG AACCCCTGTA AAGCAATTC AGCTTATGAA CCCCTGTAAA GCAATTCAG   60
CTTTGCTTCA TGGTATATTT GCTTTCTATC CAGAGGTAA TTTGTTAGTT TTCTTAAAC   120
A ATG TTT TTC TTA ATG GTC TTA TTG TTT AGA AGT AAC AAG TGG ACT GGA   169
  Met Phe Phe Leu Met Val Leu Leu Phe Arg Ser Asn Lys Trp Thr Gly
   -15                -10                -5                1
AAA GTA TAC GGG GCC CTG                                           187
Lys Val Tyr Gly Ala Leu
                   5

```

(2) INFORMATION FOR SEQ ID NO: 109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 200 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 78..164
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq FLSHVTTSLASSS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:

```

ACAACTACCA CTCTCTGTTC TGTTCACTCC GTTCCAGCCA CACCCACCTT CTTGCTGTTC   60
TTTGAACATG GCCTGGC ATG CTC CCT CTT CAG GGC CTT TGC ACT TGT TAT       110
          Met Leu Pro Leu Gln Gly Leu Cys Thr Cys Tyr
                   -25                -20
TTC CTC CAC CTA GAA TTT CTT TCC CAT GTA ACT ACC TCA CTT GCT TCA       158
Phe Leu His Leu Glu Phe Leu Ser His Val Thr Thr Ser Leu Ala Ser
   -15                -10                -5
TCA TCA GCT CCC TCA CCT AAA CCT TCA GTA ACC CTT TCT TCG           200
Ser Ser Ala Pro Ser Pro Lys Pro Ser Val Thr Leu Ser Ser
    1                5                10

```

(2) INFORMATION FOR SEQ ID NO: 110:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 276 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 205..267
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.2
 seq HFFLLNLTILLFG/CA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

```
ACAGAGAATT GTTCCACTAC ACTAAAAATC CCATGTGCTC TGCTTATTCA TCCCTCCTTC    60
TCTCCCTCTA GCCCCTGACA ACCACTGATG TCTTTACTGT CTCCCTAGTT TTGCTTTGCC    120
CAGAATGTTA TATAGATGGA ATAATATAGT ATATATTTTC ACATTGGCTT CATTCACTTA    180
GATACATGTC TTTAAGGTTC CTTC ATG TAT TTT TAT GGC TTG ACA TTT CAT      231
                Met Tyr Phe Tyr Gly Leu Thr Phe His
                -20                      -15

TTC TTC TTA TTG CTG AAT ACT ATT TTA TTG TTT GGG TGT GCC CGG          276
Phe Phe Leu Leu Leu Asn Thr Ile Leu Leu Phe Gly Cys Ala Arg
   -10                      -5                      1
```

(2) INFORMATION FOR SEQ ID NO: 111:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 271 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 152..238
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.1
 seq LWASQGSLLQDAQS/ER
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

ACTCTATTAT CTCGGCTTCT CGGGAGGAGC CTCATCTAGT CAGTCACGCA GAAGTTTCTC 60
 TTTCGCTCTT CGCGCTACAC ACCCAGATTG GCTTCCAGCG CGCAGGTAAA ACCTGGCTGT 120
 GCCCTGTTGA AATCAATTCT GTTGCACTCA T ATG CGG TGG AAT CTG TTC TTC 172
 Met Arg Trp Asn Leu Phe Phe
 -25
 TTT TGC ATC CTA CGT AAC CAG ACC AAG CTG TGG GCT TCT CAA GGT AGC 220
 Phe Cys Ile Leu Arg Asn Gln Thr Lys Leu Trp Ala Ser Gln Gly Ser
 -20 -15 -10
 CTC CAG GAT GCA CAG AGT GAG AGA GGA TGC TTT TCC CTA AAC CAG GAT 268
 Leu Gln Asp Ala Gln Ser Glu Arg Gly Cys Phe Ser Leu Asn Gln Asp
 -5 1 5 10
 GGG 271
 Gly

(2) INFORMATION FOR SEQ ID NO: 112:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 245 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 150..191
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq FIAALFTMAKTWN/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

AGTGTAATTT AGTTCAACCA TTGTGGAAGA TAGTGTGGCG ATTCCTCAAG GATCTAGAAC 60
 CAGAAATACC ATTTGACCCA GCAATCCCAT CATTGGGTAT ATACCCAAAG GATTATAAAT 120
 CATTCTAATA TAAAAACACG TGTACGCAT ATG TTT ATT GCA GCA CTA TTC ACA 173
 Met Phe Ile Ala Ala Leu Phe Thr
 -10
 ATG GCA AAG ACT TGG AAC CAA CCC GGG TGC TCA TCA ATG ATG GGC TGG 221
 Met Ala Lys Thr Trp Asn Gln Pro Gly Cys Ser Ser Met Met Gly Trp
 -5 1 5 10
 ATA AAG AAA ATG AGG CAC ATG ACG 245
 Ile Lys Lys Met Arg His Met Thr

15

(2) INFORMATION FOR SEQ ID NO: 113:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 25..156
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq LWVXLPXXXVIAS/VV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

```

ATTACATYAA ATKTATCAAA ATGY ATG CCY GGG YCA AAG CAT TTT CTC AGA      51
                        Met Pro Gly Xaa Lys His Phe Leu Arg
                        -40

GTG TTC AGA AMA TCC GCG MKG AGG AGC GTC GGA TAT KGG MAM AAG CCG      99
Val Phe Arg Xaa Ser Ala Xaa Arg Ser Val Gly Tyr Xaa Xaa Lys Pro
-35                -30                -25                -20

GGT ACT TCC AGA GCA TCA CTG TGG GTR TSG CTC CCA TTS RTG GYS GTG      147
Gly Thr Ser Arg Ala Ser Leu Trp Val Xaa Leu Pro Xaa Xaa Xaa Val
                -15                -10                -5

ATY GCC AGC GTG GTG ACC TTC TCT GKK CAT ATG ACC CTG GGC TTC GAT      195
Ile Ala Ser Val Val Thr Phe Ser Xaa His Met Thr Leu Gly Phe Asp
                1                5                10

CTG ACA GCA GCG                                          207
Leu Thr Ala Ala
15

```

(2) INFORMATION FOR SEQ ID NO: 114:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 269 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 30..176
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.1
seq VALGPLFVTGHFA/GE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

```

AAGGCTCCCC TTTCGGCCTC TCGTTCCTT ATG CGA CTT GAA TCT CCC GAC GAA      53
                               Met Arg Leu Glu Ser Pro Asp Glu
                               -45

AAT TTT GCG GTA GTT CAG GAA CAT GCC ATT CAT CAC ATC GAT GGC CCG      101
Asn Phe Ala Val Val Gln Glu His Ala Ile His His Ile Asp Gly Pro
-40                      -35                      -30

CTC CGC AGA TTC CTG CTT TTG GAA GTG CAC GAA CCC GTA GCC CTT GGG      149
Leu Arg Arg Phe Leu Leu Glu Val His Glu Pro Val Ala Leu Gly
-25                      -20                      -15                      -10

CCC CTT TTC GTC ACA GGC CAC TTT GCA GGA GAG GAT GTT GCC GAA CGC      197
Pro Leu Phe Val Thr Gly His Phe Ala Gly Glu Asp Val Ala Glu Arg
                      -5                      1                      5

CGA GAA GAT GTT GTA CAG CGC CTT GTT GTC GAT GGT CTT GCC CAG GTT      245
Arg Glu Asp Val Val Gln Arg Leu Val Val Asp Gly Leu Ala Gln Val
10                      15                      20

CTT GAT GAA GAC GTT GCC CAC CCG      269
Leu Asp Glu Asp Val Ala His Pro
25                      30

```

(2) INFORMATION FOR SEQ ID NO: 115:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 249 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 175..234
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4
seq EVLLPTVLRGSYC/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

```

ACATTAATCA ACTGTGAAAT ACAGAGCAGG TCACTTCACC TCTCAGTGTG TCCTCATTTT    60
AAAAATCAGA CCGTAACAGT AGCTATCTCA WTAGGGTTGT TAGGAGGTGT ACTGTATTAG    120
GATGTTAGGC CTTATACAWN AGAAGAAAAC GGAACAGTGA CGTAAACAAA TTTG ATG      177
                                     Met
                                     -20
GCA GGG AGT CCA GAT AGG GAG GTT CTG CTC CCG ACA GTC CTC AGA GGC      225
Ala Gly Ser Pro Asp Arg Glu Val Leu Leu Pro Thr Val Leu Arg Gly
      -15                      -10                      -5

TCA TAT TGT TTC TCC CAC CAT GGG                                249
Ser Tyr Cys Phe Ser His His Gly
      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 116:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 198 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 10..165
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4
seq RHLFLFEISLVFS/KE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

```

AATGATGCC ATG CAT GTC AGC ATG CTG GAA GGG TTC GAC GAG AAC CTG GAT    51
      Met His Val Ser Met Leu Glu Gly Phe Asp Glu Asn Leu Asp
      -50                      -45                      -40

GTG CAG GGG GAG TTG ATT CTC CAG GAT GCC TTT CAA GTG TGG GAC CCG      99
Val Gln Gly Glu Leu Ile Leu Gln Asp Ala Phe Gln Val Trp Asp Pro
      -35                      -30                      -25

AAG TCG CTG ATC CGG AAG GGG CGG GAG CGG CAC TTG TTC CTC TTT GAG     147
Lys Ser Leu Ile Arg Lys Gly Arg Glu Arg His Leu Phe Leu Phe Glu
      -20                      -15                      -10

ATC TCC TTG GTT TTT AGC AAG GAG ATC AAA GAT TCT TCA GAA CAC AAC     195
Ile Ser Leu Val Phe Ser Lys Glu Ile Lys Asp Ser Ser Glu His Asn
      -5                      1                      5                      10

```

GGG
Gly

198

(2) INFORMATION FOR SEQ ID NO: 117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 306 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 112..228
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9
seq VLAIGLLHIVLLS/IP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

```

AGCAGCCGCG GGTGTTACA GCTGCTGGAG CAGCAGCGGC CCCGCTCCC GGAACCGTT    60
CCCGGGCCGT TGATCTTCGG CCCACACGA ACAGCAGAGA GGGGCANMAG G ATG AAT    117
                               Met Asn

GTG GGC ACA GCG CAC AGC GAG GTG AAC CNC AAC ACG CGG GTG ATG AAG    165
Val Gly Thr Ala His Ser Glu Val Asn Xaa Asn Thr Arg Val Met Lys
      -35                      -30                      -25

GNC CGT GGC ATC TGG CTS TCC TAC GTG CTG GCC ATC GGT CTC CTC CAC    213
Xaa Arg Gly Ile Trp Leu Ser Tyr Val Leu Ala Ile Gly Leu Leu His
      -20                      -15                      -10

ATC GTG CTG CTG AGC ATC CCG TTT GTG AGT GTC CST GTC GTC TGG ACC    261
Ile Val Leu Leu Ser Ile Pro Phe Val Ser Val Xaa Val Val Trp Thr
      -5                      1                      5                      10

CTC ACC AAC CTC ATT CAC AAC ATG GGC ATG TAT ATC TTC CTG CAC    306
Leu Thr Asn Leu Ile His Asn Met Gly Met Tyr Ile Phe Leu His
              15                      20                      25

```

(2) INFORMATION FOR SEQ ID NO: 118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 433 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Homo Sapiens
 - (D) DEVELOPMENTAL STAGE: Fetal
 - (F) TISSUE TYPE: brain
- (ix) FEATURE:
- (A) NAME/KEY: sig_peptide
 - (B) LOCATION: 371..418
 - (C) IDENTIFICATION METHOD: Von Heijne matrix
 - (D) OTHER INFORMATION: score 3.8
seq FVXAIXXYIPTNS/VQ
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

```
ATTTCCTTCTA TGTCTTGTTT ATTTCACTTA GCATAATGTC CTCTAGGTTC ACTCATGCTG   60
TCACAACTTG GCAGGATTTT CTTCTTTTTT AAGGCTTGAA AATATTCCAT TTTGTGTGTA  120
TATGTGTATG TATGTATACA TACACACACA CATAACAAG CACACATACA CTATAATTTT  180
TTAATCCATT CAACTATTGA TCGATACTTA AATTGATTCC AAATCTTGGC TATTGTGAAT  240
AATGCTGCAA TGAACATGGG AGTACAGATA TCTCTTCAAC ATACTGAGTT CAAATCTTTT  300
GGGTAAATAC CCAGATGTGG GATTGCTGGA TCATATGATA ATTCTATTTT TAATTTTTTG  360
AGTGACVTCC ATG CTA TCA TTC GTA NTG GCT ATA KDA NTT TAC ATT CCT   409
      Met Leu Ser Phe Val Xaa Ala Ile Xaa Xaa Tyr Ile Pro
      -15                               -10                               -5

ACT AAT AGT GTA CAA GAA TTC CTT   433
Thr Asn Ser Val Gln Glu Phe Leu
      1                               5
```

(2) INFORMATION FOR SEQ ID NO: 119:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 403 base pairs
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: DOUBLE
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Homo Sapiens
 - (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
- (A) NAME/KEY: sig_peptide
 - (B) LOCATION: 284..379
 - (C) IDENTIFICATION METHOD: Von Heijne matrix
 - (D) OTHER INFORMATION: score 3.8
seq TFINITLWLGLSLC/QR
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

```

ACAGCTGGGG CTTTGTCTTC TTTATTGCTA GGAGAATGTA GCAATAGAAG TTCTCATCGC 60
CCTGTATTGC ACTTTTGGTT TTAAGGACTG GACCCAGAGT TCCTGAAAGC CAAACTCCAT 120
AAGCTGCTCA GTAAGTTCCA AGCACATAGC CGGCTKHGGG ATGCGATTCTG GTCGAGGTCT 180
GTTGAATGAA GGTAGACGCA GCAGGCAGTT TGTCCTTACC AGTGACCTGG AAGACGGTGG 240
CACTTCCTGA GTGAGCTCAC TTACCTTCCC TGAATGGTGA GGC ATG GAT GAA TAT 295
                               Met Asp Glu Tyr
                               -30

TCC TGG TGG TGC CAC GTG TTA GAG GTG GTA AAG GGT CAA ATG TTT ACT 343
Ser Trp Trp Cys His Val Leu Glu Val Val Lys Gly Gln Met Phe Thr
      -25              -20              -15

TTT ATT AAT ATT ACA TTA TGG CTT GGT TCT CTG TGT CAG CGA TTT TTC 391
Phe Ile Asn Ile Thr Leu Trp Leu Gly Ser Leu Cys Gln Arg Phe Phe
      -10              -5              1

TAT GCC TCG GGT 403
Tyr Ala Ser Gly
      5

```

(2) INFORMATION FOR SEQ ID NO: 120:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 181 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 95..163
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8
seq FIFLIQIWKTCLS/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

```

ACATGATGGA TGAATGATGT TATCTTTCAC ATGTGGCTTT CTTGCCTTGT CCTAAGTGCC 60
TGCTGTAGTC GTTGACATTT TCCAGCAAAC AGGA ATG AGG AGA AAA GGT CAA GGA 115
                               Met Arg Arg Lys Gly Gln Gly
                               -20

CAT CTA GCC TTT ATC TTC CTG ATT CAG ATT TGG AAA ACA TGC CTT TCG 163
His Leu Ala Phe Ile Phe Leu Ile Gln Ile Trp Lys Thr Cys Leu Ser
      -15              -10              -5

```

TTT TCT CCC ACC TCT GGG
 Phe Ser Pro Thr Ser Gly
 1 5

181

(2) INFORMATION FOR SEQ ID NO: 121:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 129 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 37..111
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8
seq NILFLAVSSFSMP/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

AATATTGTAA GGTACCTTAT TCCTTAGTGG TATGGA ATG TTT TTA ATC AGT GGA	54
Met Phe Leu Ile Ser Gly	
-25 -20	
CAT GTG CAT TTA ATT TAT AAC ATC CTG TTC CTG GCA GTA TCG TCT TTT	102
His Val His Leu Ile Tyr Asn Ile Leu Phe Leu Ala Val Ser Ser Phe	
-15 -10 -5	
TCC ATG CCC CTG CCC TGC CTC TAC AGG	129
Ser Met Pro Leu Pro Cys Leu Tyr Arg	
1 5	

(2) INFORMATION FOR SEQ ID NO: 122:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 99..290
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8
seq LFIVVCVICVTLN/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

```

ATTGGATTAG TAGAATTGCT TTTGTCATTC CATTGTTTTT ATATATTTGT TTGGGACATT    60
TTACTTTTTT CTGTTAACGC TTACCCTAGR AATTAGAA ATG ACA CCA CGT ATT CTT    116
                                     Met Thr Pro Arg Ile Leu
                                     -60

AGC GAA GTC CAG TTT TCA GCA TTT TGT CCT TAT TGG ACA ATA GCA AGG    164
Ser Glu Val Gln Phe Ser Ala Phe Cys Pro Tyr Trp Thr Ile Ala Arg
      -55                      -50                      -45

ATA TTA GAA CGT GTT GGT TCC GCG TGC TTC CGT CTT GAG TTA TGT GCT    212
Ile Leu Glu Arg Val Gly Ser Ala Cys Phe Arg Leu Glu Leu Cys Ala
      -40                      -35                      -30

GCT ATT GTC GGA TAT TTT GTC TTA GAT GTA CGT ACT TTC CTG TTC ATT    260
Ala Ile Val Gly Tyr Phe Val Leu Asp Val Arg Thr Phe Leu Phe Ile
      -25                      -20                      -15

GTG GTA TGT GTA ATT TGC GTT ACT TTG AAT TTT CCA CGT THN KAC TTT    308
Val Val Cys Val Ile Cys Val Thr Leu Asn Phe Pro Arg Xaa Xaa Phe
      -10                      -5                      1                      5

CTT TGT CTC TCA TCA CTT ACC GCG    332
Leu Cys Leu Ser Ser Leu Thr Ala
      10

```

(2) INFORMATION FOR SEQ ID NO: 123:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 225 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 25..66
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7
seq CSLLSGWGQLLRC/VQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 123:

[illegible]

-10

-5

1

5

ACA CAC ATG GGT
Thr His Met Gly

281

(2) INFORMATION FOR SEQ ID NO: 125:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 364 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 284..328
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq WILRILFVIGSXL/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 125:

```

ATGACTGCCA TTTGAAGTGG CTA CTTCAAT GGT TGGTGA TAATAACTTT CAACATTCTG   60
TGAATGTAAG CTGTGCACAC CCTGAATGGC TAGCAGGGCA AAGCATCCTG AATGTGGATC   120
TGANAGATTT WRTCTGTGAT GATTTTCTCA AGCCACAGAT AAGGACACAT CCWGAAACCA   180
TAATTGCTCT AAGAGGCATG ANTGTGACTC TGACGTGCAC TGCAGTGAAG CAGMAGTGAT   240
TCACCCATGT CCACTGTGTG GCGCAAAGAC AGTGAAATCC TGT ATG ACG TGG ATA   295
                               Met Thr Trp Ile
                               -15

CTG AGA ATT TTG TTC GTT ATT GGC AGC ARD CTG GAG AAG CTC TGG AAT   343
Leu Arg Ile Leu Phe Val Ile Gly Ser Xaa Leu Glu Lys Leu Trp Asn
-10                               -5                               1                               5

ATA CTA GTA TCT TAC ATC TTT   364
Ile Leu Val Ser Tyr Ile Phe
                               10

```

(2) INFORMATION FOR SEQ ID NO: 126:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 290 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 123..266
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq ILCIFLGLLIIRC/FK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 126:

```

AGACAAACGT GCCAACACTT AAGTCTACTG GCTGGACTTC ATCTCCATGG CAACAAGCAT    60
GGAAGGCAAA GAGTTGATTC CAGAAGGAAC TGTGAAGAGC CACAACAATG TGCCAGTGAA    120
TA ATG AGT AGT ACC TAC TGT GGC AAC TCT TCA GCT AAG ATG AGT GTC    167
Met Ser Ser Thr Tyr Cys Gly Asn Ser Ser Ala Lys Met Ser Val
      -45                      -40                      -35
AAC GAA GTA TCA GCT TTC TCA TTG AGT CTG GAG CAA AAA ACT GGC TTT    215
Asn Glu Val Ser Ala Phe Ser Leu Ser Leu Glu Gln Lys Thr Gly Phe
      -30                      -25                      -20
GCT TTT GTT GGG ATT TTG TGT ATC TTC TTG GGA CTT CTT ATT ATC CGA    263
Ala Phe Val Gly Ile Leu Cys Ile Phe Leu Gly Leu Leu Ile Ile Arg
      -15                      -10                      -5
TGC TTC AAA ATC CTG CTA GNS CAA TCG                                290
Cys Phe Lys Ile Leu Leu Xaa Gln Ser
      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 127:

-(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 208 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 143..196
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq LTMLSMIVGATCY/AM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 127:

```

AGCTGCAGGA CTTCCCGCGC AACTGCTGGG TGTCCATCAA TGGCATGGTG AACCACTCGT    60
GGAGTTAACT GTACTCCTTC GCACTCTTCA AGGCCATGAG CCACATGCTG TGCATCGGGT    120
ACGGCCGGCA GSGCCCGAGA GC ATG ACG GAC ATC TGG CTG ACC ATG CTC AGC    172
                        Met Thr Asp Ile Trp Leu Thr Met Leu Ser
                        -15                               -10

ATG ATT GTG GGT GCC ACC TGC TAC GCC ATG ATC GGG    208
Met Ile Val Gly Ala Thr Cys Tyr Ala Met Ile Gly
      -5                               1

```

(2) INFORMATION FOR SEQ ID NO: 128:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 214 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 59..109
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq WIYAFISLGYILG/SG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 128:

```

AAGTTTGGGT TGTTTCTGCT TTTCGTTAAA AATAGTAGTA CTTCTCTGAA CACTGTGG    58
ATG AGH NTT TGC TGG ATA TAT GCT TTC ATT TCT CTT GGG TAT ATA CTT    106
Met Xaa Xaa Cys Trp Ile Tyr Ala Phe Ile Ser Leu Gly Tyr Ile Leu
      -15                               -10                               -5

GGG AGT GGA ATT GTT GGG TTA TTT GGT AAT TTT ATG TTT AAA CTT TTG    154
Gly Ser Gly Ile Val Gly Leu Phe Gly Asn Phe Met Phe Lys Leu Leu
      1                               5                               10                               15

AGG AAC TGC CAG ACC GTT TTC CAG GAT GGC TAT GCT ATA TTA CCC TTC    202
Arg Asn Cys Gln Thr Val Phe Gln Asp Gly Tyr Ala Ile Leu Pro Phe
      20                               25                               30

CCA CCA ACG GGG    214
Pro Pro Thr Gly
      35

```

(2) INFORMATION FOR SEQ ID NO: 129:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 156 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 19..111
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq QLALSWVPPXCRV/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 129:

ACTTTTAGGC CCATTGGG ATG TTC ATT AGA ACT CTG AAA ACT ACA GTT CTC	51
Met Phe Ile Arg Thr Leu Lys Thr Thr Val Leu	
-30 -25	
CCC TTT ATG AGG ACT GCA CCA CAG CTC GCC CTC TCC TGG GTT CCG CCT	99
Pro Phe Met Arg Thr Ala Pro Gln Leu Ala Leu Ser Trp Val Pro Pro	
-20 -15 -10 -5	
RRT TGC AGA GTG AGC CCA TGG GAC AGC CCT CTG AAA TTA TAC TGC TTA	147
Xaa Cys Arg Val Ser Pro Trp Asp Ser Pro Leu Lys Leu Tyr Cys Leu	
1 5 10	
CAA CCG CAG	156
Gln Pro Gln	
15	

(2) INFORMATION FOR SEQ ID NO: 130:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 297 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 211..282
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5

seq SVLIFCLLPYIYH/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 130:

ATAAAAATAA TTTATATTTG ATGTTATAAT ACATTATTTA ATTTTATAG TAAATTCAC T 60
 AATCTGTTTT GTTAGACCA GKRTGAGAGA AAAGSSAGAC TKCGAGTTAA TACTTTGTAA 120
 GGTTATCTGC ACTCTCATCT GTGGTTGGCA ATATTGATG CAGTTTATAT TAGATTTATG 180
 TTGTTGTTAT TTAGATAGTT TGGAGCTGAG ATG AGG ACA GGA GCT GAG ATG AGG 234
 Met Arg Thr Gly Ala Glu Met Arg
 -20
 ACA AAC TCT TCA GTT TTA ATT TTT TGT TTG CTC CCA TAT ATT TAT CAT 282
 Thr Asn Ser Ser Val Leu Ile Phe Cys Leu Leu Pro Tyr Ile Tyr His
 -15 -10 -5
 TTT TTT CCA GCT TCG 297
 Phe Phe Pro Ala Ser
 1 5

(2) INFORMATION FOR SEQ ID NO: 131:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 310 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 146..214
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq EGLELGFSHRTFA/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 131:

ATAGGGCAAT TANNGGCTCA TCTCATTTGT TTCCWNWNCT CTCAGGGGCC ACAGCCCTGC 60
 ATTACCTGTT TGCCAATGTG TGAAAGCAAT TGTTTCATGT ATTTTGCCCA GTTTTCTAGT 120
 TTCTTATGGC AGGAAGTTAA GTCCA ATG ATT GTT ATT CCA TCA TGG CTG GAA 172
 Met Ile Val Ile Pro Ser Trp Leu Glu
 -20 -15
 AAC GAG GGT CTG GAA TTG GGA TTT TCA CAT AGG ACC TTT GCT TTT CCT 220
 Asn Glu Gly Leu Glu Leu Gly Phe Ser His Arg Thr Phe Ala Phe Pro
 -10 -5 1

GTG ACA CAT GCT TCC TCT CAG TAC ATA TGG ATG AAC TNK CTG ACT AGG 268
 Val Thr His Ala Ser Ser Gln Tyr Ile Trp Met Asn Xaa Leu Thr Arg
 5 10 15

ACT ACA GTA GCA ATA TCA GTT TAT TTT TGG ACC CAC ACA GGG 310
 Thr Thr Val Ala Ile Ser Val Tyr Phe Trp Thr His Thr Gly
 20 25 30

(2) INFORMATION FOR SEQ ID NO: 132:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 409 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 266..394
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq VISVFLSFLPSYP/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 132:

ACAAATCTAA ATCTAAAGAT AATTTTCAAG TTTGGCCAGG GAATCCTTAT GTTATATTGA 60
 TTCCATAAAA ACAGGGGTAT GTATTACCTA GCAGCTGAAG AAGAGATGGC CTACATTAGG 120
 CACCTGAGTG TCAGAAGGGG TTGTCAAGAA AGGTACAGT TAATACATAA GTCTGTCCTG 180
 ATTGAATTAA TTAATCAAGA ACTACAGGTG TTGAAAAGAG AAATGGGTAC CTGGAGAAGG 240
 GCATCATCTA TATTGGCATC TTGGA ATG TTA AAA AAG GAA ATA GCT CAC CAC 292
 Met Leu Lys Lys Glu Ile Ala His His
 -40 -35
 AGC CCT AGC CTG GTG AGC TGC CCT GTC TGC ACC ACA AAA TAT AGA ACT 340
 Ser Pro Ser Leu Val Ser Cys Pro Val Cys Thr Thr Lys Tyr Arg Thr
 -30 -25 -20
 CTG AGA CTC CTG AGG GTT ATC TCA GTT TTT CTG TCT TTT CTT CCT TCT 388
 Leu Arg Leu Leu Arg Val Ile Ser Val Phe Leu Ser Phe Leu Pro Ser
 -15 -10 -5
 TAC CCA GGG TTC AGC ATG CAA 409
 Tyr Pro Gly Phe Ser Met Gln
 1 5

(2) INFORMATION FOR SEQ ID NO: 133:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 343 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 112..344
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 36..268
id AA256780
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 248..337
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq CAYSLPGVALTLG/RQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 133:

```

TGATCGGACC ATTTCACTGC AGCAAGCAAC ACAGTATTCT GAGCAGAAGA TCGGGACTTG      60
AGGCCATGTT GCGGAGGGCC AGTGACATTA TCTGGACTCT GGAGTGTGRR GRAATATBGA    120
STCCACKCTT CACTATATTC ACAGCGATTC AGACTTGAGC AACAAATAGCA GTTTTAGCCC    180
TGATGAGGAA AGGAGAACTR RAGTACAAGA TGTGTACCT CAGGCCTTGT TAGATCAGKA     240
TTTATCT ATG ACT GRS CCT TCT CGT GCA CAG ACG GTT GAC ASK GGA ATT      289
      Met Thr Xaa Pro Ser Arg Ala Gln Thr Val Asp Xaa Gly Ile
      -30                -25                -20

GCT AAG CAC TGT GCA TAT AGC CTC CCT GGT GTG GCC TTG ACA CTC GGA      337
Ala Lys His Cys Ala Tyr Ser Leu Pro Gly Val Ala Leu Thr Leu Gly
      -15                -10                -5

AGA CAG
Arg Gln
1

```

343

(2) INFORMATION FOR SEQ ID NO: 134:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 151 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 52..133

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 90
region 55..136
id W81722
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 90..150

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 90
region 1..61
id R67182
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 80..114

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 97
region 1..35
id R57498
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 20..50

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 90
region 297..327
id N21080
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 71..133

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 9.8
seq LGLLCALLPQHHG/AP.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 134:

AGCAGCTCCC AGGATGAACT GGTTCAGTG GCTGCTGCTG CTGCGGGGGC GCTGAGAGGA 60

CACGAGCTCT ATG CCT TTC CGG CTG CTC ATC CCG CTC GGC CTC CTG TGC 109
Met Pro Phe Arg Leu Leu Ile Pro Leu Gly Leu Leu Cys
-20 -15 -10

GCG CTG CTG CCT CAG CAC CAT GGT GCG CCA GGT CCC GAY KGG 151

Ala Leu Leu Pro Gln His His Gly Ala Pro Gly Pro Asp Xaa
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 135:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 244 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 68..245
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 1..178
id T08712
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 79..245
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 2..168
id R88049
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 80..154
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 1..75
id AA094697
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 170..202
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 89..121
id AA094697
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 112..245
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 35..168

id H30765
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 79..121
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 1..43
id H30765
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 112..245
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 21..154
id H38484
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 134..187
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.2
seq VFLCSLLAPMVLA/SA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 135:

```

AAGGASGCAG GGGAGGCGGG AAAGCAGCTC AAGCCTCACC CACCGCCCTG CCCCCAGCCC      60
CGCSACTCCC AGGCTCCTCG GGA CTGGCG GGTCTCCTG GGAGTCTCGS AGGGGACCGG      120
CTGTGCAGAC GCC ATG SAG TTG GTG CTG GTC TTC CTC TGC AGC CTG CTG      169
           Met Xaa Leu Val Leu Val Phe Leu Cys Ser Leu Leu
           -15                               -10
GCC CCC ATG GTC CTG GCC AGT GCA GCT GAR AAG GAG RAG GAM ATG GAS      217
Ala Pro Met Val Leu Ala Ser Ala Ala Glu Lys Glu Xaa Xaa Met Xaa
          -5              1              5              10
CCT TTT CAT TAT GAT TAC CAG ACC CTG      244
Pro Phe His Tyr Asp Tyr Gln Thr Leu
           15

```

(2) INFORMATION FOR SEQ ID NO: 136:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 281 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 52..283
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 1..232
 id AA111270
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 15..104
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 9
 seq FFLLLLFRGCLIG/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 136:

AAGCAACCCT CGAC ATG GCG CTG AGG CGG CCA CCG CGA CTC CGG CTC TGC	50
Met Ala Leu Arg Arg Pro Pro Arg Leu Arg Leu Cys	
-30 -25 -20	
GCT CGG CTG CCT GAC TTC TTC CTG CTG CTG CTT TTC AGG GGC TGC CTG	98
Ala Arg Leu Pro Asp Phe Phe Leu Leu Leu Leu Phe Arg Gly Cys Leu	
-15 -10 -5	
ATA GGG GCT GTA AAT CTC AAA TCC AGC AAT CGA ACC CCA GTG GTA CAG	146
Ile Gly Ala Val Asn Leu Lys Ser Ser Asn Arg Thr Pro Val Val Gln	
1 5 10	
GAA TTT GAA AGT GTG GAA CTG TCT TGC ATC ATT ACG GAT TCG CAG ACA	194
Glu Phe Glu Ser Val Glu Leu Ser Cys Ile Ile Thr Asp Ser Gln Thr	
15 20 25 30	
AGT GAC CCC AGG ATC GAG TGG AAG AAA ATT CAA GAT GAA CAA ACC ACA	242
Ser Asp Pro Arg Ile Glu Trp Lys Lys Ile Gln Asp Glu Gln Thr Thr	
35 40 45	
TAT GTG TTT TTT GAC AAC AAA ATT CAG GGA GAC TTG GCG	281
Tyr Val Phe Phe Asp Asn Lys Ile Gln Gly Asp Leu Ala	
50 55	

(2) INFORMATION FOR SEQ ID NO: 137:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 461 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 105..234
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 1..130
 id N77056
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 123..302
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 7.2
 seq VLLTLLLI AFIFL/II

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 137:

```

AAAGTAATCT TTATTCGTC ATTTTGGARA CATAGAAGCC GTAACGGAAG CAAGTGAAAT    60
GCTCAGTCTT AGACGACTGC GTCGTGCTAT GACCGGACTT TTTCTTGAAA GGGGATGACA    120
GC ATG GGA GGC AAT GGC TCC ACA TGT AAA CCC GAC ACT GAA AGA CAA    167
Met Gly Gly Asn Gly Ser Thr Cys Lys Pro Asp Thr Glu Arg Gln
-60                      -55                      -50

GGC ACT CTC TCC ACA GCA GCC CCA ACA ACT AGC CCT GCA CCC TGT CTC    215
Gly Thr Leu Ser Thr Ala Ala Pro Thr Thr Ser Pro Ala Pro Cys Leu
-45                      -40                      -35                      -30

TCT AAC CAC CAC AAC AAA AAA CAT TTA ATC CTT GCC TTT TGT GCT GGG    263
Ser Asn His His Asn Lys Lys His Leu Ile Leu Ala Phe Cys Ala Gly
-25                      -20                      -15

GTT CTA CTG ACA CTG CTG CTG ATA GCC TTT ATC TTC CTC ATC ATA ABB    311
Val Leu Leu Thr Leu Leu Leu Ile Ala Phe Ile Phe Leu Ile Ile Xaa
-10                      -5                      1

AGC TAC AGA AAA TAT CAC TCC AAG CCC CAG GCC CCA GAT CCT CAC TCA    359
Ser Tyr Arg Lys Tyr His Ser Lys Pro Gln Ala Pro Asp Pro His Ser
-5                      10                      15

GAT CCT CCA GCG VNG CTT TCA TBS ATC CCA GGG GAA ATC ACT TAC CTA    407
Asp Pro Pro Ala Xaa Leu Ser Xaa Ile Pro Gly Glu Ile Thr Tyr Leu
20                      25                      30                      35

TGC CAG CAC AAC TTT CAA ACT CTY NRN WBW AAA AGA GCA ATC ACT TGG    455
Cys Gln His Asn Phe Gln Thr Leu Xaa Xaa Lys Arg Ala Ile Thr Trp
40                      45                      50

CTG AGA    461
Leu Arg

```

(2) INFORMATION FOR SEQ ID NO: 138:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 380 base pairs
 (B) TYPE: NUCLEIC ACID

- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 233..381
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 157..305
id N78069
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 72..129
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 1..58
id N78069
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 117..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 45..108
id N78069
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 173..224
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 100..151
id N78069
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 233..381
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 183..331
id AA022144
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 117..234
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 69..186
id AA022144

est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 89..129
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 42..82
id AA022144
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 242..369
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 170..297
id AA043627
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 89..129
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 21..61
id AA043627
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 117..233
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 194..310
id W68800
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 251..304
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 329..382
id W68800
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 89..129
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 167..207
id W68800
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 117..224
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94

region 70..177
id H91637
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 251..337
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 202..288
id H91637
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 89..129
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 43..83
id H91637
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 90..206
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 7.2
seq SALAKLLLLTCCSA/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 138:

```

AAACTCTCAA CCCACTTCTC CAGCCAGCGC CCCAGCCCTC CCGCCGCCCG CTCGCAGGTC      60
CCGAGGAGCG CAGACTGTGK CCCTGGCAA ATG GGA ACA GCH GAC AGT GAT GAG      113
                               Met Gly Thr Ala Asp Ser Asp Glu
                               -35
ATG GCC CCG GAG GCC CCA CAG CAC ACC CAC ATC GAT GTG CAC ATC CAC      161
Met Ala Pro Glu Ala Pro Gln His Thr His Ile Asp Val His Ile His
-30                -25                -20
CAG GAG TCT GCC CTG GCC AAG CTC CTG CTC ACC TGC TGC TCT GCG CTG      209
Gln Glu Ser Ala Leu Ala Lys Leu Leu Leu Thr Cys Cys Ser Ala Leu
-15                -10                -5                1
CGG CCC CGG GCC ACC CAG GCC AGG GGC AGC AGC CGG CTG CTG GTG GCC      257
Arg Pro Arg Ala Thr Gln Ala Arg Gly Ser Ser Arg Leu Leu Val Ala
          5                10                15
TCG TGG GTG ATG CAG ATC GTG CTG GGG ATC DTG AGT GCA GTC MTA GGA      305
Ser Trp Val Met Gln Ile Val Leu Gly Ile Xaa Ser Ala Val Xaa Gly
          20                25                30
GGA TTT TTC TAC ATC CGC GAC TAS ACC CTC MTC GTC ACC TCG GGA GCW      353
Gly Phe Phe Tyr Ile Arg Asp Xaa Thr Leu Xaa Val Thr Ser Gly Ala
          35                40                45
GCC ATC TGG ACA GGG GCT GTG GCT GTG      380
Ala Ile Trp Thr Gly Ala Val Ala Val
          50                55

```


(2) INFORMATION FOR SEQ ID NO: 139:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 237 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 201..234
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..34
id R74380
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 175..216
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.6
seq SLLSFLFARVNLG/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 139:

```

ACTATCCTAT CTTCTTTAAG GAAAACTTAC CAAGTATGAT TTGGTCCAAG GATTTCAGGT    60
GGGCTCTCAG TGCTGCTCCC AATATATTAG AGGTCTCCTT CCTTACTAT TTCCTAACCA    120
GATGTAAAT  TAGCTTTTCC CCCCTTCTAC ATCACCTAAC CCGTTTCTCA TTGG  ATG    177
                                   Met
TCC TTG CTT TCG TTT TTA TTT GCC AGA GTA AAT CTA GGA TCT CCC TTG    225
Ser Leu Leu Ser Phe Leu Phe Ala Arg Val Asn Leu Gly Ser Pro Leu
      -10                -5                1

TCT GCC AAC GGG                                237
Ser Ala Asn Gly
      5

```

(2) INFORMATION FOR SEQ ID NO: 140:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 375 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 90..376
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 5..291
id R59435
est .

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 206..376
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..171
id R19220
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 199..371
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 1..173
id HSC32F041
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 283..371
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 1..89
id HSC2VA021
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 136..264
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq WWCCPARLTLTSG/WP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 140:

```
ACTCTTTTGG AAGGTCTCCT TTGCCAGCGC ACACGGCTCC CTGGGCTGGA ATGTCTGTTC    60
ATTCATCCCT GCAGTTGTTT GCGGATGTCC CGGGGGHWAA CGTGAGTYAG TTAATGAAGT    120
CCAAAGCCAA GCCCA ATG GCA AGA AGC CCG CTG CGG AGG AGA GGA AGG CCT    171
      Met Ala Arg Ser Pro Leu Arg Arg Arg Gly Arg Pro
                -40                      -35
```

```

ACC TGG AGC CTG AGC ACA CCA AGG CCA GGA TCA CCG ACT TCC AGT TCA    219
Thr Trp Ser Leu Ser Thr Pro Arg Pro Gly Ser Pro Thr Ser Ser Ser
  -30                      -25                      -20

AGG AGC TGG TGG TGC TGC CCC GCG AGA TTG ACC TTA ACG AGT GGC TGG    267
Arg Ser Trp Trp Cys Cys Pro Ala Arg Leu Thr Leu Thr Ser Gly Trp
-15                      -10                      -5                      1

CCA GCA ACA CCA CGA CGT TTT TCC ACC ACA TCA ACC TGC AGT ATA GCA    315
Pro Ala Thr Pro Arg Arg Phe Ser Thr Thr Ser Thr Cys Ser Ile Ala
          5                      10                      15

CCA TCT CGG AGT TCT GCA CAG GAG AGA CGT GTC AGA CGA TGG CCG TGT    363
Pro Ser Arg Ser Ser Ala Gln Glu Arg Arg Val Arg Arg Trp Pro Cys
          20                      25                      30

GCA ACA CAC AGT                                                    375
Ala Thr His Ser
    35

```

(2) INFORMATION FOR SEQ ID NO: 141:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 268 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 40..264
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..225
id T05872
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 64..264
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 2..202
id T34681
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..264
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..180
id T05903

est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 157..228
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 25..96
id N56217
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 223..264
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 92..133
id N56217
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 158..214
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7
seq TLLLACHLQLEVG/VV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 141:

```

AATCATTTTG AAGACAGAGG AGCAGCTGTT AAGGCTATGG ATGAAATGAA TGGCAAAGAA   60
ATAGAAGGGG AAGAAATTGA AATAGTCTTA GCCAAGCCAC CAGACAAGAA AAGGAAAGAG   120
CGCCAAGCTG CTAGACAGGC CTCCAGAAGC ACTGCGT ATG AAG ATT ATT ACT ACC   175
                               Met Lys Ile Ile Thr Thr
                               -15
ACC CTC CTC CTC GCA TGC CAC CTC CAA TTA GAG GTC GGG GTC GTG GTG   223
Thr Leu Leu Leu Ala Cys His Leu Gln Leu Glu Val Gly Val Val Val
      -10                      -5                      1
GGG GGA GAG GTG GAT ATG GCT ACC CTC CAG ATT ACT ACG GCA TCG   268
Gly Gly Glu Val Asp Met Ala Thr Leu Gln Ile Thr Thr Ala Ser
      5                      10                      15

```

(2) INFORMATION FOR SEQ ID NO: 142:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 195 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 118..191
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 112..185
id R60698
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 122..191
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 1..70
id H17558
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 113..191
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 229..307
id N26943
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 118..191
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 145..218
id W24886
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 76..186
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.7
seq FLGVLALLGYLAV/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 142:

```
AATAAGTTCT CGCGAGACGC AKAAGGCACC GGCTCGAACT GGGGCGGGCC ACTGCCAGGA    60
AAGCAACGCC CCTGA ATG CTT ATG CCG GTG GTT GGT AGA GGA AAT GGA ATT    111
      Met Leu Met Pro Val Val Gly Arg Gly Asn Gly Ile
      -35                               -30

CCC CAG ACT GTT TCA GAA TGG CTT CGG TTA TTG CCT TTC CTT GGT GTA    159
Pro Gln Thr Val Ser Glu Trp Leu Arg Leu Leu Pro Phe Leu Gly Val
-25          -20          -15          -10

CTC GCA CTT CTT GGC TAC CTT GCA GTT CGT CCC GGG    195
Leu Ala Leu Leu Gly Tyr Leu Ala Val Arg Pro Gly
      -5                               1
```

(2) INFORMATION FOR SEQ ID NO: 143:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 386 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 114..343
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 76..305
id T91418
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 41..118
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 2..79
id T91418
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 150..296
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq PPFFLCLQCFTRG/FE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 143:

```

AAGGCGCGCG CGACCGGCGG CTCTTTGGCG CGGATTAGGG GGTCTCGGCG AGGGAGTCAT   60
CAAGCTTTGG TGTATGTGTT GGCCGGTTCT GAAGTCTTGA AGAAGCTCTG CTGAGGAAGA   120
CCAAAGCAGC ACTCGTTGCC AATTAGGGA ATG GAC CGT TTG GGT TCC TTT AGC   173
                               Met Asp Arg Leu Gly Ser Phe Ser
                               -45
AAT GAT CCC TCT GAT AAG CCA CCT TGC CGA GGC TGC TCC TCC TAC CTC   221
Asn Asp Pro Ser Asp Lys Pro Pro Cys Arg Gly Cys Ser Ser Tyr Leu
-40                               -35                               -30
ATG GAG CCT TAT ATC AAG TGT GCT GAA TGT GGG CCA CCT CCT TTT TTC   269
Met Glu Pro Tyr Ile Lys Cys Ala Glu Cys Gly Pro Pro Pro Phe Phe
-25                               -20                               -15                               -10
CTC TGC TTG CAG TGT TTC ACT CGA GGC TTT GAG TAC AAG AAA CAT CAA   317

```

Leu Cys Leu Gln Cys Phe Thr Arg Gly Phe Glu Tyr Lys Lys His Gln
 -5 1 5

AGC GAT CAT ACT TAT GAA ATA ATG GCA GGA TGT AGC CAA TCA AAT GTG 365
 Ser Asp His Thr Tyr Glu Ile Met Ala Gly Cys Ser Gln Ser Asn Val
 10 15 20

CAC CAA GAC CAA GGA GGT CAG 386
 His Gln Asp Gln Gly Gly Gln
 25 30

(2) INFORMATION FOR SEQ ID NO: 144:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 389 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 101..384
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
 region 1..284
 id R35114
 est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 90..365
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
 region 196..471
 id W65227
 est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 45..96
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
 region 152..203
 id W65227
 est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 90..349
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
 region 173..432
 id AA116536

est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 90..384
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 3..297
 id AA065524
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 154..384
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 98
 region 1..231
 id R52412
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 57..317
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 3.6
 seq GLLLLYMVYLTIV/EP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 144:

```

AAGAAAGCCG ACGGGGTGCT GTGGTCAGGA GGC GCGTGGG CGGCAGAATT TCGAGG ATG   59
                                     Met
TCA GAT GTA AAT GTA TCT GCC CTC CCT ATA AAG AAA AAT TCT GGG CAT   107
Ser Asp Val Asn Val Ser Ala Leu Pro Ile Lys Lys Asn Ser Gly His
-85                               -80                               -75
ATT TAT AAT AAG AAC ATA TCT CAG AAA GAT TGT GAT TGC CTT CAT GTC   155
Ile Tyr Asn Lys Asn Ile Ser Gln Lys Asp Cys Asp Cys Leu His Val
-70                               -65                               -60                               -55
GTG GAG -CCC ATG CCT GTG CGG GGG CCT GAT GTA GAA GCA TAC TGT CTA   203
Val Glu Pro Met Pro Val Arg Gly Pro Asp Val Glu Ala Tyr Cys Leu
-50                               -45                               -40
CGC TGT GAA TGC AAA TAT GAA GAA AGA AGC TCT GTC ACA ATC AAG GTT   251
Arg Cys Glu Cys Lys Tyr Glu Glu Arg Ser Ser Val Thr Ile Lys Val
-35                               -30                               -25
ACC ATT ATA ATT TAT CTC TCC ATT TTG GGC CTT CTA CTT CTG TAC ATG   299
Thr Ile Ile Ile Tyr Leu Ser Ile Leu Gly Leu Leu Leu Leu Tyr Met
-20                               -15                               -10
GTA TAT CTT ACT CTG GTT GAG CCC ATA CTG VAG AGG CGC CTC TTT GGA   347
Val Tyr Leu Thr Leu Val Glu Pro Ile Leu Xaa Arg Arg Leu Phe Gly
-5                               1                               5                               10
CAT GCA CAG TTG ATA CAG AGT GAT GAT RAT ATT GGG GGA TTG   389
His Ala Gln Leu Ile Gln Ser Asp Asp Xaa Ile Gly Gly Leu
15                               20

```


(2) INFORMATION FOR SEQ ID NO: 145:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 45..150
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 25..130
id H30752
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 150..222
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 129..201
id H30752
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 93..222
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 76..205
id AA072341
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 73..156
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 1..84
id T31153
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 150..222
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 77..149
id T31153
est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 79..150
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 98
 region 87..158
 id W74452
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 152..222
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 100
 region 158..228
 id W74452
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 79..150
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 98
 region 81..152
 id H26405
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 150..222
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 151..223
 id H26405
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 85..222
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 3.5
 seq ALSLSLSMAPPNP/GP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 145:

```

ACACCTCCCC GCCTTGTTGT CCAACTTCTC CCGGAGCAGC CGGAGAGCAG GCGTCGGGAC   60
GCAGCAAAGA GAGGAGAGGC CACC ATG GCG GAS TGC AGG AGG TGC AGA TCA   111
               Met Ala Xaa Cys Arg Arg Cys Arg Ser
               -45                               -40

CAG AGG AGA AGC CAC TGT TGC CAG GAC AGA CGC CTG AGG CGG CCA AGA   159
Gln Arg Arg Ser His Cys Cys Gln Asp Arg Arg Leu Arg Arg Pro Arg
   -35                               -30                               -25

CTC ACT CTG TGG AGA CAC CAT ACG GCT CTG TCA CTT TCA CTG TCT ATG   207
Leu Thr Leu Trp Arg His His Thr Ala Leu Ser Leu Ser Leu Ser Met
   -20                               -15                               -10

GCA CCC CCA AAC CCA GGC CCG   228

```

Ala Pro Pro Asn Pro Gly Pro
-5 1

(2) INFORMATION FOR SEQ ID NO: 146:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 376 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 246..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 9..138
id N56074
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 89..196
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq LALTALSVXRKXS/XX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 146:

```

ACTGTCTGGC AGCCTGCAAG TCTCCATTCA GAAGCGGCTC CGTGCTGCCC AGCGATGGCG      60
CCCTGGCGGC GCGGAAGCCC GCGGCCAA ATG ACA CGA CTT GGG GGC AAA GGA      112
                               Met Thr Arg Leu Gly Gly Lys Gly
                               -35                      -30

GGA CAA CAG TTC CCA CCA GGA CAA AAA ATA ATA TCC AAA GAT ATT TTG      160
Gly Gln Gln Phe Pro Pro Gly Gln Lys Ile Ile Ser Lys Asp Ile Leu
-25                      -20                      -15

GCA CTA ACG GCG CTA TCT GTA GSA AGR AAG TTK AGC ART GKG RAM YKG      208
Ala Leu Thr Ala Leu Ser Val Xaa Arg Lys Xaa Ser Xaa Xaa Xaa Xaa
-10                      -5                      1

TKN RAG ACT TCC AAG GAG ACT TBA GAC AAC CAA GAC AGT GTA AAG GAA      256
Xaa Xaa Thr Ser Lys Glu Thr Xaa Asp Asn Gln Asp Ser Val Lys Glu
5                      10                      15                      20

AAC AGA GAA AAA GAC TTG TTA GAC ATT ATT AAG GGC ACG AAA GTT GAA      304
Asn Arg Glu Lys Asp Leu Leu Asp Ile Ile Lys Gly Thr Lys Val Glu
25                      30                      35

TTG AGC ACA GTA AAT GTA CAA ACA ACA AAG CCA CCC AAC AGA AGT TCA      352

```

Leu Ser Thr Val Asn Val Gln Thr Thr Lys Pro Pro Asn Arg Ser Ser
40 45 50

CTT AAA AGC TAC AAC TGG CGG GCG
Leu Lys Ser Tyr Asn Trp Arg Ala
55 60

376

(2) INFORMATION FOR SEQ ID NO: 147:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 265 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 123..265.
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 187..329
id N57089
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..127
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 110..190
id N57089
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..218
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 2..217
id AA136163
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 154..266
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 52..164
id R22491
est

(ix) FEATURE:

- (A) NAME/KEY: other

(B) LOCATION: 104..160
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 3..59
id R22491
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 47..150
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 1..104
id R29291
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 194..253
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 12.4
seq ALLLGALLGTAWA/RR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 147:

```
AAGGCGGTCTG CCGGGACACC CCGTGTGTGG CAGGCGGCGA ASGCTCTGGA GAATCCCGGA    60
CAGCCCTGCT CCCTGCAGCC AGGTGTAGTT TCGGGAGCCA CTGGGGCCAA AGTGAGAGTC   120
CAGCGGTCTT CCACGCCTTG GGCCACGGCG GCGGCCCTGG GAGCAGAGGT GGAGCGACCC   180
CATTACGCTA AAG ATG AAA GGC TGG GGT TGG CTG GCC CTG CTT CTG GGG       229
      Met Lys Gly Trp Gly Trp Leu Ala Leu Leu Gly
      -20          -15          -10

GCC CTG CTG GGA ACC GCC TGG GCT CGG AGG AGC CGG                265
Ala Leu Leu Gly Thr Ala Trp Ala Arg Arg Ser Arg
      -5          1
```

(2) INFORMATION FOR SEQ ID NO: 148:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 222 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 134..220
(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 91
region 14..100
id AA001733
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 88..135
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 11.4
seq LLCLLLLFGGGDP/RR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 148:

```

AAAGGTGCGC GTGCTCGCTG GTTCTAACCC TTCTGTTGGG CGTTTCTGCG GAGAGGCGGG      60
AGGCGCTGAG AGTCTGTGCG GAGGTCC ATG CAC AGA CTG CTT TGC CTG TTG TTG      114
                               Met His Arg Leu Leu Cys Leu Leu Leu
                               -15                               -10

CTC TTC GGA GGC GGC GAT CCC CGA AGG CGA GCT GAA ATA CGG CTG CAG      162
Leu Phe Gly Gly Asp Pro Arg Arg Arg Ala Glu Ile Arg Leu Gln
      -5              1              5

GCT ACA ATT TGC AGC CGA CCA TTA AGG AAG ACG ACG AGC GGG AGA GGT      210
Ala Thr Ile Cys Ser Arg Pro Leu Arg Lys Thr Thr Ser Gly Arg Gly
      10              15              20              25

GGC CCA CCC TGG
Gly Pro Pro Trp
                                         222

```

(2) INFORMATION FOR SEQ ID NO: 149:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 472 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 245..466
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 74..295
id R61190
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 181..258
(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 94
region 11..88
id R61190
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 89..154
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 11.1
seq QLLALFFLPFCLC/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 149:

```

AGCTCCAGTC CTGGCATCTG CCCGAGGAGA CCACGCTCCT GGAGCTCTGC TGTCTTCTCA    60
GGGAGACTCT GAGGCTCTGT TGAGAATC ATG CTT TGG AGG CAG CTC ATC TAT    112
                               Met Leu Trp Arg Gln Leu Ile Tyr
                               -20                               -15

TGG CAA CTG CTG GCT TTG TTT TTC CTC CCT TTT TGC CTG TGT CAA GAT    160
Trp Gln Leu Leu Ala Leu Phe Phe Leu Pro Phe Cys Leu Cys Gln Asp
                               -10                               -5                               1

GAA TAC ATG GAG TCT CCA CAA ACC GGA GGA CTA CCC CCA GAC TGC AGT    208
Glu Tyr Met Glu Ser Pro Gln Thr Gly Gly Leu Pro Pro Asp Cys Ser
                               5                               10                               15

AAG TGT TGT CAT GGA GAC TAC AGC TTT CGA GGC TAC CAA GGC CCC CCT    256
Lys Cys Cys His Gly Asp Tyr Ser Phe Arg Gly Tyr Gln Gly Pro Pro
                               20                               25                               30

GGG CCA CCG GGC CCT CCT GGC ATT CCA GGA AAC CAT GGA AAC AAT GGC    304
Gly Pro Pro Gly Pro Pro Gly Ile Pro Gly Asn His Gly Asn Asn Gly
                               35                               40                               45                               50

AAC AAT GGA GCC ACT GGT CAT GAA GGA GCC AAA GGT GAG AAG GGC GAC    352
Asn Asn Gly Ala Thr Gly His Glu Gly Ala Lys Gly Glu Lys Gly Asp
                               55                               60                               65

AAA GGT GAC CTG GGG CCT CGA GGG GAG CGG GGG CAG CAT GGC CCC AAA    400
Lys Gly Asp Leu Gly Pro Arg Gly Glu Arg Gly Gln His Gly Pro Lys
                               70                               75                               80

GGA GAG AAG GGC TAC CCG GGG ATT CCA CCA GAA CTT CAG ATT GCA TTC    448
Gly Glu Lys Gly Tyr Pro Gly Ile Pro Pro Glu Leu Gln Ile Ala Phe
                               85                               90                               95

ATG GCT TCT CTG GMA CCC ACT TCA    472
Met Ala Ser Leu Xaa Pro Thr Ser
                               100                               105

```

(2) INFORMATION FOR SEQ ID NO: 150:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 191 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 37..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..157
id T30099
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 53..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..141
id T35974
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 58..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..136
id T35248
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 58..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..136
id T32601
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 59..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..135
id T31945
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 81..131
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 9.2
seq LLLVAASAMVRS/XA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 150:

```

AAGTGGGGAG CAGCTCGCTC CTGGGCTTTG GGCTGGCTGC AGTCTGTCTG AGGGCGGCCG      60
AAGTGGCTGG CTCATTKAAG ATG AGG CTT CTA CTG CTT CTC CTA GTG GCG GCG      113
                Met Arg Leu Leu Leu Leu Leu Val Ala Ala
                  -15                      -10

TCT GCG ATG GTC CGG AGC GAK GCC TCG GCC AAT CTG GGC GGC GTG CCC      161
Ser Ala Met Val Arg Ser Xaa Ala Ser Ala Asn Leu Gly Gly Val Pro
   -5                1                5                10

AGC AAG AGA TTA AAG ATG CAG TAC ACC ACG      191
Ser Lys Arg Leu Lys Met Gln Tyr Thr Thr
          15                20

```

(2) INFORMATION FOR SEQ ID NO: 151:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 561 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 136..210
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92
region 177..251
id H23535
est

{ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 166..240
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9
seq ALLVLLGVAASLC/VR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 151:

```

AGGATGTACG GATGATTCAG TGGCTGGCAG GAAGCCCGCC CTGCCCGCCC GCCAGTGTCA      60
GTGGTGTGG CATCAGCTTG GGCAGGTGTG CGGGCTCAGG ATGGGGCGGC CGTGGTGAGG      120
AACCCTGGAC TCTCAGCATC ACAAGAGGCA ACACCAGGAG CCAAC ATG AGC TCG GGC      177
                Met Ser Ser Gly
                  -25

RCT GAA CTG CTG TGG CCC GGA GCA GCG CTG CTG GTG CTG TTG GGG GTG      225
Xaa Glu Leu Leu Trp Pro Gly Ala Ala Leu Leu Val Leu Leu Gly Val

```

-20	-15	-10	
GCA GCC AGT CTG TGT GTG CGC TGC TCA CGC CCA GGT GCA AAG AGG TCA			273
Ala Ala Ser Leu Cys Val Arg Cys Ser Arg Pro Gly Ala Lys Arg Ser			
-5	1	5	10
GAG AAA ATC TAC CAG CAG AGA AGT CTG CGT GAG GAC CAA CAG AGC TTT			321
Glu Lys Ile Tyr Gln Gln Arg Ser Leu Arg Glu Asp Gln Gln Ser Phe			
15	20	25	
ACG GGG TCC CGG ACC TAC TCC TTG GTC GGG CAG GCA TGG CCA GGA CCC			369
Thr Gly Ser Arg Thr Tyr Ser Leu Val Gly Gln Ala Trp Pro Gly Pro			
30	35	40	
CTG GCG GAC ATG GCA CCC ACA AGG AAG GAC AAG CTG TTG CAA TTC YAC			417
Leu Ala Asp Met Ala Pro Thr Arg Lys Asp Lys Leu Leu Gln Phe Xaa			
45	50	55	
CCC AGC CTG GAG GMT CCA AGC ATC TTC CAG GKR MCA GAA MTT CAG CCA			465
Pro Ser Leu Glu Xaa Pro Ser Ile Phe Gln Xaa Xaa Glu Xaa Gln Pro			
60	65	70	75
GTC TGT GTG TGC GCT GCT CAC GCC CAG GTG CAA ANN NVT CAG AGA AAA			513
Val Cys Val Cys Ala Ala His Ala Gln Val Gln Xaa Xaa Gln Arg Lys			
80	85	90	
TCT ACC AGC AGA GAA GTC TGC GTG AGG ACC AAC AGA GCT TTA CGG GGC			561
Ser Thr Ser Arg Glu Val Cys Val Arg Thr Asn Arg Ala Leu Arg Gly			
95	100	105	

(2) INFORMATION FOR SEQ ID NO: 152:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 375 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 128..305
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 106..283
id H14437
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 24..116
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100

region 2..94
id H14437
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 317..376
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 36..95
id HSA86D041
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 1..234
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 8.8
seq SCLGLTLMPFASS/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 152:

ATG ACT AAG GAG ATT TTT TTT TTC ACA GTT GAG TTA GTT TGT GAA AAT	48
Met Thr Lys Glu Ile Phe Phe Phe Thr Val Glu Leu Val Cys Glu Asn	
-75 -70 -65	
AAA GAA CTC TGT AGC TCA CCA AGG TGG AGA AAC GCA ATT CAG AAA AGT	96
Lys Glu Leu Cys Ser Ser Pro Arg Trp Arg Asn Ala Ile Gln Lys Ser	
-60 -55 -50	
AAT TTC TCC AAG GTC ACT TCT TTT TTT ATG TCT TGC CAT CAC TTT AAA	144
Asn Phe Ser Lys Val Thr Ser Phe Phe Met Ser Cys His His Phe Lys	
-45 -40 -35	
GGA CTA GCC CCA CTC CCC CAT GTG TAT ACA CAA GGA AAT TGC AGA CCA	192
Gly Leu Ala Pro Leu Pro His Val Tyr Thr Gln Gly Asn Cys Arg Pro	
-30 -25 -20 -15	
ATT AGT TGT CTT GGC CTG ACT CTA ATG CCT TTT GCA AGT AGC TTT CCA	240
Ile Ser Cys Leu Gly Leu Thr Leu Met Pro Phe Ala Ser Ser Phe Pro	
-10 -5 1	
GAA GTA AAA GTC CCA GTG ATG TAT TCC CAT AGA AAT ATT TTT CAG TTG	288
Glu Val Lys Val Pro Val Met Tyr Ser His Arg Asn Ile Phe Gln Leu	
5 10 15	
TTT ATG TCG TTT ACT ACA AAA AAA AAG ATT CAG AGT GGA TGG AGT ACA	336
Phe Met Ser Phe Thr Thr Lys Lys Lys Ile Gln Ser Gly Trp Ser Thr	
20 25 30	
ACT CTG AGT ATT TTT CTA GTC CGG AAT TTT TTA TTA ATA	375
Thr Leu Ser Ile Phe Leu Val Arg Asn Phe Leu Leu Ile	
35 40 45	

(2) INFORMATION FOR SEQ ID NO: 153:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 339 base pairs

- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 209..332
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 39..162
id N42351
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 209..324
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 39..154
id N42357
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 209..284
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 25..100
id T57420
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 253..315
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5
seq YFRALCLPRGAWG/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 153:

```
AAGGTGAGCT GAGGTACCTG GCTTTAGTGA AGACCCTGTG GGGGCTTCCT GGCTGCGGCT    60
TCGACRWGCC TGASTCCASA CGCCCCTAAG GTTTGATGAA AACCTGCTGG AGGTTTGGGA    120
CCTAACCGCA TCAAAGTCGC CTTTAGCGGT GCCTGGACCC AGTTCGCACG GGAGGAAGTA    180
GGAGGCAGAA TCCCCTTTGG GCCACAGAAA GCTCACCTGT TACTCGGCCT CCCAGAAAGA    240
TGGATAGGAG AA ATG ACT ACG GAT ATA GGG TGC CTC TAT TTC AGG GCC CTC    291
      Met Thr Thr Asp Ile Gly Cys Leu Tyr Phe Arg Ala Leu
      -20                      -15                      -10

TGC CTC CCC CGG GGA GCC TGG GGC TTC CCT TCC CTC CAG ATT AAG GGG    339
Cys Leu Pro Arg Gly Ala Trp Gly Phe Pro Ser Leu Gln Ile Lys Gly
```

-5

1

5

(2) INFORMATION FOR SEQ ID NO: 154:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 10..109
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 1..100
id W30713
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 24..109
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..86
id HUM402E06B
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 44..109
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 30..95
id H50196
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 38..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.4
seq LTCLFLFLNLRWS/RH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 154:

AGTGTCTGCA CTTGGGCTGC TCTCGGGTTA GCACCCT ATG GTG CCT TCT CTT GTG 55
Met Val Pro Ser Leu Val
-20

ATC CCT GAC CTA ACC TGT CTC TTC CTT TTC CTC AAC CTC AGG TGG AGC 103

Ile Pro Asp Leu Thr Cys Leu Phe Leu Phe Leu Asn Leu Arg Trp Ser
-15 -10 -5

CGC CAC GTA

112

Arg His Val

1

(2) INFORMATION FOR SEQ ID NO: 155:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 191 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 89..189
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 71..171
id R11825
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 17..84
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 1..68
id R11825
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 98..189
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 83..174
id H08475
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 13..84
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 1..72
id H08475
est

(ix) FEATURE:

- (A) NAME/KEY: other

(B) LOCATION: 30..84
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 49..103
id AA113990
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 150..189
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 163..202
id AA113990
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 89..141
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 67..119
id C14102
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 46..89
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 25..68
id C14102
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 160..189
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 93
region 141..170
id C14102
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 24..68
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 7
seq LRLKLAATSASA/RV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 155:

GATCCTGAGC TGACCGGGTA GCC ATG GCC TTG CGG CTC CTG AAG CTG GCA GCG 53
Met Ala Leu Arg Leu Lys Leu Ala Ala
-15 -10

ACG TCC GCG TCC GCC CGG GTC GTG RMG GCG GRM GCC CAG CGC GTG AGA 101
Thr Ser Ala Ser Ala Arg Val Val Xaa Ala Xaa Ala Gln Arg Val Arg
-5 1 5 10

GGA ATT CAT AGC AGT GTG CAG TGC AAG CTG CGC TAT GGA ATG TGG CAT	149
Gly Ile His Ser Ser Val Gln Cys Lys Leu Arg Tyr Gly Met Trp His	
15 20 25	
TTC CTA CTT GGG GAT AAA GCA AGC AAA AGA CTG ACA GTA CAG	191
Phe Leu Leu Gly Asp Lys Ala Ser Lys Arg Leu Thr Val Gln	
30 35 40	

(2) INFORMATION FOR SEQ ID NO: 156:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 160 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 24..161
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 1..138
id AA046377
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 75..161
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..87
id AA112337
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 33..62
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 333..362
id R72972
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 29..88
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9
seq VLLFLYSVLLTKG/IE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 156:

AAAAGATGCT GTCTTGGACC AGTATTCA ATG TGG GGA AAT AAA TTT GGA GTA	52
Met Trp Gly Asn Lys Phe Gly Val	
-20 -15	
TTG CTT TTT CTG TAT TCT GTA TTA CTG ACA AAG GGC ATT GAA AAC ATA	100
Leu Leu Phe Leu Tyr Ser Val Leu Leu Thr Lys Gly Ile Glu Asn Ile	
-10 -5 1	
AAA AAC GAA ATT GAA GAT GCA AGT GAA CCC TTG ATA GAT CCT GTA TAT	148
Lys Asn Glu Ile Glu Asp Ala Ser Glu Pro Leu Ile Asp Pro Val Tyr	
5 10 15 20	
GGA CAT GGC ARC	160
Gly His Gly Xaa	

(2) INFORMATION FOR SEQ ID NO: 157:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 410 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(2..353)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..352
id N43180
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(2..332)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..331
id W56791
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 70..396
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 1..327
id T32797
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(123..396)

(C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 62..335
 id T08229
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: complement(109..396)
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 62..349
 id HSC27A061
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 279..359
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 6.8
 seq VFVCSSVLGQSWG/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 157:

```

ATGTAGGACT TGTTCTGTG GGCTTCAGTG ATGGGATAGT ACACTTCACT CAGAGGCATT   60
TGCATCTTTA AATAATTTCT TAAAAGCCTC TAAAGTGATC AGTGCCTTGA TGCCAACTAA  120
GGAAATTTGT TTAGCATTGA ATCTCTGAAG GCTCTATGAA AGGAATAGCA TGATGTGCTG  180
TTAGAATCAG ATGTTACTGC TAAATTTTAC ATGTTGTGAT GTAAATTGTG TAGAAAACCA  240
TTAAATCATT CAAAATAATA AACTATTTTT ATTAGAGA ATG TAT ACT TTT AGA AAG   296
                               Met Tyr Thr Phe Arg Lys
                               -25

CTG TCT CCT TAT TTA AAT AAA ATA GTG TTT GTC TGT AGT TCA GTG TTG   344
Leu Ser Pro Tyr Leu Asn Lys Ile Val Phe Val Cys Ser Ser Val Leu
-20                               -15                               -10

GGG CAA TCT TGG GGG GGA TTC TTC TCT AAT CTT TCA GAA ACT TTG TCT   392
Gly Gln Ser Trp Gly Gly Phe Phe Ser Asn Leu Ser Glu Thr Leu Ser
-5                               1                               5                               10

GCG ACA CTC TTT AAT GGA                                           410
Ala Thr Leu Phe Asn Gly
15

```

(2) INFORMATION FOR SEQ ID NO: 158:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 188 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 3..186
id T31677
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 3..186
id HUM76142
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 39..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 1..148
id M85529
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 120..182
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4
seq TFCLIFGLGAVWG/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 158:

```

AAAAATCCGA GGCAGCAGCA GGAGAGACAA ACGTTATTTT CCCGCTTGAT TCCAAGAACC   60
TCTTCGATAT TTATTTTAT TTTTAAAGAG GGAGACGATG GACTGAGCTG ATCCGCACC   119
ATG GAG TCT CGG GTC TTA CTG AGA ACA TTC TGT TTG ATC TTC GGT CTC   167
Met Glu Ser Arg Val Leu Leu Arg Thr Phe Cys Leu Ile Phe Gly Leu
  -20                      -15                      -10

GGA GCA GTT TGG GGG CTG GTG                               188
Gly Ala Val Trp Gly Leu Val
  -5                      1

```

(2) INFORMATION FOR SEQ ID NO: 159:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 218 base pairs
- (B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(37..103)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 2..68
id HSC07B041
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(2..38)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 68..104
id HSC07B041
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(1..38)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 60..97
id W23273
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(2..38)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 60..96
id W22817
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(2..37)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 91
region 54..89
id W23007
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 33..143
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.4
seq ILIFLGFFLGFLFH/QF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

```

ACGGCCCGGC AGGAACAACT CATTCCGCTT AT ATG CTT GTT CTT AAA AAG CAC      53
                               Met Leu Val Leu Lys Lys His
                               -35

TCG GTA AAC ATT GCG GCC CAG ACG TGT TTT AAA TTC AAT TTT ATT TTC      101
Ser Val Asn Ile Ala Ala Gln Thr Cys Phe Lys Phe Asn Phe Ile Phe
-30                -25                -20                -15

AGG ATC CTC ATC TTT CTT GGT TTC TTT CTG GGG CTT TTC CAT CAG TTC      149
Arg Ile Leu Ile Phe Leu Gly Phe Phe Leu Gly Leu Phe His Gln Phe
                -10                -5                1

CTC TTC CTC TTT CTC TTT GCT GGC AAT CTC AGC TCG TAC CTT TTG AAG      197
Leu Phe Leu Phe Leu Phe Ala Gly Asn Leu Ser Ser Tyr Leu Leu Lys
                5                10                15

CAG AGC AAA ATC CAA GCC AGG      218
Gln Ser Lys Ile Gln Ala Arg
    20                25

```

(2) INFORMATION FOR SEQ ID NO: 160:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 314 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(110..315)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 186..391
id AA046808
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(8..105)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 394..491
id AA046808
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(110..315)

(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 204..409
id AA156232
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(6..75)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 441..510
id AA156232
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 110..265
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 101..256
id AA147488
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 264..315
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 256..307
id AA147488
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 110..282
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 108..280
id AA157472
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 2..75
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 3..76
id AA157472
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 110..315
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 100..305
id AA046825
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 8..75
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..68
id AA046825
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 165..260
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.3
seq TVVLCVGCSTVLC/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 160:

```

AGCTTGCAGC TTGCTAGCTG TGTGGGCTGG GAGGTCTGGT AGGGCTGAGC TTGCAAGAGG      60
RTCAACATGC CTTTGGCTAG AGAYKTAMTA CATRCGTMCY TGGRRWGWGG YABAAGAVMA      120
MACATAAAAA GAANCGCCTA GTACAAAGTC CAAATTCTTA CTTT ATG GAT GTA AAA      176
                                   Met Asp Val Lys
                                   -30

TGT CCA GGT TGC TAC AAG ATC ACC ACG GTT TTC AGC CAT GCT CAG ACA      224
Cys Pro Gly Cys Tyr Lys Ile Thr Thr Val Phe Ser His Ala Gln Thr
      -25                      -20                      -15

GTG GTT CTT TGT GTA GGT TGT TCA ACA GTG TTG TGC CAG CCT ACA GGA      272
Val Val Leu Cys Val Gly Cys Ser Thr Val Leu Cys Gln Pro Thr Gly
      -10                      -5                      1

GGA AAG GCC AGA CTC ACA GAA GGG TGT TCA TTT AGG MGA AAG      314
Gly Lys Ala Arg Leu Thr Glu Gly Cys Ser Phe Arg Arg Lys
  5                      10                      15

```

(2) INFORMATION FOR SEQ ID NO: 161:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 120..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 183..290

id T34489
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 80..119
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 144..183
id T34489
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 120..227
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 141..248
id HUM416B01B
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 80..119
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 102..141
id HUM416B01B
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 120..220
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 140..240
id HUM425D11B
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 80..119
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 101..140
id HUM425D11B
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 120..227
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 28..135
id AA147546
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 120..154
(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 91
region 26..60
id AA103632
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 163..210
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.2
seq ILSVLHALPAGIA/WS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 161:

```
ACATTATCAA GAGAGAGAGA AGGTTAGGAT GAAAGAGATG AGCAGAGGCA AATCTTAAGT    60
GTTGACACCA TGATTGAAAA GGTTATTTGG AAATAGCGGT GCTTGCAGCG CYTGCGGAMC   120
RGTCGATTCC TCGGAGTGAA CTCGTCATGA GGGCGCAAGG CA ATG TGT ATC ATC       174
                                   Met Cys Ile Ile
                                   -15
TTA AGT GTT TTA CAT GCT CTA CCT GCC GGA ATC GCC TGG TCC CGG GAG       222
Leu Ser Val Leu His Ala Leu Pro Ala Gly Ile Ala Trp Ser Arg Glu
      -10                      -5                      1
AAA GGG                                                                228
Lys Gly
  5
```

(2) INFORMATION FOR SEQ ID NO: 162:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 255 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Surrrenals

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 58..255
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 176..373
id AA082793
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(97..255)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98

region 352..510
id AA129762
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(66..97)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 511..542
id AA129762
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 89..214
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 11..136
id H56715
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 207..255
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 93
region 128..176
id H56715
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(191..255)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 355..419
id AA101128
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(151..189)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 423..461
id AA101128
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(119..151)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 462..494
id AA101128
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 46..225

(C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.9
 seq TWLLLGALPASE/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 162:

```

AGCAGCTGTT TCGGTAAC TG CTTGCCTCC CGGCTCCCGC AGWGG ATG CTG GTG GTG   57
                                     Met Leu Val Val
                                     -60

GAG GCT TCT TCC TCA GTG CGG CTG GCA AGT TCG GAG GTG ACT TCC TGG   105
Glu Ala Ser Ser Ser Val Arg Leu Ala Ser Ser Glu Val Thr Ser Trp
-55                               -50                               -45

TCT ATC CTG GTG ACC CCC TCC GCT TCC ACG CCC ATT ATA TCG CTC AGT   153
Ser Ile Leu Val Thr Pro Ser Ala Ser Thr Pro Ile Ile Ser Leu Ser
-40                               -35                               -30                               -25

GCT GGG CCC CTG AGG ACA CCA TCC CAC TCC AAG ACC TGG TTG CTG CTG   201
Ala Gly Pro Leu Arg Thr Pro Ser His Ser Lys Thr Trp Leu Leu Leu
                               -20                               -15                               -10

GGC GCC TTG GAA CCA GCG TCA GAA AGA CCC TGC TCC TCT GTT CTC CGC   249
Gly Ala Leu Glu Pro Ala Ser Glu Arg Pro Cys Ser Ser Val Leu Arg
                               -5                               1                               5

AGC CGG   255
Ser Arg
10
  
```

(2) INFORMATION FOR SEQ ID NO: 163:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 285 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

-(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 22..201
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 5..184
id W07565
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 200..282
- (C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 100
region 184..266
id W07565
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 132..209
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 102..179
id R58430
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 208..282
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 177..251
id R58430
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 29..80
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 1..52
id R58430
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 82..132
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 53..103
id R58430
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 7..118
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 1..112
id W19152
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 175..282
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 109..216
id W19152
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 175..282
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 95..202
id AA095767
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 69..118
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 49..98
id AA095767
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 20..74
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 1..55
id AA095767
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 175..282
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 99..206
id T86637
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 22..118
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 6..102
id T86637
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 181..261
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.7
seq LSLQLIAFPTVSC/EI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 163:

```
CGCCTCCTCC GCTTTGGGAG CMCCGGGCTA MTCTTTCACA GCCCTGTTG CCCTGTGATC   60
TG TAGGTCCT TGGGGACGCA CAGTTAAGAT GACAGGACAT CCTGGAAGCT GGGAAATGGC  120
TGAATGCTAT CCCAGTGAAT ATACGTGCCC TGTTTGTGTA ATCTACTCAT CCTTAAAGAT  180
ATG TAT TCA TTT CCT ACC ACC GTA GTG GAA GAG ATA CTA TCC CTA TCT    228
Met Tyr Ser Phe Pro Thr Thr Val Val Glu Glu Ile Leu Ser Leu Ser
```

(2) INFORMATION FOR SEQ ID NO: 164:

(A) LENGTH: 174 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(A) NAME/KEY: other
(B) LOCATION: 75..168
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 91
region 200..293
id N57841
est

(A) NAME/KEY: sig_peptide
(B) LOCATION: 115..162
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.6
seq LLPLRSLLALVRE/SR

AAAGCTTTTAA CGAGCTTATA AATCACTTG	TGAC CCCACTTTAT GTAATGTGAT	60
TCTGCAGGTT TGAAAAAGGT CCATAAATAG	GTGTTTTAAA CAAGTTTCCT GTCA ATG	117
	Met	
CTG ATG TTG CTC CCC CTT CGC TCA TTA TTA GCA	TTA GTT AGA GAA AGC	165
Leu Met Leu Leu Pro Leu Arg Ser Leu Leu Ala	Leu Val Arg Glu Ser	
-15	-10 -5 1	
AGG GCA CGG		174
Arg Ala Arg		

(2) INFORMATION FOR SEQ ID NO: 165:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 407 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Cerebellum
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 37..217
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 1..181
 id AA057016
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 216..378
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 179..341
 id AA057016
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 34..217
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 98
 region 1..184
 id AA133917
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 216..327
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 94
 region 182..293
 id AA133917
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 325..366
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 290..331
 id AA133917
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 216..342
 (C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 99
region 119..245
id R13065
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 114..217
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 18..121
id R13065
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 99..173
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.5
seq AQLFACLLRLGTQ/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 165:

```

AAGTCCAGGC CTTGAGACCC AGAAGGGAGC GAAGGTTTTT GCTGCGCCAA CGCAGTGAGC   60
CGAAGCTCCG CTCACGCCCG GCCTGATCCT GCCTGAAG ATG GTG CCA CTG GTG GCT   116
                               Met Val Pro Leu Val Ala
                               -25                -20

GTG GTA TCA GGG CCC CGT GCC CAG CTC TTT GCC TGC CTG CTC AGG CTG   164
Val Val Ser Gly Pro Arg Ala Gln Leu Phe Ala Cys Leu Leu Arg Leu
                               -15                -10                -5

GGC ACT CAG CAG GTC GGC CCC CTT CAG CTG CAC ACC GGG GCC AGC CAT   212
Gly Thr Gln Gln Val Gly Pro Leu Gln Leu His Thr Gly Ala Ser His
                               1                5                10

GCG GCC AGG AAC CAT TAT GAG GTG CTG GTG CTG GGT GGG GGC AGT GGC   260
Ala Ala Arg Asn His Tyr Glu Val Leu Val Leu Gly Gly Gly Ser Gly
                               15                20                25

GGA ATC ACC ATG GCT GCC CGC ATG AAG AGG AAA GTG GGT GCA GAG AAT   308
Gly Ile Thr Met Ala Ala Arg Met Lys Arg Lys Val Gly Ala Glu Asn
                               30                35                40                45

GTG GCC ATT GTT GAG CCC AGT GAG AGA CAT TTC TAC CAG CCA ATC TGG   356
Val Ala Ile Val Glu Pro Ser Glu Arg His Phe Tyr Gln Pro Ile Trp
                               50                55                60

ACA CTG GTG GGT GCT GGT GCA ADC AAT TGT CCT CAT CTG GTC GTC CCA   404
Thr Leu Val Gly Ala Gly Ala Xaa Asn Cys Pro His Leu Val Val Pro
                               65                70                75

CGG   407
Arg

```

(2) INFORMATION FOR SEQ ID NO: 166:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 184 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 18..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 18..180
id AA110680
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 40..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 64..204
id W30470
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 91..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 46..135
id HSCOCC021
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 91..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 23..112
id HUMHG5097
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 80..180
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 16..116
id AA089700
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 110..169
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.4
seq AFVIACVLSLIST/IY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 166:

```
AGAGCTAGAG GGTGAAGCTG GCGGAGCAGG AGGATGGGCG TATGCAGGTG ATAGACTAGA    60
GAACAAGACC TCTGTCTCCG TAGCATCCTG GAGCAGTCTG AATGCCAGA ATG GAT AAC    118
                                         Met Asp Asn
                                         -20

CGT TTT GCT ACA GCA TTT GTA ATT GCT TGT GTG CTT AGC CTC ATT TCC    166
Arg Phe Ala Thr Ala Phe Val Ile Ala Cys Val Leu Ser Leu Ile Ser
   -15                -10                -5

ACC ATC TAC ATG GCC CGG                                184
Thr Ile Tyr Met Ala Arg
   1                5
```

(2) INFORMATION FOR SEQ ID NO: 167:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 371 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(1..223)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 277..499
id AA059399
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(236..330)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 170..264
id AA059399
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(332..371)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 130..169
id AA059399

est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(6..143)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 357..494
id N35134
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(236..371)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 128..263
id N35134
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(141..223)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 276..358
id N35134
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(44..223)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 223..402
id H49423
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(236..371)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 75..210
id H49423
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 14..190
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 1..177
id H43799
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 236..342
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98

region 224..330
id H43799
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 338..369
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 93
region 327..358
id H43799
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(85..223)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 223..361
id H27862
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(266..346)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 101..181
id H27862
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(236..269)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 177..210
id H27862
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 12..137
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.4
seq WLLLLTSLDGHLL/EP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 167:

AGGATGACCT G ATG CCT GAG TAC TGT GGC AAT GAG GTG ACT CCA ACC GAG	50
Met Pro Glu Tyr Cys Gly Asn Glu Val Thr Pro Thr Glu	
-40 -35 -30	
GCT GCC CAA GCG CCA GAG GTG ACC TAT GAG GCA GAA GAG GGC TCC TTG	98
Ala Ala Gln Ala Pro Glu Val Thr Tyr Glu Ala Glu Glu Gly Ser Leu	
-25 -20 -15	
TGG ACG TTG CTA CTC ACT AGC TTG GAT GGG CAC CTG CTG GAG CCA GAT	146
Trp Thr Leu Leu Leu Thr Ser Leu Asp Gly His Leu Leu Glu Pro Asp	
-10 -5 1	

```

GCT GAG TAC CTC CAC TGG CTG CTA ACC AAC ATC CCG GGT AAC CGG GTG      194
Ala Glu Tyr Leu His Trp Leu Leu Thr Asn Ile Pro Gly Asn Arg Val
   5                      10                      15

GCT GAA GGA CAG GTG ACG TGT CCC TAC CTC CCC CCC TTC CCT GCC CGA      242
Ala Glu Gly Gln Val Thr Cys Pro Tyr Leu Pro Pro Phe Pro Ala Arg
  20                      25                      30                      35

GGC TCC GGC ATC CAC CGT CTT GCC TTC CTG CTC TTC AAG CAG GAC CAG      290
Gly Ser Gly Ile His Arg Leu Ala Phe Leu Leu Phe Lys Gln Asp Gln
                     40                      45                      50

CCG ATT GAC TTC TCT GAG GAC GCA CGC CCC TCA CCC TGC TAT CAG CTG      338
Pro Ile Asp Phe Ser Glu Asp Ala Arg Pro Ser Pro Cys Tyr Gln Leu
   55                      60                      65

GYS CAG CGG ACC TTC CGC ACT TTT GAT TTC TAC                          371
Xaa Gln Arg Thr Phe Arg Thr Phe Asp Phe Tyr
   70                      75

```

(2) INFORMATION FOR SEQ ID NO: 168:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 118 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrrenals

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 51..114
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 112..175
id AA029014
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 20..55
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..36
id AA029014
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 5..55
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..51

id AA029165
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 70..114
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 66..110
id AA029165
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 29..58
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 8..37
id W79853
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 23..67
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.4
seq RVLCPAAGAVRA/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 168:

```
AAGTCGAGGA GTCAAGGCAG CA ATG AAT CGT GTC TTG TGT GCC CCG GCG GCC      52
                        Met Asn Arg Val Leu Cys Ala Pro Ala Ala
                        -15                               -10

GGG GCC GTC CGG GCG CTG AGG CTC ATA GGC TGG GCT TCC CGA AGC CTT      100
Gly Ala Val Arg Ala Leu Arg Leu Ile Gly Trp Ala Ser Arg Ser Leu
-5              1              5              10

CAT CCG TTG CCC GGA AAG      118
His Pro Leu Pro Gly Lys
              15
```

(2) INFORMATION FOR SEQ ID NO: 169:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 183 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 123..181
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 133..191
id W02973
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 4..55
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 13..64
id W02973
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 56..102
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 66..112
id W02973
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 55..181
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 38..164
id T97581
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 19..56
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 1..38
id T97581
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 21..181
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 91
region 58..218
id W70849
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 21..181
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 91
region 47..207
id AA049525
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 22..181
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 4..163
id H32732
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 49..117
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3
seq MLALLLTAALIFF/AI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 169:

```
ACTTTCTCCG CTGGCAACGG CGCCGCTCCC CGCTCCTCCT CCCAGGCC ATG GCG TTC      57
                                     Met Ala Phe

ACG TTC GCG GCC TTC TGC TAC ATG CTG GCG CTG CTG CTC ACT GCC GCG      105
Thr Phe Ala Ala Phe Cys Tyr Met Leu Ala Leu Leu Leu Thr Ala Ala
-20                -15                -10                -5

CTC ATC TTC TTC GCC ATT TGG CAC ATT ATA GCA TTT GAT GAG CTG AAG      153
Leu Ile Phe Phe Ala Ile Trp His Ile Ile Ala Phe Asp Glu Leu Lys
                        1                5                10

ACT GAT TAC AAG AAT CCT ATA GAC CAG TTG                                183
Thr Asp Tyr Lys Asn Pro Ile Asp Gln Leu
      15                20
```

(2) INFORMATION FOR SEQ ID NO: 170:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 169 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 107..143
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 125..161
id N86955
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 89..139
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3
seq XEXLLAFHHDCEA/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 170:

```

AATAGGTAAT GAGTCTTATG AGATCTGTTG GTTTTAAAAA CAGGAGTTTC TCTGCGCAAS    60
CTCTGTCTTT TTTTGCCCTGC TGGCATCC ATG CRA RRC RWG AST GAG YTC CTC    112
                               Met Xaa Xaa Xaa Xaa Glu Xaa Leu
                               -15                               -10

CTT GCC TTC CAC CAT GAT TGT GAG GCT TCC CCA GCC ACG TGG AAC TTA    160
Leu Ala Phe His His Asp Cys Glu Ala Ser Pro Ala Thr Trp Asn Leu
                -5                1                5

AGT CCA AGG                                169
Ser Pro Arg
                10

```

(2) INFORMATION FOR SEQ ID NO: 171:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 240 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(15..236)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 314..535
id AA194996
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 13..120
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq PLRLNLLILIEG/SV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 171:

```

TTATATAGAG CC ATG GGG CCT TAC AAC GTG GCA GTG CCT TCA GAT GTA TCT    51
Met Gly Pro Tyr Asn Val Ala Val Pro Ser Asp Val Ser

```

	-35	-30	-25	
CAT GCC CGC TTT TAT TTC TTA TTT CAT CGA CCA TTA AGG CTG TTA AAT				99
His Ala Arg Phe Tyr Phe Leu Phe His Arg Pro Leu Arg Leu Leu Asn				
	-20	-15	-10	
CTG CTC ATC CTT ATT GAG GGC AGT GTC GTC TTC TAT CAG CTC TAT TCC				147
Leu Leu Ile Leu Ile Glu Gly Ser Val Val Phe Tyr Gln Leu Tyr Ser				
	-5	1	5	
TTG CTG CGG TCG GAG AAG TGG AAC CAC ACA CTT TCC ATG GCT CTC ATC				195
Leu Leu Arg Ser Glu Lys Trp Asn His Thr Leu Ser Met Ala Leu Ile				
	10	15	20	25
CTC TTC TGC AAC TAC TAT GTT TTA TTT AAA CTT CTC CGG GAT CAG				240
Leu Phe Cys Asn Tyr Tyr Val Leu Phe Lys Leu Leu Arg Asp Gln				
	30	35	40	

(2) INFORMATION FOR SEQ ID NO: 172:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 31..207
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..177
id W44975
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 45..207
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..163
id N43016
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..207
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..161
id R10507
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 49..207
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..159
id H70117
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 48..207
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..160
id H61870
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 67..135
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq IGVGLYLLASAAA/FY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 172:

```
AGCGGCGGCA TCCGGGACGG CGGGCGGGCT GGCCACCACG GGACAGGAAG GCACAGAGCA    60
TGGAGA ATG ATG AAC TTC CGT CAG CGG ATG GGA TGG ATT GGA GTG GGA    108
      Met Met Asn Phe Arg Gln Arg Met Gly Trp Ile Gly Val Gly
                -20                -15                -10

TTG TAT CTG TTA GCC AGT GCA GCA GCA TTT TAC TAT GTT TTT GAA ATC    156
Leu Tyr Leu Leu Ala Ser Ala Ala Ala Phe Tyr Tyr Val Phe Glu Ile
                -5                1                5

AGT GAG ACT TAC AAC AGG CTG GCC TTG GAA CAC ATT CAA CAG CAC CHT    204
Ser Glu Thr Tyr Asn Arg Leu Ala Leu Glu His Ile Gln Gln His Xaa
      10                15                20

GGG
Gly    207
```

(2) INFORMATION FOR SEQ ID NO: 173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 487 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 71..252
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 149..330
id R18686
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 252..304
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 331..383
id R18686
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 23..73
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..51
id R18686
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 186..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 1..190
id R54039
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 376..489
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 190..303
id R54039
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 94..303
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 70..279
id HSC2LA121
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 26..97
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 1..72

id HSC2LA121
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 236..375
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 1..140
id N91698
est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 376..458
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 95
 region 140..222
 id N91698
 est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 458..489
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 223..254
id N91698
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 98..451
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.1
seq AFXVVCWLGPCA/MH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 173:

AAGCGCCCCG	VGSSCGCGTC	CCCGGCCCAA	CCATGGCGTC	CTCCGCGGCC	GGCTGCGTGG	60
TGATCGTTGG	CAGTGGAGTC	ATTGGGCGAR	STGGGCC	ATG CTG TTT GCC AGT GGA	115	
				Met Leu Phe Ala Ser Gly		
				-115		
GGC TTC CAK GTG AAA CTC TAT GAC ATT GAG CAA CAG CAG ATA AGG AAC	163					
Gly Phe Xaa Val Lys Leu Tyr Asp Ile Glu Gln Gln Gln Ile Arg Asn						
-110 -105 -100						
GCC CTG GAA AAC ATC AGA AAG GAG ATG AAG TTG CTG GAG CAG GCA GGT	211					
Ala Leu Glu Asn Ile Arg Lys Glu Met Lys Leu Leu Glu Gln Ala Gly						
-95 -90 -85						
TCT CTG AAA GGC TCC CTG AGT GTG GAA GAG CAG CTG TCA CTC ATC AGT	259					
Ser Leu Lys Gly Ser Leu Ser Val Glu Glu Gln Leu Ser Leu Ile Ser						
-80 -75 -70 -65						
GGT TGT CCC AAT ATC CAA GAA GCA GTA GAG GGT GCC ATG CAC ATT CAG	307					
Gly Cys Pro Asn Ile Gln Glu Ala Val Glu Gly Ala Met His Ile Gln						
-60 -55 -50						

GAA TGT GTT CCA GAA GAT CTA GAA CTG AAG AAG AAG ATT TTT GCT CAG	355
Glu Cys Val Pro Glu Asp Leu Glu Leu Lys Lys Lys Ile Phe Ala Gln	
-45 -40 -35	
TTA GAT TCC ATC ATT GAT GAA TCG AGT GAT CTT AAG CGT TTC CAM TTC	403
Leu Asp Ser Ile Ile Asp Glu Ser Ser Asp Leu Lys Arg Phe Xaa Phe	
-30 -25 -20	
TTG TCT CAT GCC TTC VAA GTT GTT TGC TGG CTT GGT CCA TGT GAA GCA	451
Leu Ser His Ala Phe Xaa Val Val Cys Trp Leu Gly Pro Cys Glu Ala	
-15 -10 -5	
ATG CAT CGT GGM TCA TCC TGT GAA TCC GCC ATA CTA	487
Met His Arg Gly Ser Ser Cys Glu Ser Ala Ile Leu	
1 5 10	

(2) INFORMATION FOR SEQ ID NO: 174:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(47..141)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 236..330
id R62451
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(78..141)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 224..287
id C14686
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 78..141
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..64
id C15352
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 78..141
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..64
id C15019
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(78..141)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 225..288
id C14869
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 39..104
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq LCSLPLSPSAVCP/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 174:

```
ACACACAAAA ATTTTCCTTT GTTTCGCGGG GGCTGGGG ATG CAG TGT TTT TTG GGG    56
                                     Met Gln Cys Phe Leu Gly
                                     -20

GGT CTT GGT TTA TGC TCC CTG CCC TTG AGC CCC TCA GCC GTT TGC CCT    104
Gly Leu Gly Leu Cys Ser Leu Pro Leu Ser Pro Ser Ala Val Cys Pro
-15                      -10                      -5

GCC CCC ACC TCG GCT CCA TGG TGG GAG GGG GCT CTG    140
Ala Pro Thr Ser Ala Pro Trp Trp Glu Gly Ala Leu
 1                      5                      10
```

(2) INFORMATION FOR SEQ ID NO: 175:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 363 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 37..288
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 4..255

id R13070
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 287..353
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 255..321
id R13070
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 175..213
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.9
seq MSSFLLSFSQSLS/NV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 175:

```

ACTCTTTACT CCCCTGTGAG TGATTCACTG CCTGTGCATT ATTACGATAG ATGTGTTTGT      60
ATTGKTTTTT TTCTGATGAT ACTGATGTTG ATGAATTTTT AATTTTATTT GATGTGGTAG      120
AGTTGGGAGG TTTCAGGGTT TTTTCCCCTC TTTTACTTTC CATTGAGGAA GGA ATG      177
                                         Met
AGC TCC TTT CTC CTC TCC TTC AGC CAA TCA TTA TCA AAT GTT CCT TCA      225
Ser Ser Phe Leu Leu Ser Phe Ser Gln Ser Leu Ser Asn Val Pro Ser
      -10                      -5                      1

GCC CTG CAG TKG CCC CAA ATA ACC TTT TTT CAG CAT CCT CTG TCC TCA      273
Ala Leu Gln Xaa Pro Gln Ile Thr Phe Phe Gln His Pro Leu Ser Ser
      5                      10                      15                      20

GTC ATG CCA GTC TGG ACA TGC TCT GTT GTG CCC TGT GAC AAA ACT GST      321
Val Met Pro Val Trp Thr Cys Ser Val Val Pro Cys Asp Lys Thr Xaa
      25                      30                      35

CAG TAT TCC TAT TGC TTT TAC TGT GTT TTA GGT ACT GTG AAG      363
Gln Tyr Ser Tyr Cys Phe Tyr Cys Val Leu Gly Thr Val Lys
      40                      45                      -50

```

(2) INFORMATION FOR SEQ ID NO: 176:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 379 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(216..354)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 82..220
id W28597
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(351..381)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 54..84
id W28597
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 216..347
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99
region 121..252
id T90405
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 249..310
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90
region 141..202
id AA085837
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 293..338
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 186..231
id AA085837
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 216..248
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 107..139
id AA085837
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 351..381
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 246..276
id AA085837

est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 152..190
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq MLTASLAFQLVDG/VS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 176:

```

AGAATCTGCT CGCTCTTCTG GAGCTGAGAC ACTCATCTTC TCCTGCCCTG GCACATCAAA   60
ACTCCAGGTT CTCTGATCTT TGAACACTGG GACTTAATCC AGCATCCCCT TACTGCCTGA  120
AGTTCTCAGG ACTTTGGACT TGGACTGAAC G ATG CTA ACA GCT TCC CTG GCT   172
                               Met Leu Thr Ala Ser Leu Ala
                               -10

TTC CAG CTT GTA GAT GGC GTA TCA TGG AAT TTC TCA GTC TCC AAA ATG   220
Phe Gln Leu Val Asp Gly Val Ser Trp Asn Phe Ser Val Ser Lys Met
   -5              1              5              10

CTG GCA TCA CCA TCT ACA TCA GGT CAG CTG TCT CAG TTT GGG GCA AGT   268
Leu Ala Ser Pro Ser Thr Ser Gly Gln Leu Ser Gln Phe Gly Ala Ser
              15              20              25

TTA TAC GGG CAA CAA AGT GCA CTA GGC CTT CCA ATG AGG GGG ATG AGC   316
Leu Tyr Gly Gln Gln Ser Ala Leu Gly Leu Pro Met Arg Gly Met Ser
              30              35              40

AAC AAT ACC CCT CAG TTA AAT CGC AGC TTA TCA CAA GSA CTC AGT TAC   364
Asn Asn Thr Pro Gln Leu Asn Arg Ser Leu Ser Gln Xaa Leu Ser Tyr
              45              50              55

CGA GCC ACG TCA CGC                                           379
Arg Ala Thr Ser Arg
   60

```

(2) INFORMATION FOR SEQ ID NO: 177:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 198 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(2..132)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99

region 105..235
id T58540
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 118..192
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq LSAFNFLVCLSLG/RG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 177:

```
ATATGATGCT GGCAAGACAC CCAGAAAGGC TATTTTCAGA TGAAATCGAT ATTAGAAGCT 60
ATATTAGCTG AAACAACCTCC TTTTACTGCG TAGAACCTAT ATCGAGAGTG TGTGTAT 117
ATG TAT TAW AGG AGG GAG CTC TCA ATT TTA TGT ATT CTT TCT GCC TTT 165
Met Tyr Xaa Arg Arg Glu Leu Ser Ile Leu Cys Ile Leu Ser Ala Phe
-25 -20 -15 -10
AAT TTT CTT GTT TGT TTG AGC TTA GGG AGA GGG 198
Asn Phe Leu Val Cys Leu Ser Leu Gly Arg Gly
-5 1
```

(2) INFORMATION FOR SEQ ID NO: 178:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 292 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 56..93
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 213..250
id N58549
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 206..271
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8
seq IVFGVSWVMLVYS/AS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 178:

```

ATGWAGGCTG TTGTTTCAGAT TCCTTTGTCC CATGGGGTGT TCCCTTATTA TAGTACTCTC   60
GCCCTTTTCC TATGGATGTG GCTTCCTGAG AGCGGMSTGA TASTGATTGT KATCTTTCTT   120
KTGGATCTAS CCACCCAGCT ATTCTACCAK GCTCTGGGCT GGTACTGGCG GTTGTCTGCA   180
CAGAGTCTTG TGACCTGAAC CATCT ATG GGT CTC TCW GCC ATG GAT ACC AGC   232
                               Met Gly Leu Ser Ala Met Asp Thr Ser
                               -20                               -15
ATA GTA TTT GGG GTG TCC TGG GTC ATG CTG GTG TAC TCT GCT TCC TTC   280
Ile Val Phe Gly Val Ser Trp Val Met Leu Val Tyr Ser Ala Ser Phe
                               -10                               -5                               1
AGG AGG TGT GRN   292
Arg Arg Cys Xaa
5

```

(2) INFORMATION FOR SEQ ID NO: 179:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 307 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(80..196)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 68..184
id R85971
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(31..86)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 179..234
id R85971
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(80..196)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 68..184
id R85965
est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: complement(31..86)
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 179..234
 id R85965
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 205..305
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 95
 region 128..228
 id N33907
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 205..305
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 95
 region 99..199
 id T19230
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 205..305
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 95
 region 96..196
 id T31783
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 101..295
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.8
 seq IILFSAIVGFIYG/YV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 179:

```

AAAAAAGGTC AAGGTAGCCT CCTCCAACT CCACGTTGAG CTGCACCTCC CCTGAGCTCC      60
CACATCTGGG CCTCCCTCTC CCAGGGCTAA TCCAGCTGTC ATG TAT TTC TGG CGT      115
                               Met Tyr Phe Trp Arg
                               -65

GAT GTG GCT GTC TCC CTT GAC ACG CTC TGG GCT CTT CCA AGG CAA CAG      163
Asp Val Ala Val Ser Leu Asp Thr Leu Trp Ala Leu Pro Arg Gln Gln
-60                -55                -50                -45

CCT GGT CTT GGT AAC AAC CGT GTC CTC GGT CTG CTG TCT GGC ACA AAC      211
Pro Gly Leu Gly Asn Asn Arg Val Leu Gly Leu Leu Ser Gly Thr Asn
          -40                -35                -30

AAG GAT TAC AAG GGC CAG AAG CTA GCT GAA CAG ATG TTT CAG GGA ATT      259

```

ATT CTT TTT TCT GCA ATA GTT GGA TTT ATC TAC GGG TAC GTG GCC GCG 307
Ile Leu Phe Ser Ala Ile Val Gly Phe Ile Tyr Gly Tyr Val Ala Ala
-10 -5 1

ATTACGCAGA GAGAAAGTTA CGAGGTTTCGT GGCCGCGGTT TCCCCAGGCA GCTGGCGCTG	60
GAGGCTTCGG CGTCACGTGC TGGTCTGGRT TTTTCTCG ATG CAC TGG GGA AAG CGG	116
Met His Trp Gly Lys Arg	
-25	
TGG RCT CTT RTC GWR GGA GGG CTC TTG ATC TGT GRT TTA TRG ATA GGC	164
Trp Xaa Leu Xaa Xaa Gly Gly Leu Leu Ile Cys Xaa Leu Xaa Ile Gly	
-20 -15 -10 -5	
ACA GCT ACT CCC GTT CGG GRA CCC AAC GGC AGA CAG GTS CTA GTG CCC	212
Thr Ala Thr Pro Val Arg Xaa Pro Asn Gly Arg Gln Val Leu Val Pro	
1 5 10	
RTC GGW TAC CCG CGG CCG GGA CTC GGA GCT GTG GGG TGT GGG GAG GCG	260
Xaa Gly Tyr Pro Arg Pro Gly Leu Gly Ala Val Gly Cys Gly Glu Ala	
15 20 25	

(2) INFORMATION FOR SEQ ID NO: 181:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 416 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 158..364
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 153..359
id N25870
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 358..416
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 355..413
id N25870
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 148..332
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 134..318
id AA045920
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 332..395
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 319..382
id AA045920
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 205..416
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 1..212
id AA150024
est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 148..315
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 133..300
 id H99323
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 352..403
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 92
 region 337..388
 id H99323
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: complement(202..384)
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 93
 region 215..397
 id R83259
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 309..395
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.6
 seq ALGLXTCLSVLFG/YA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 181:

```

AGCASGGAAC AGCGGGTGCG GACATTACGG CGGAAGGCTC TSGAGGAAGC AGAAGTGAAG   60
GACCTCGCAS TCCTGGGACG GTGGGGCMCA GASAGAGAAA GGGAGCCCCG GGC GCGGCGC   120
GGTGAGGATG CGAGCAGAGG AAGGACACGC GGCGCCGSAA AATATTTACA CCAGCAGCTC   180
CAGTTCATAC CMMTAAAGAM SATCCTGCTA CCCAMACTAM TTTGGGATTT ATCCMYGCAT   240
TTGTCGCTGC CATATCAGTT ATTATTGTAT CTGAATTGGG TGATAAGACA TTTTTTATAG   300
CAGCCATC ATG GCA MCG CGC TAT AAC CGC CTG ACC GTG CTG GCT GGT GCM   350
      Met Ala Xaa Arg Tyr Asn Arg Leu Thr Val Leu Ala Gly Ala
      -25                               -20

MTG CTT GCC TTG GGA CTA MTG ACA TGC YTG TCA GTT TTG TTT GGC TAT   398
Xaa Leu Ala Leu Gly Leu Xaa Thr Cys Leu Ser Val Leu Phe Gly Tyr
-15                -10                -5                1

GCA CCA CAG TCA TCC CCA                                           416
Ala Pro Gln Ser Ser Pro
      5

```

(2) INFORMATION FOR SEQ ID NO: 182:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 247 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 55..152
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 92
 region 188..285
 id N94950
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 54..152
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 122..220
 id T09182
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: 170..233
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 90
 region 296..359
 id N71787
 est
- (ix) FEATURE:
 (A) NAME/KEY: other
 (B) LOCATION: complement(54..142)
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 93
 region 199..287
 id T64627
 est
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: 143..229
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.5
 seq FMTCILCRPPISS/CV
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 182:

```

GTATTTGTAA ACTGTCATGG TGCTGTTGGG AGTGTAGCAG TGAGGACGAC CAGAGGTCAC 120
TCTTGTCGCC ATCTTGTTT TG ATG GGT TTT ACT GGC TTC TTT ACT GCA ACC 172
      Met Gly Phe Thr Gly Phe Phe Thr Ala Thr
                        -25                -20

TGT TTT ATC AGC AAG GTC TTT ATG ACC TGT ATC TTG TGC AGA CCT CCT 220
Cys Phe Ile Ser Lys Val Phe Met Thr Cys Ile Leu Cys Arg Pro Pro
      -15                -10                -5

ATC TCA TCC TGT GTC TTA GAA TGC GGG 247
Ile Ser Ser Cys Val Leu Glu Cys Gly
      1                5

```

(2) INFORMATION FOR SEQ ID NO: 183:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(61..347)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 88..374
id W84548
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(1..48)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 387..434
id W84548
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(61..282)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 154..375
id N66815
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(279..347)

(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 88..156
id N66815
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(12..48)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 388..424
id N66815
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(89..282)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 155..348
id N24160
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(279..347)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 89..157
id N24160
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(200..347)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 88..235
id N66833
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(239..347)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 11..119
id H88111
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(208..239)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 120..151
id H88111
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 258..362
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.4
 seq LIVLLPVLFSLK/NF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 183:

```

AACACAAATA GCACTGTCAC CTCTAATATG AACATTAGTT TGAGGTAGTT TTTTCTAAA   60
GCAAAAATTT TAACTGTTTT CTAATTGTCA AGCACTATTT TCATTAAAAG TGTCTAATGA  120
ATCATGATAT ACTCTCCAT TTGTTGTGTC TATTTTTTAT ATATTGGTA TTTTTTGAAA  180
ATTCCAAATA CTCATGTCTC AAGTAAGCTT AAACACAAC TTGTCACATA AAGGAAGTCT  240
TAAGTGGAGT TCACAGA ATG ATA ATG TAT CTA TTT GTC ATT TGT GTT ATA   290
          Met Ile Met Tyr Leu Phe Val Ile Cys Val Ile
          -35                      -30                      -25

TTT GAA ATT ATT AGA AAT TAT GCT TTT TCC ATT TTA ATT GTA TTG CTG   338
Phe Glu Ile Ile Arg Asn Tyr Ala Phe Ser Ile Leu Ile Val Leu Leu
          -20                      -15                      -10

CCA GTG CTA TTT TTT TCT TTA AAA AAT TTT ATT CTT AGC ACA CAG       383
Pro Val Leu Phe Phe Ser Leu Lys Asn Phe Ile Leu Ser Thr Gln
          -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 184:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 349 base pairs
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: DOUBLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 63..281
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 1..219
 id R06709
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 166..336
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 96
 region 104..274
 id W92876

est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 63..165
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 2..104
id W92876
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 165..351
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 63..249
id N31364
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 102..165
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..64
id N31364
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 166..351
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 64..249
id N75642
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 105..165
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 4..64
id N75642
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(269..324)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 259..314
id W92774
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(322..351)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93

region 231..260
id W92774
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 176..286
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3
seq LWEKLTLLSPGIA/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 184:

```
AAGTAAACAT GAGCCTCTGG TAGATAAGAG AAACCGCGAT CGGAGTACGG CGCGTGCGWG      60
NATCAGGGAT CGCGATTGCG AATCCTCCGC TGAGGTGATT TGGATATCCC TAGAACGTTG      120
AGGGCACGAG TCGGGTCCTG AGACCAGGTC CTCAGCCAGC AGAGCCACGT TCCTT ATG      178
                                         Met
AGC ACC GTG GGT TTA TTK CAT TTT CCT ASR CCA CTG ACC CGA ATA TGC      226
Ser Thr Val Gly Leu Xaa His Phe Pro Xaa Pro Leu Thr Arg Ile Cys
-35                               -30                               -25

CCG GCG CCA TGG GGA CTC CGG CTT TGG GAG AAG CTG ACG TTG TTA TCC      274
Pro Ala Pro Trp Gly Leu Arg Leu Trp Glu Lys Leu Thr Leu Leu Ser
-20                               -15                               -10                               -5

CCA GGA ATA GCT GTS ACT CCG GTC CAG ATG GCA GGC AAG AAG GAC TAS      322
Pro Gly Ile Ala Val Thr Pro Val Gln Met Ala Gly Lys Lys Asp Xaa
                               1                               5                               10

CCT GCA CTG CTT TCC TTG GAT GAG AAT      349
Pro Ala Leu Leu Ser Leu Asp Glu Asn
15                               20
```

(2) INFORMATION FOR SEQ ID NO: 185:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 268 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 64..107
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 336..379
id AA100380
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 32..70
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3
seq MLALAXHLSTVES/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 185:

```

CATTGAGCTC GGGCTGAGTG AGGCCAGGT G ATG CTG GCT CTA GCC ASN CAC      52
                               Met Leu Ala Leu Ala Xaa His
                               -10

CTG AGC ACA GTG GAG TCG GAG AAA CAG AAG CTG CGG GCT CAG GTG CGG    100
Leu Ser Thr Val Glu Ser Glu Lys Gln Lys Leu Arg Ala Gln Val Arg
-5                      1                      5                      10

CGG CTA TGC CAG GAG AAC CAG TGG CTG CGG GAT GAG CTG GCT GGC ACC    148
Arg Leu Cys Gln Glu Asn Gln Trp Leu Arg Asp Glu Leu Ala Gly Thr
                      15                      20                      25

CAG CAG CGG CTA CAG CGC AGT GAA CAG GCT GTG GCT CAG CTG GAG GAG    196
Gln Gln Arg Leu Gln Arg Ser Glu Gln Ala Val Ala Gln Leu Glu Glu
                      30                      35                      40

GAA AAG AAG CAC CTG GAG TTC CTG GGG CAG CTG CGG CAG TAT GAT GAG    244
Glu Lys Lys His Leu Glu Phe Leu Gly Gln Leu Arg Gln Tyr Asp Glu
                      45                      50                      55

GAT GGA CAT ACC TCG GAG GCG GGG                                268
Asp Gly His Thr Ser Glu Ala Gly
60                      65

```

(2) INFORMATION FOR SEQ ID NO: 186:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 305 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 14..122
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..109
id HUM429E03B
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 114..230
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 100..216
id HUM429E03B
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 226..307
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 211..292
id HUM429E03B
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 114..274
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92
region 85..245
id T31768
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 29..122
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..94
id T31768
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 262..307
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 234..279
id T31768
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 103..307
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 95..299
id T80259
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 21..120
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 15..114
id T80259

est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 114..307
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 68..261
id N32697
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 45..87
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 1..43
id N32697
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 92..122
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 93
region 47..77
id N32697
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 114..307
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 59..252
id N44613
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 55..122
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 94
region 1..68
id N44613
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 147..248
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.3
seq QLFAFLNLLPVEA/DI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 186:

AGTCGTCCCT GCTAGTACTC CGGGCTGTGG GGGTCGGTGC GGATATTGAG TCATGAAATC 60

AGGGTAGGGA CTTCTCCCGC AGCGACGCGG CTGGCAAGAC TGTTTGTGTT GCGGGGGCCG 120

```

GAACTTCAAG GTGATTTTAC AACGAG ATG CTG CTC TCC ATA GGG ATG CTC ATG   173
                        Met Leu Leu Ser Ile Gly Met Leu Met
                        -30

CTG TSA GCC ACA CAA GTC TAC ACC ATC TTG ACT GTC CAG CTC TTT GCA   221
Leu Xaa Ala Thr Gln Val Tyr Thr Ile Leu Thr Val Gln Leu Phe Ala
-25                      -20                      -15                      -10

TTC TTA AAC CTA CTG CCT GTA GAA GCA GAC ATT TTA GCA TAT AAC TTT   269
Phe Leu Asn Leu Leu Pro Val Glu Ala Asp Ile Leu Ala Tyr Asn Phe
                      -5                      1                      5

GAA AAT GCA TCT CAG ACA TTT GAT GAC CTC CCC GCA   305
Glu Asn Ala Ser Gln Thr Phe Asp Asp Leu Pro Ala
      10                      15

```

(2) INFORMATION FOR SEQ ID NO: 187:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 333 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 95..297
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 63..265
id N31364
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 32..95
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..64
id N31364
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 293..329
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 262..298
id N31364
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 2..211
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 10..219
id R06709
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 203..235
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 212..244
id R06709
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 96..329
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 64..297
id N75642
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 35..95
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 4..64
id N75642
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 96..266
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 99
region 104..274
id W92876
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 2..95
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 96
region 11..104
id W92876
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(252..329)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 183..260
id W92774
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(199..254)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 259..314
id W92774
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 106..216
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3
seq LWEKLTLLSPGIA/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 187:

```

AGCGATTGCG AATCCTCCGC TGAGGTGATT TGGATATCCC TAGAACGTTG AGGGCACGAG      60
TCGGGTCCTG AGACCAGGTC CTCAGCCAGC AGAGCCACGT TCCTT ATG AGC ACC GTG      117
                                     Met Ser Thr Val
                                     -35

GGT TTA TTT CAT TTT CCT ACA CCA CTG ACC CGA ATA TGC CCG GCG CCA      165
Gly Leu Phe His Phe Pro Thr Pro Leu Thr Arg Ile Cys Pro Ala Pro
      -30                      -25                      -20

TGG GGA CTC CGG CTT TGG GAG AAG CTG ACG TTG TTA TCC CCA GGA ATA      213
Trp Gly Leu Arg Leu Trp Glu Lys Leu Thr Leu Leu Ser Pro Gly Ile
      -15                      -10                      -5

GCT GTC ACT CCG GTC CAG ATG GCA GGC AAG AAG GAC TAC CCT GCA CTG      261
Ala Val Thr Pro Val Gln Met Ala Gly Lys Lys Asp Tyr Pro Ala Leu
      1                      5                      10                      15

CTT TCC TTG GAT GAG VAT GAA CTC GAA GAG CAG TTT GTG AAA GGA CAC      309
Leu Ser Leu Asp Glu Xaa Glu Leu Glu Glu Phe Val Lys Gly His
      20                      25                      30

GGT CCA GGG GGC CAG GCA ACG CGG      333
Gly Pro Gly Gly Gln Ala Thr Arg
      35

```

(2) INFORMATION FOR SEQ ID NO: 188:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 510 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 188..485
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95
region 171..468
id R76663
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 17..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..170
id R76663
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 183..367
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 169..353
id H67124
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 14..185
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..172
id H67124
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 214..430
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 81..297
id R53683
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 133..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93
region 1..95
id R53683
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 16..114
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq LLNFLGLWSWICK/KW

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 188:

```

ATTCCCTTCT GAGCA ATG GAG CTT ACC ATC TTT ATC CTG AGA CTG GCC ATT      51
      Met Glu Leu Thr Ile Phe Ile Leu Arg Leu Ala Ile
                -30                      -25

TAC ATC CTG ACA TTT CCC TTG TAC CTG CTG AAC TTT CTG GGC TTG TGG      99
Tyr Ile Leu Thr Phe Pro Leu Tyr Leu Leu Asn Phe Leu Gly Leu Trp
      -20                      -15                      -10

AGC TGG ATA TGC AAA AAA TGG TTC CCC TAC TTC TTG GTG AGG TTC ACT      147
Ser Trp Ile Cys Lys Lys Trp Phe Pro Tyr Phe Leu Val Arg Phe Thr
      -5                      1                      5                      10

GTG ATA TAC AAC GAA CAG ATG GCA AGC AAG AAG CGG GAG CTC TTC AGT      195
Val Ile Tyr Asn Glu Gln Met Ala Ser Lys Lys Arg Glu Leu Phe Ser
                15                      20                      25

AAC CTG CAG GAG TTT GCG GGC CCC TCC GGG AAA CTC TCC CTG CTG GAA      243
Asn Leu Gln Glu Phe Ala Gly Pro Ser Gly Lys Leu Ser Leu Leu Glu
                30                      35                      40

GTG GGC TGT GGC ACG GGG GCC AAC TTC AAG TTC TAC CCA CCT GGG TGC      291
Val Gly Cys Gly Thr Gly Ala Asn Phe Lys Phe Tyr Pro Pro Gly Cys
      45                      50                      55

AGG GTG ACC TGT ATT GAC CCC AAC CCC AAC TTT GAG AAG TTT TTG ATC      339
Arg Val Thr Cys Ile Asp Pro Asn Pro Asn Phe Glu Lys Phe Leu Ile
      60                      65                      70                      75

AAG AGC ATT GCA GAG AAC CGA CAC CTG CAG TTT GAG CGC TTT GTG GTA      387
Lys Ser Ile Ala Glu Asn Arg His Leu Gln Phe Glu Arg Phe Val Val
                80                      85                      90

GCT GCC GGG GAG AAC ATG CAC CAG GTG GCT GAT GGC TCT GTG GAT GTG      435
Ala Ala Gly Glu Asn Met His Gln Val Ala Asp Gly Ser Val Asp Val
                95                      100                      105

GTG GTC TGC ACC CTG GTG CTG TGC TCT GTG AAG AAC CAG GAG CGG ATT      483
Val Val Cys Thr Leu Val Leu Cys Ser Val Lys Asn Gln Glu Arg Ile
      110                      115                      120

CTC CGC GAG GTG TGC AGA GTG CTG AGA      510
Leu Arg Glu Val Cys Arg Val Leu Arg
      125                      130

```

(2) INFORMATION FOR SEQ ID NO: 189:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 155 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(2..122)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 9..129
id R16550
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..150
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..66
id N42850
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 9..119
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq LLLYLCCMINIHH/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 189:

```

ATTGAACT ATG AGT CTC CTG CAT GGC AAC AAA ATG TGT GTC ACC ATC AGG      50
  Met Ser Leu Leu His Gly Asn Lys Met Cys Val Thr Ile Arg
        -35                      -30                      -25

CCA ACA GGC CAG CCC TTG AAT GGG GAT TTA TTA CTG TTG TAT CTA TGT      98
Pro Thr Gly Gln Pro Leu Asn Gly Asp Leu Leu Leu Leu Tyr Leu Cys
        -20                      -15                      -10

TGC ATG ATA AAC ATT CAT CAC CTT CCT CCT GTA GTC CTG CCT CGT ACT      146
Cys Met Ile Asn Ile His His Leu Pro Pro Val Val Leu Pro Arg Thr
        -5                      1                      5

CCC CAA GGG
Pro Gln Gly
10

```

155

(2) INFORMATION FOR SEQ ID NO: 190:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

- (D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
(B) LOCATION: complement(90..123)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 1..34
id H04995
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: 122..172
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.9
seq ISYFIAFPNLSQA/EL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 190:

```
ATAATCATAT TTCAAATGA ATAGCAAAGA GGGTTATAAT CAAGTTTAT AAAAATTCCA    60
AATGTAATAA AGTTATATTT GTAACCTACA TATACTGCAA AAATGGTAGT GATTCAAATG   120
T ATG TCT TTC AAT ATA TCG TAT TTT ATT GCC TTT CCA AAT CTC TCC CAG   169
  Met Ser Phe Asn Ile Ser Tyr Phe Ile Ala Phe Pro Asn Leu Ser Gln
    -15                      -10                      -5

GCA GAA TTA ACA CAT CCC AGG TGC TCT TAC ACA GGG TTG AGT AGT TCC    217
Ala Glu Leu Thr His Pro Arg Cys Ser Tyr Thr Gly Leu Ser Ser Ser
   1             5             10             15

TGT GGA TTT CAG TTG AGT GAT ACC CCC CAC AGG                        250
Cys Gly Phe Gln Leu Ser Asp Thr Pro His Arg
          20             25
```

(2) INFORMATION FOR SEQ ID NO: 191:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 379 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: other
(B) LOCATION: 193..356
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 131..294

id AA148880
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 93..193
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 97
region 30..130
id AA148880
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 63..97
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 91
region 1..35
id AA148880
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 122..307
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 131..316
id H92506
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 9..97
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 92
region 1..89
id H92506
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 302..361
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.7
seq LTIILLPVHLLIT/IY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 191:

```
ACTTTTTGCG AGCTTTCCGA GTGCCAGGCG CCGCCCGGCT GCGAAGACGC GGTGGGCCGC   60
CCCTCCGATT GAAATCACAG AAGATATTCG KGKTCCTTCT TAAGAGAAAA AGAGGACATT  120
TGCGTACTTT ATTGTGCGCT TCCAAAGATT ACTAACTTTT ATCTGTATCA CTAAGATTGA  180
ACTGCCTTGG CTGTACTGCT ATTCTTACTG CTGCTTCTAT TATTGCCTTC TTCAGCACAA  240
TAAGGCTTTC AAAAGCCAAA GAATAACAAG AAATAAGCAC CATTTTAGAA GCCTTTCCAC  300
T ATG AAA CTT AAG CWA AAT GTG CTC ACC ATT ATT TTG CTG CCT GTC CAC  349
Met Lys Leu Lys Xaa Asn Val Leu Thr Ile Ile Leu Leu Pro Val His
```

-20

-15

-10

-5

TTG TTA ATA ACA ATA TAC AGT GCC CTT ATA
Leu Leu Ile Thr Ile Tyr Ser Ala Leu Ile
1 5

379

(2) INFORMATION FOR SEQ ID NO: 192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 176 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 25..173
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91
region 67..215
id H46464
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 73..173
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 44..144
id C17500
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 32..67
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94
region 1..36
id C17500
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(112..173)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 215..276
id AA143237
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(65..117)

(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 90
region 270..322
id AA143237
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(125..173)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 241..289
id W57931
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: complement(8..50)
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 95
region 368..410
id W57931
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 9..50
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.7
seq AALVTVLFTGVRR/LH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 192:

AGCTCAGC	ATG	GCT	GCT	TTA	GTG	ACT	GTT	CTC	TTC	ACA	GGT	GTC	CGG	AGG	50	
	Met	Ala	Ala	Leu	Val	Thr	Val	Leu	Phe	Thr	Gly	Val	Arg	Arg		
					-10										-5	
CTG	CAC	TGC	AGC	GCR	SCG	CTT	GGG	CGG	GCG	GCC	AGT	GGC	GRC	TAC	AGC	98
Leu	His	Cys	Ser	Ala	Xaa	Leu	Gly	Arg	Ala	Ala	Ser	Gly	Xaa	Tyr	Ser	
1				5					10					15		
AGG	AAC	TGG	CTG	CCA	ACC	CCT	CCG	GCT	ACG	GGC	CCC	TTA	CCG	AGC	TCC	146
Arg	Asn	Trp	Leu	Pro	Thr	Pro	Pro	Ala	Thr	Gly	Pro	Leu	Pro	Ser	Ser	
			20					25						30		
CAG	ACT	GGT	CAT	ATG	CGG	ATG	GCC	GCC	AGG							176
Gln	Thr	Gly	His	Met	Arg	Met	Ala	Ala	Arg							
			35					40								

(2) INFORMATION FOR SEQ ID NO: 193:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 236 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..236
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 91..242
id AA031580
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..87
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 10..94
id AA031580
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..236
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96
region 82..233
id W80981
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..87
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 1..85
id W80981
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 90..236
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97
region 89..235
id R69999
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..88
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100
region 3..88
id R69999
est

(ix) FEATURE:

- (A) NAME/KEY: other

(B) LOCATION: 90..236
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 89..235
 id R76832
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 3..88
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 100
 region 3..88
 id R76832
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 90..236
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 97
 region 89..235
 id R80120
 est

(ix) FEATURE:

(A) NAME/KEY: other
 (B) LOCATION: 3..88
 (C) IDENTIFICATION METHOD: blastn
 (D) OTHER INFORMATION: identity 100
 region 3..88
 id R80120
 est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 6..131
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 3.7
 seq ILMRDFSPSGIFG/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 193:

ATAAA ATG GCG TCA GTT GGT GAG TGT CCG GCC CCA GTA CCA GTG AAG GAC	50
Met Ala Ser Val Gly Glu Cys Pro Ala Pro Val Pro Val Lys Asp	
-40 -35 -30	
AAG AAA CTT CTG GAG GTC AAA CTG GGG GAG CTG CCA AGC TGG ATC TTG	98
Lys Lys Leu Leu Glu Val Lys Leu Gly Glu Leu Pro Ser Trp Ile Leu	
-25 -20 -15	
ATG CGG GAC TTC AGT CCT AGT GGC ATT TTC GGA GCG TTT CAA AGA GGT	146
Met Arg Asp Phe Ser Pro Ser Gly Ile Phe Gly Ala Phe Gln Arg Gly	
-10 -5 1 5	
TAC TAS CGG TAC TAC AAC VAG TAC ATC AAT GTG RAG AAG GGG AGC ATC	194
Tyr Xaa Arg Tyr Tyr Asn Xaa Tyr Ile Asn Val Xaa Lys Gly Ser Ile	
10 15 20	
TCG GGG ATT AMC ATG GTG CTG GCA TGC TAC GTG CTC TTT AGC	236

Ser Gly Ile Xaa Met Val Leu Ala Cys Tyr Val Leu Phe Ser
 25 30 35

(2) INFORMATION FOR SEQ ID NO: 194:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 155 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 61..155
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 98
region 4..98
id HUMPASP
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 77..155
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98
region 1..79
id W44779
est

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 81..125
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.9
seq ALLVLVTVALASA/HH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 194:

```

AGCAATCAGG AGAAGAAAGC AAAGAACACT CAGGATTATA AAAGCAGATG AGACCTACCC      60
ACTAGACCTG GTCAGACACA ATG TTG GCA CTC TTG GTT CTG GTG ACT GTG GCC      113
           Met Leu Ala Leu Leu Val Leu Val Thr Val Ala
           -15                -10                -5

CTG GCA TCT GCT CAT CAT GGT GGT GAG CAC TTT GAA GGC GCG      155
Leu Ala Ser Ala His His Gly Gly Glu His Phe Glu Gly Ala
           1                5                10

```

(2) INFORMATION FOR SEQ ID NO: 195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 214 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..211
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 98
region 1..165
id HSU16129
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..211
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 92
region 117..326
id RATGLUR4A
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 35..211
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 91
region 1..177
id GGGLUR4
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..211
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 93
region 1..165
id RATAMPASGD
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..211
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 93
region 1..165
id RATGLURD
vrt

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..211
- (C) IDENTIFICATION METHOD: fasta

(D) OTHER INFORMATION: identity 93
region 1..165
id S94371
vrt

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 61..211
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 100
region 93..243
id R93859
est

(ix) FEATURE:

(A) NAME/KEY: other
(B) LOCATION: 136..210
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION: identity 98
region 1..75
id R16185
est

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: 47..109
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 8.9
seq VLLFSGFWGLAMG/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 195:

```

AAGAGAGAAA GAGAGAGAGC GCGCGCCAGG GAGAGGAGAA AAGAAG ATG AGG ATT      55
                                     Met Arg Ile
                                     -20

ATT TCC AGA CAG ATT GTC TTG TTA TTT TCT GGA TTT TGG GGA CTC GCC      103
Ile Ser Arg Gln Ile Val Leu Leu Phe Ser Gly Phe Trp Gly Leu Ala
   -15          -10          -5

ATG GGA GCC TTT CCG AGC AGC GTG CAA ATA GGT GGT CTC TTC ATC CGA      151
Met Gly Ala Phe Pro Ser Ser Val Gln Ile Gly Gly Leu Phe Ile Arg
   1              5              10

AAC ACA GAT CAG GAA TAC ACT GCT TTT CGA TTA GCA ATT TTT CTT CAT      199
Asn Thr Asp Gln Glu Tyr Thr Ala Phe Arg Leu Ala Ile Phe Leu His
  15          20          25          30

AAC ACC AGC CCC GGG      214
Asn Thr Ser Pro Gly
          35

```

(2) INFORMATION FOR SEQ ID NO: 196:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -21..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 14.3
seq LLLCAVLLSLASA/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 196:

Met Arg Val Arg Ile Gly Leu Thr Leu Leu Leu Cys Ala Val Leu Leu
-20 -15 -10

Ser Leu Ala Ser Ala Ser Ser Asp Glu Glu Gly Asn Gly
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 197:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -28..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 11.1
seq GLLFLLLLMLLA/DP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 197:

Met Gly Leu His Leu Arg Pro Tyr Arg Val Gly Leu Leu Pro Asp Gly
-25 -20 -15

Leu Leu Phe Leu Leu Leu Leu Leu Met Leu Leu Ala Asp Pro Ala Leu
-10 -5 1

Pro Ala Ala Arg
5

(2) INFORMATION FOR SEQ ID NO: 198:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 134 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.6
seq LLLGAVSWPPASA/SG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 198:

```

Met Ala Thr Leu Ser Phe Val Phe Leu Leu Leu Gly Ala Val Ser Trp
  -20                      -15                      -10

Pro Pro Ala Ser Ala Ser Gly Gln Glu Phe Trp Pro Gly Gln Ser Ala
  -5                      1                      5                      10

Ala Asp Ile Leu Ser Gly Ala Ala Ser Arg Arg Arg Tyr Leu Leu Tyr
          15                      20                      25

Asp Val Asn Pro Pro Glu Gly Phe Asn Leu Arg Arg Asp Val Tyr Ile
          30                      35                      40

Arg Ile Ala Ser Leu Leu Lys Thr Leu Leu Lys Thr Glu Glu Trp Val
          45                      50                      55

Leu Xaa Leu Pro Pro Trp Gly Arg Leu Xaa Xaa Trp Gln Ser Xaa Asp
          60                      65                      70                      75

Ile His Gln Val Arg Ile Pro Trp Ser Glu Phe Phe Asp Leu Pro Ser
          80                      85                      90

Leu Asn Lys Asn Ile Pro Val Ile Glu Tyr Glu Gln Phe Ile Ala Glu
          95                      100                      105

Ser Gly Gly Pro Phe Ile
          110

```

(2) INFORMATION FOR SEQ ID NO: 199:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 59 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -54..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1
seq VLCLRGLVSLAFQ/GP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 199:

```

Met Phe Leu Phe Leu Ser Pro Ala Thr Pro Val Leu Pro Pro Ser Leu
      -50                      -45                      -40

Asp Ser Arg Asp Leu Leu Pro His Leu Phe Trp Gly Arg Ala Gly Ser
      -35                      -30                      -25

Ser Ser Ser Ser Pro Ala Leu Ser Pro Val Leu Cys Leu Arg Gly Leu
      -20                      -15                      -10

Val Ser Leu Ala Phe Gln Gly Pro His Pro Glu
      -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 200:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.8
seq LLWALLFMQSLWP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 200:

```

Met Trp Ala Met Glu Ser Gly His Leu Leu Trp Ala Leu Leu Phe Met
      -20                      -15                      -10

Gln Ser Leu Trp Pro Gln Leu Thr Asp Gly Ala Thr Arg Val Tyr Tyr
      -5                      1                      5                      10

Leu Gly Ile Arg Asp Val Gln Trp Asn Tyr Ala Pro Lys Gly Arg Asn
      15                      20                      25

```

Val Ile Thr Asn Gln Pro Leu Asp Ser Asp Ile Val Ala Ser Ser Phe
30 35 40

Leu Lys Ser Asp Asp Gly
45

(2) INFORMATION FOR SEQ ID NO: 201:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.7
seq ILGLCCVLATMA/NP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 201:

Met Thr His Tyr Arg Asn Ile Leu Gly Leu Leu Cys Cys Val Leu Ala
-15 -10 -5

Thr Met Ala Asn Pro Gly
1

(2) INFORMATION FOR SEQ ID NO: 202:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.4
seq LLLLLASLIERS/KT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 202:

Met Lys Leu Leu Leu Leu Ala Ser Leu Ile Glu Arg Ser Ser Lys
 -15 -10 -5 1

Thr Ser Cys Xaa Xaa Gln His Tyr Ser Ser Gln
 5 10

(2) INFORMATION FOR SEQ ID NO: 203:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -38..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.2
seq SFXLFLALCASFS/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 203:

Met Ala Arg Asn Gln Ala Leu Val Cys Leu Pro Ser Phe Gln Asn Ala
 -35 -30 -25

Phe Ile Pro Val Glu Asp Leu Pro Thr Ser Phe Xaa Leu Phe Leu Ala
 -20 -15 -10

Leu Cys Ala Ser Phe Ser Phe Phe Leu Xaa Leu Ser Leu Ser Leu Pro
 -5 1 5 10

Ser Phe Phe Phe

(2) INFORMATION FOR SEQ ID NO: 204:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide

(B) LOCATION: -35..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 8.2
seq SLLLLFYSFYVLA/VK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 204:

Met Pro Asn Glu Ser Trp Gln Ile Pro Cys Gly Lys Gln Glu Ala Glu
-35 -30 -25 -20
Thr Leu Phe Asn Phe Gln Ser Leu Leu Leu Leu Phe Tyr Ser Phe Tyr
-15 -10 -5
Val Leu Ala Val Lys Arg Gly Pro Gly
1 5

(2) INFORMATION FOR SEQ ID NO: 205:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -36..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 8
seq LVLLICLVSSYLP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 205:

Met Gln Thr Thr Phe Ile Asp Val Thr Val Asp Gln His Val Ala Lys
-35 -30 -25
Ser Asn Asp His Leu Ser Val Leu Val Leu Leu Ile Cys Leu Val Ser
-20 -15 -10 -5
Ser Tyr Leu Pro Gln Leu Pro
1

(2) INFORMATION FOR SEQ ID NO: 206:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -73..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 7.8
 seq YLPLLAGLGLTLA/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 206:

Met Ala Leu Gly Glu Glu Lys Ala Glu Ala Glu Ala Ser Xaa Asp Thr
 -70 -65 -60

Lys Ala Gln Ser Tyr Gly Arg Gly Ser Cys Arg Glu Arg Glu Leu Asp
 -55 -50 -45

Ile Pro Gly Pro Met Ser Gly Glu Gln Pro Pro Arg Leu Glu Ala Glu
 -40 -35 -30

Gly Gly Leu Ile Ser Pro Val Trp Gly Ala Glu Xaa Tyr Leu Pro Leu
 -25 -20 -15 -10

Leu Ala Gly Leu Gly Leu Thr Leu Ala Ala Pro Leu Glu Pro Thr Thr
 -5 1 5

(2) INFORMATION FOR SEQ ID NO: 207:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -21..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 7.7
 seq LLCISPFVPFTSG/NK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 207:

Met Thr Ser Leu Tyr Leu Lys His Leu Leu Cys Ile Ser Pro Phe Val
 -20 -15 -10

Pro Phe Thr Ser Gly Asn Lys Leu Tyr Tyr Thr Met Ile Tyr Trp Leu
 -5 1 5 10

Phe Lys Thr Val Leu Asn Met His Gly
 15 20

(2) INFORMATION FOR SEQ ID NO: 208:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -47...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.6
seq CLATLTLFHTSFS/FQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 208:

Met Thr Asp Ser Pro Asn Ala His Gly Leu Ala Leu Thr Thr Lys Trp
 -45 -40 -35

Met Met Pro Ala Val Ser Leu Asn Leu Thr Tyr Tyr Leu Pro Ser Trp
 -30 -25 -20

Tyr Leu Cys Leu Ala Thr Leu Thr Leu Phe His Thr Ser Phe Ser Phe
 -15 -10 -5 1

Gln Ala Ser Glu Ser Val Lys Ala Ile Thr
 5 10

(2) INFORMATION FOR SEQ ID NO: 209:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -48...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5

seq FVILLFIFTVVS/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 209:

```

Met Ala Ser Ser His Trp Asn Glu Thr Thr Thr Ser Val Tyr Gln Tyr
      -45                -40                -35

Leu Gly Phe Gln Val Gln Lys Ile Tyr Pro Phe His Asp Asn Trp Asn
      -30                -25                -20

Thr Ala Cys Phe Val Ile Leu Leu Leu Phe Ile Phe Thr Val Val Ser
      -15                -10                -5

Leu Val Val Leu Ala Phe Leu Tyr Glu Val Leu Xaa Xaa Cys Cys Cys
  1              5              10              15

Val Lys Asn Lys Thr Val Lys Asp Leu Lys Ser Glu Pro Lys
      20              25              30

```

(2) INFORMATION FOR SEQ ID NO: 210:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 44 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -25..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.4
seq IFLLNMWVACLLS/GE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 210:

```

Met Ser Leu Leu Phe Val Phe Cys Leu Glu Cys Ser Ile Phe Leu Leu
-25                -20                -15                -10

Asn Met Trp Val Ala Cys Leu Leu Ser Gly Glu Ile Pro His Ser Ser
      -5              1              5

Trp Xaa Xaa Lys Leu Ile Gly Thr Leu Pro Thr Ser
      10              15

```

(2) INFORMATION FOR SEQ ID NO: 211:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids

(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -18..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 7.4
seq VLLLLPLVAFITL/KF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 211:

```
Met Phe Val Val Thr Val Leu Leu Leu Leu Pro Leu Val Ala Phe Ile
    -15                      -10                      -5

Thr Leu Lys Phe Cys Asn Leu Ile Asn Phe Pro Thr Xaa Arg His Gly
    1                      5                      10
```

(2) INFORMATION FOR SEQ ID NO: 212:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 67 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -22..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 7.2
seq IIYALQFLFLVFA/PS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 212:

```
Met Asn Arg Ser Cys Arg Asn Thr Gly Ile Ile Tyr Ala Leu Gln Phe
    -20                      -15                      -10

Leu Phe Leu Val Phe Ala Pro Ser Ser Leu Gly Tyr Phe Glu Trp Ile
    -5                      1                      5                      10

Val Ala Ile Asn Gln Asp Leu Val Leu Phe Val Phe Cys Leu Ser Phe
    15                      20                      25

Ser Leu Arg Ile Ser Ile Ile Gln Gly Lys Arg Lys Ala Ala Phe Pro
    30                      35                      40
```

Thr Pro Pro
45

(2) INFORMATION FOR SEQ ID NO: 213:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -35..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.2
seq LSLLLAWVTLTHL/LS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 213:

Met	Thr	Gln	Thr	Thr	Trp	Gly	Ala	Pro	Thr	Arg	Ala	Ser	Asn	His	Pro
-35					-30					-25					-20
Leu	Pro	Ala	Trp	Leu	Thr	Leu	Ser	Leu	Leu	Leu	Ala	Trp	Val	Thr	Leu
				-15					-10					-5	
Thr	His	Leu	Leu	Ser	Val	Leu	Thr	His	Pro	Thr	Leu	Leu			
		1					5					10			

(2) INFORMATION FOR SEQ ID NO: 214:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1
seq XXAVLCVCAAWC/SQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 214:

```

Met Leu Lys Xaa Xaa Ala Val Leu Cys Val Cys Ala Ala Ala Trp Cys
  -15                -10                -5

Ser Gln Ser Leu Ala Ala Ala Ala Val Ala Ala Xaa Gly Gly Arg
  1          5          10          15

Ser Asp Gly Gly Asn Phe Leu Asp Asp Lys Gln Trp Leu Thr Thr Ile
          20                25                30

Ser Gln Tyr Asp Lys Glu Val Gly Gln Trp Asn Lys Phe Arg Asp Asp
          35                40                45

Asp Tyr Phe Arg Thr Trp Ser Pro Gly Lys Pro Phe Asp Gln Ala Leu
          50                55                60

Asp Xaa Ala Asn Gly
          65

```

(2) INFORMATION FOR SEQ ID NO: 215:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9
seq LHLGSSISPASA/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 215:

```

Met Ile Ser Ala His Cys Asn Leu His Leu Leu Gly Ser Ser Ile Ser
-20                -15                -10                -5

Pro Ala Ser Ala Ser Leu
          1

```

(2) INFORMATION FOR SEQ ID NO: 216:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -28..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.9
seq FLPFLLSLPLDQT/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 216:

```

Met Xaa Xaa Lys Ala Cys Arg Thr Leu Ala Trp Leu Pro Xaa Pro Phe
      -25                      -20                      -15

Leu Pro Phe Leu Leu Ser Leu Pro Leu Asp Gln Thr Leu Pro Arg Gln
      -10                      -5                      1

Gly Pro Gly Gln Ser Leu Ser Phe Pro Glu Asn Tyr Gln Thr Leu Pro
  5                      10                      15                      20

Lys Ser Thr Arg His Pro Gly
      25

```

(2) INFORMATION FOR SEQ ID NO: 217:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -19..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.8
seq GLLLVFLPHQRG/GQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 217:

```

Met Ala Val Lys Arg Leu Gly Leu Leu Val Phe Leu Pro His Pro
      -15                      -10                      -5

Gln Arg Gly Gly Gln Glu Arg Ser Ala His Thr Pro Arg Gln His Pro
      1                      5                      10

Ala Arg Pro Thr Ser Leu Ser Gln Gly Glu Arg Pro Gly Arg Gly Gly
  15                      20                      25

```

Gly Trp Gly Asn Gly Arg Asp Ala His Gln
30 35

(2) INFORMATION FOR SEQ ID NO: 218:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 119 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -83..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8
seq LLMILTFPFKILS/DA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 218:

Met Gln Ala Val Asp Asn Leu Thr Ser Ala Pro Gly Asn Thr Ser Leu
-80 -75 -70

Cys Thr Arg Asp Tyr Lys Ile Thr Gln Val Leu Phe Pro Leu Leu Tyr
-65 -60 -55

Thr Val Leu Phe Phe Val Gly Leu Ile Thr Asn Gly Leu Ala Met Arg
-50 -45 -40

Ile Phe Phe Gln Ile Arg Ser Lys Ser Asn Phe Ile Ile Phe Leu Lys
-35 -30 -25 -20

Asn Thr Val Ile Ser Asp Leu Leu Met Ile Leu Thr Phe Pro Phe Lys
-15 -10 -5

Ile Leu Ser Asp Ala Lys Leu Gly Thr Gly Pro Leu Arg Thr Phe Val
1 5 10

Cys Gln Val Thr Ser Val Ile Phe Tyr Xaa Xaa Met Tyr Ile Ser Ile
15 20 25

Ser Phe Leu Gly Leu Ile Thr
30 35

(2) INFORMATION FOR SEQ ID NO: 219:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -36..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 6.8
seq LWFFLPSLXCPEC/CP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 219:

```

Met Trp Thr Leu Pro Ser Leu Ser Ala Ser Phe Gln Pro Phe Leu Gly
-35                      -30                      -25

Ser Leu Arg Pro Ser His Ile Leu Trp Phe Phe Leu Pro Ser Leu Xaa
-20                      -15                      -10                      -5

Cys Pro Glu Cys Cys Pro Pro Asp Pro Gly Ser Pro Ala Ser Arg Asp
                      1                      5                      10

Pro Asn Val Ala Cys Glu Arg
                      15

```

(2) INFORMATION FOR SEQ ID NO: 220:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 84 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -22..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 6.7
seq LLLFQPSSHSATG/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 220:

```

Met Ser Leu Thr Asp Val Pro Met Ser Leu Leu Leu Phe Gln Pro Ser
-20                      -15                      -10

Ser His Ser Ala Thr Gly Ser Ser Ile Lys Ile Ile Ile Leu Asn Tyr
-5                      1                      5                      10

```


Ile Ile Leu Gln Phe Lys Thr Leu Gln Thr Leu Pro Asn Ala Leu Arg
 15 20 25
 Ile His Ile Lys Val Phe His Ile Tyr Cys Ser Phe Val Ser Arg Phe
 30 35 40
 His Tyr Tyr Lys Asn Thr Ala Thr Val Phe Phe Arg Ser Val Leu Lys
 45 50 55
 Arg Arg Met Gly
 60

(2) INFORMATION FOR SEQ ID NO: 221:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -17..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.6
seq LAFLLVSLYWSHM/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 221:

Met Asp Trp Ser Leu Ala Phe Leu Leu Val Ser Leu Tyr Trp Ser His
 -15 -10 -5
 Met His Pro Cys Tyr Trp Ser Trp Pro Cys Ser Cys Gly Phe Val Asp
 1 5 10 15
 Ser Pro Cys Ile Cys Thr Ala Ser Thr Arg Cys Cys Cys Ser Ser Leu
 20 25 30
 Cys Cys Leu Trp Arg Leu Ala Leu Trp Asp Trp Thr Ser Asn Gly Ser
 35 40 45
 Arg Ser Gly Ile Ala Cys Val Cys
 50 55

(2) INFORMATION FOR SEQ ID NO: 222:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5
seq LLILFFMVGRIP/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 222:

Met Tyr Leu Leu Ile Leu Phe Phe Met Val Gly Arg Ile Ile Pro Ser
-15 -10 -5 1

Pro His Arg

(2) INFORMATION FOR SEQ ID NO: 223:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -48..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5
seq LLVSCCLLFHQ/IA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 223:

Met Asn Lys Pro Pro Trp Glu Glu Ser Trp Gly Gln Asn Gln Leu Ser
-45 -40 -35

Gly Glu Pro Ala Thr Trp Ser Leu Cys Ile Ser Pro Leu Pro Gly Arg
-30 -25 -20

Glu Pro Ser Leu Leu Val Val Ser Cys Cys Leu Leu Phe His Gln Ala
-15 -10 -5

Ile His Asn Lys Leu Leu Trp Arg
1 5

(2) INFORMATION FOR SEQ ID NO: 224:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -24..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.5
seq FLILLSIDSLVSG/FL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 224:

Met Leu Ile Leu Glu Leu Thr Met Met Leu Ser Phe Leu Ile Leu Leu
-20 -15 -10

Ser Ile Asp Ser Leu Val Ser Gly Phe Leu Ser Lys Arg Lys Gly Leu
-5 1 5

Arg Val Cys Asp Gly Ser Arg Ser Gly
10 15

(2) INFORMATION FOR SEQ ID NO: 225:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -52..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.4
seq CLLGAAWASRLRT/OP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 225:

Met Lys Leu Gln Arg Ser Arg Ala Phe Arg Ile Glu Cys Ser Ala Ile
-50 -45 -40

Leu Arg Arg Ala Glu Arg Leu Val Trp Asn Asp Val Cys Ser Glu Ser
-35 -30 -25
Gln Ser Gln Ser Arg Asp Ser Cys Leu Leu Gly Ala Ala Trp Ala Ser
-20 -15 -10 -5
Arg Leu Arg Thr Gln Pro His Pro
1

(2) INFORMATION FOR SEQ ID NO: 226:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq FTLCVFTLPFLCA/CL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 226:

Met Val Ile Phe Thr Leu Cys Val Phe Thr Leu Pro Phe Leu Cys Ala
-15 -10 -5
Cys Leu Pro Arg
1

(2) INFORMATION FOR SEQ ID NO: 227:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq VLVVGTWSSQGQA/NS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 227:

Met Trp Gly Ala Leu Pro Val Leu Val Val Gly Thr Trp Ser Ser Gln
 -15 -10 -5
 Gly Gln Ala Asn Ser Cys Ala Gly Arg Gly Met Gly Pro Asp Val Cys
 1 5 10
 Gly Ala
 15

(2) INFORMATION FOR SEQ ID NO: 228:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq LVCGFLQISLSLA/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 228:

Met Thr Arg Leu Val Cys Gly Phe Leu Gln Ile Ser Leu Ser Leu Ala
 -15 -10 -5
 Ser Leu Phe Leu Thr Ile Pro Leu Met Trp Tyr Met Gln Ser Lys Trp
 1 5 10 15
 Trp Arg Gly

(2) INFORMATION FOR SEQ ID NO: 229:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -15..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.6
seq FLLPLLLHLLTFH/GR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 229:

Met Asn Phe Leu Leu Pro Leu Leu Leu His His Leu Thr Phe His Gly
-15 -10 -5 1

Arg Pro Leu Lys
5

(2) INFORMATION FOR SEQ ID NO: 230:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 81 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -58..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.5
seq LIIFICXTASISA/YM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 230:

Met Leu Ser Ala Arg Asp Arg Arg Asp Arg His Pro Glu Glu Gly Val
-55 -50 -45

Val Ala Glu Leu Gln Gly Phe Ala Val Asp Lys Ala Phe Leu Thr Ser
-40 -35 -30

His Lys Gly Ile Leu Leu Glu Thr Glu Leu Ala Leu Thr Leu Ile Ile
-25 -20 -15

Phe Ile Cys Xaa Thr Ala Ser Ile Ser Ala Tyr Met Ala Ala Ala Leu
-10 -5 1 5

Leu Glu Phe Phe Ile Thr Leu Ala Phe Leu Phe Leu Tyr Ala Thr Pro
10 15 20

Ala

(2) INFORMATION FOR SEQ ID NO: 231:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -13..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5
seq MLTMSVTLSPQRS/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 231:

Met Leu Thr Met Ser Val Thr Leu Ser Pro Leu Arg Ser Gln Asp Leu
 -10 -5 1
Asp Pro Met Ala Thr Asp Ala Ser Pro Met Ala Ile Asn Met Thr Pro
 5 10 15
Thr Val Glu Gln Gly Leu
 20 25

(2) INFORMATION FOR SEQ ID NO: 232:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -46..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq FLLISSVRPISQ/TF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 232:

Met Phe Xaa Pro Val Ala Leu Ile Phe Pro Ile Ser Val Ser Asp Pro


```

-45          -40          -35
Thr Ile His Pro Ile Thr Gln Ala Gln Asn Leu Glu Ser Xaa Leu Gln
-30          -25          -20          -15
Ser Phe Phe Leu Leu Ile Ser Ser Val Arg Pro Ile Ser Gln Thr Phe
          -10          -5          1
Lys Ile Asp Leu Ser Pro Ser Val Arg Ala Thr Gly
          5          10

```

(2) INFORMATION FOR SEQ ID NO: 233:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -31..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq WCAVLRSLAASS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 233:

```

Met Leu Leu Phe Phe Pro Phe Phe Gly Glu Thr Val Ser Leu His His
-30          -25          -20
Pro Cys Trp Cys Ala Val Leu Arg Ser Trp Leu Ala Ala Ser Ser Ala
-15          -10          -5          1
Pro Arg

```

(2) INFORMATION FOR SEQ ID NO: 234:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide

(B) LOCATION: -35..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.4
 seq LVVVCYLSWRVSS/RS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 234:

```

Met Pro Leu Lys Asn Leu Phe Ser Val Gly Leu Trp Asp Pro Tyr Asn
-35                               -30           -25           -20

Leu Leu Lys Lys His Val Leu Val Val Val Cys Tyr Leu Ser Trp Arg
                               -15           -10           -5

Val Ser Ser Arg Ser Trp Thr Leu Leu Ile Thr Pro Val Thr Leu His
                               1             5             10

Ala Ser Leu Ser Thr Gln Ala Arg
15                               20
  
```

(2) INFORMATION FOR SEQ ID NO: 235:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -19..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.4
 seq LSHLLPSLRQVIQ/EP.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 235:

```

Met Ala Met Ala Gln Lys Leu Ser His Leu Leu Pro Ser Leu Arg Gln
                               -15           -10           -5

Val Ile Gln Glu Pro Gln Leu Ser Leu Gln Pro Glu Xaa Val Phe Thr
                               1             5             10

Val Asp Arg Ala Glu Val Pro Pro Leu Phe Trp Lys Pro Tyr Ile Tyr
15                               20           25

Ala Gly Xaa Arg Pro Leu His Gln Thr Trp Arg Phe Tyr Phe Arg Thr
30                               35           40           45

Leu Phe Gln Gln His Asn
                               50
  
```

(2) INFORMATION FOR SEQ ID NO: 236:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -28..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq LALVALAPHSVQK/SX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 236:

Met Ile Ala Pro Thr Leu Lys Gly Thr Pro Ser Ser Ser Ala Pro Leu
-25 -20 -15

Ala Leu Val Ala Leu Ala Pro His Ser Val Gln Lys Ser Xaa Xaa Phe
-10 -5 1

Pro Pro Asn Leu Leu Thr Ser Pro Pro Ser Val Ala Xaa Ala Glu Ser
5 10 15 20

Gly Ser Val Ile Thr Leu Ser Ala Xaa Ile Ala Pro Ser Glu Pro Lys
25 30 35

Thr Asn Leu Asn Lys Val Pro Ser Glu Val
40 45

(2) INFORMATION FOR SEQ ID NO: 237:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -37..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.3
seq LVESLCLVFNLLS/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 237:

```
Met Cys Leu Phe Pro Val Ser Pro Cys Pro Ala Tyr Ser Phe Ser Ser
   -35                -30                -25
Glu Xaa Xaa Gly Ala Val Leu Leu Leu Val Glu Ser Leu Cys Leu Val
   -20                -15                -10
Phe Asn Leu Leu Ser Leu Pro Pro Arg
   -5                  1
```

(2) INFORMATION FOR SEQ ID NO: 238:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 91 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -15..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.3
seq IAVLFCFLLLIIF/QT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 238:

```
Met Lys Ile Ala Val Leu Phe Cys Phe Leu Leu Ile Ile Phe Gln
-15                -10                -5                1
Thr Asp Phe Gly Lys Asn Glu Glu Ile Pro Arg Lys Gln Arg Arg Lys
      5                10                15
Ile Tyr His Arg Arg Leu Arg Lys Ser Ser Thr Ser His Lys His Arg
      20                25                30
Ser Asn Arg Gln Leu Gly Ile Pro Gln Thr Thr Val Phe Thr Pro Val
      35                40                45
Ala Arg Leu Pro Ile Val Asn Phe Asp Tyr Ser Met Glu Glu Lys Phe
      50                55                60                65
Glu Ser Phe Ser Ser Phe Pro Gly Val Glu Ser
      70                75
```

(2) INFORMATION FOR SEQ ID NO: 239:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -27..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq VGAVLLSSLPISP/QY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 239:

Met Cys Ser Pro Arg Ser Pro Leu Asn Leu Ser Leu Val Pro Val Gly
-25 -20 -15

Ala Val Leu Leu Ser Ser Leu Pro Ile Ser Pro Gln Tyr Gly
-10 -5 1

(2) INFORMATION FOR SEQ ID NO: 240:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -48..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq EVVTLPLTSHCLA/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 240:

Met Gly Leu His Ile Ser Leu Ile Lys Phe Leu Leu Ala Asn Gly Pro
-45 -40 -35

His Ile Pro Ser His Gln Arg Pro Phe Glu Pro Lys Gly Glu Lys Ser
-30 -25 -20

Cys Arg Ile Glu Val Val Thr Leu Pro Leu Thr Ser His Cys Leu Ala
 -15 -10 -5

Gln Val Ala Ser Ser Asp Leu Ile His Arg Met Arg Thr Ile Thr Gly
 1 5 10 15

Thr Ser Ser His Gly
 20

(2) INFORMATION FOR SEQ ID NO: 241:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq LLTLVVFVASSMQ/IY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 241:

Met Lys Thr Thr Tyr Val Ile Phe Met Gln Ser Lys Ala Leu Leu Thr
 -25 -20 -15

Leu Tyr Val Phe Val Ala Ser Ser Met Gln Ile Tyr Val Leu His Ile
 -10 -5 1 5

Ser Asn-Tyr Pro Thr Asp Glu His Phe Pro Ile Ile Lys His Phe Tyr
 10 15 20

Phe Thr Phe Lys Ile His Phe Ser Lys Ile Ile Tyr Val Gln Tyr Ser
 25 30 35

(2) INFORMATION FOR SEQ ID NO: 242:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq ALVFLIFLRFINI/SE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 242:

Met Asn Ala Leu Val Phe Leu Ile Phe Leu Arg Phe Ile Asn Ile Ser
-15 -10 -5 1

Glu Val Thr Thr Lys Cys Gln Ala Gly
 5 10

(2) INFORMATION FOR SEQ ID NO: 243:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq WGFLLTGHSLSHS/SK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 243:

Met Gln Leu Gly Pro Leu His Thr Val Ser Thr Pro Phe Phe Phe Cys
 -25 -20 -15

Trp Gly Phe Leu Leu Thr Gly His Ser Leu Ser His Ser Ser Lys Ser
 -10 -5 1

Cys His Leu
 5

(2) INFORMATION FOR SEQ ID NO: 244:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 62 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -57..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq VGSVCCCVGPLRG/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 244:

Met Gly Arg Gly Trp Glu Arg Thr Val Cys Ser Leu Gly Trp Arg Gly
-55 -50 -45
Gly Pro Asp Pro Leu Ser Trp Ala Thr Cys Trp Ser Gly Ala Arg Ser
-40 -35 -30
Arg His Thr Arg Val Ser Ser Ile Val Asn Gly Tyr Val Gly Ser Val
-25 -20 -15 -10
Cys Cys Cys Val Gly Pro Leu Arg Gly Leu Val Trp Ala Pro
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 245:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5
seq HLFVTWSSQRALS/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 245:

Met Pro Glu Ala Val Glu Gln Ser Ala His Leu Phe Val Thr Trp Ser
-20 -15 -10
Ser Gln Arg Ala Leu Ser His Pro Ala Pro Leu
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 246:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 49 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -24..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5
seq CWLIALSVPLVFW/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 246:

Met Pro Gly Thr His Thr Phe Thr Phe Lys Ser Cys Trp Leu Ile Ala
 -20 -15 -10

Leu Ser Val Pro Leu Val Phe Trp Val Thr Phe Trp Pro Cys Asn Phe
 -5 1 5

Tyr Pro Ser Leu Asp Phe Cys Met Leu Thr Lys Xaa Lys Ser Ile Phe
 10 15 20

Ile
25

(2) INFORMATION FOR SEQ ID NO: 247:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq LFCLIGLDLLCQV/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 247:

Met Leu Leu Leu Thr Phe Lys Trp Phe Leu Phe Cys Leu Ile Gly Leu
 -20 -15 -10

Asp Leu Leu Cys Gln Val Phe Ser Pro Tyr Phe Leu Ser Glu Lys Val
 -5 1 5 10

Ala Asp Leu Leu Phe Tyr Met Ser Leu Phe Phe
 15 20

(2) INFORMATION FOR SEQ ID NO: 248:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -52..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq VPNLHLLLPLTTP/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 248:

Met Ala Thr Thr Gly Arg Arg Gln Ala Glu Pro Pro Pro Val Arg Pro
 -50 -45 -40

Ala His Ser Arg Pro Pro Pro Arg Val Pro Gly Ser Ser Ser Leu Gly
 -35 -30 -25

Leu Ala Gly Leu Met Ser Pro Val Pro Asn Leu His Leu Leu Leu Pro
 -20 -15 -10 -5

Leu Thr Thr Pro Gln Pro Arg
 1

(2) INFORMATION FOR SEQ ID NO: 249:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9
seq FIYLQAHFTLCSG/WS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 249:

Met Val Pro Phe Ile Tyr Leu Gln Ala His Phe Thr Leu Cys Ser Gly
-15 -10 -5

Trp Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr
1 5 10 15

Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile
20 25 30

Pro His Gly Pro Lys
35

(2) INFORMATION FOR SEQ ID NO: 250:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8
seq FFSFLLTINLVSL/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 250:

Met Phe Phe Ser Phe Leu Leu Thr Ile Asn Leu Val Ser Leu Gln Val
-10 -5 1

Val Ile Leu Asn Arg Val Tyr Leu Asn Gln Pro Asp Ala Arg
5 10 15

(2) INFORMATION FOR SEQ ID NO: 251:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -21..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8
seq NWGLLCFASECTT/DR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 251:

Met Trp Pro Gly Arg Glu Cys Lys Asn Trp Gly Leu Leu Cys Phe Ala
-20 -15 -10

Ser Glu Cys Thr Thr Asp Arg
-5 1

(2) INFORMATION FOR SEQ ID NO: 252:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 42 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(D) DEVELOPMENTAL STAGE: Fetal

(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -37..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8
seq CLLLTLRQPPTHSGI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 252:

Met Leu Thr Pro Phe Ser Leu Glu Glu Lys Leu Leu Glu Cys His Tyr
-35 -30 -25

Val Leu Ala Lys Leu Ala Gly Ala Cys Leu Leu Leu Thr Leu Arg Gln
-20 -15 -10

Pro Pro Thr His Ser Gly Ile Pro His Gly
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 253:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -40..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7
seq IVFVGLIFLKSSA/HR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 253:

Met Lys Arg Ile Lys Ser Met Met Gly Lys Val Glu His Ile Lys Ile
-40 -35 -30 -25

Lys Gly Glu Lys Gln Arg Ser Arg His Val Lys Ile Val Phe Val Gly
-20 -15 -10

Leu Ile Phe Leu Lys Ser Ser Ala His Arg
-5 1

(2) INFORMATION FOR SEQ ID NO: 254:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -17..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7
seq LCLFCKICPFTHG/VA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 254:

Met Gln Ser Ala Leu Cys Leu Phe Cys Lys Ile Cys Pro Phe Thr His
-15 -10 -5

Gly Val Ala Thr Pro Ala Trp Glu Leu Ser Ser Lys Arg Lys Ala Ser

1

5

10

15

His Pro Pro Arg

(2) INFORMATION FOR SEQ ID NO: 255:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 92 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -69..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7
seq SLLCLAFLLGKFL/HM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 255:

```

Met Met His Asn Ile Ile Val Lys Glu Leu Ile Val Thr Phe Phe Leu
      -65                      -60                      -55

Gly Ile Thr Val Val Gln Met Leu Ile Ser Val Thr Gly Leu Lys Gly
      -50                      -45                      -40

Val Glu Ala Gln Asn Gly Ser Glu Ser Glu Val Phe Val Gly Lys Tyr
      -35                      -30                      -25

Glu Thr Leu Val Phe Tyr Trp Pro Ser Leu Leu Cys Leu Ala Phe Leu
      -20                      -15                      -10

Leu Gly Arg Phe Leu His Met Phe Val Lys Ala Leu Arg Val His Leu
      -5                      1                      5                      10

Gly Trp Glu Leu Gln Val Glu Glu Lys Ser Val Leu
      15                      20

```

(2) INFORMATION FOR SEQ ID NO: 256:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 102 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -27..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6
seq RLLCSRLCQQLRS/KR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 256:

```

Met Ser Asn Cys Leu Gln Asn Phe Leu Lys Ile Thr Ser Thr Arg Leu
  -25                      -20                      -15

Leu Cys Ser Arg Leu Cys Gln Gln Leu Arg Ser Lys Arg Lys Phe Phe
  -10                      -5                      1                      5

Gly Thr Val Pro Ile Ser Arg Leu His Arg Arg Val Val Ile Thr Gly
          10                      15                      20

Ile Gly Leu Val Thr Pro Leu Gly Val Gly Thr His Leu Val Trp Asp
          25                      30                      35

Arg Leu Ile Gly Gly Glu Ser Gly Ile Val Ser Leu Val Gly Glu Glu
          40                      45                      50

Tyr Lys Ser Ile Pro Cys Ser Val Ala Ala Tyr Val Pro Arg Gly Ser
          55                      60                      65

Asp Glu Gly Gln Ser Gly
  70                      75

```

(2) INFORMATION FOR SEQ ID NO: 257:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6
seq XGLFLRTTAAARA/CR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 257:

```

Met Ser Gly Xaa Gly Leu Phe Leu Arg Thr Thr Ala Ala Ala Arg Ala
  -15                      -10                      -5

Cys Arg Gly Leu Val Val Ser Thr Ala Asn Arg Arg Leu Leu Arg Thr

```

1 5 10 15
Ser Pro Pro Val Arg Ala Phe Ala Lys Glu Leu Phe Leu Gly Lys Ile
 20 25 30
Xaa Lys Val Thr Arg Ala Leu Gly
 35 40

(2) INFORMATION FOR SEQ ID NO: 258:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -25..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq LTWLHLLLSHLKS/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 258:

Met Asn Pro Leu His Lys His Cys Ala Ala Gly Pro Leu Thr Trp Leu
-25 -20 -15 -10
His Leu Leu Leu Ser His Leu Lys Ser Ser Leu Val Met Pro Pro Lys
 -5 1 5

(2) INFORMATION FOR SEQ ID NO: 259:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5

seq FAVLRVLHLPALT/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 259:

Met Pro Lys Asp Lys Arg Gly Ala Arg His Asn Ser Pro His Phe Ser
 -25 -20 -15

Phe Ala Val Leu Arg Val Leu His Leu Pro Ala Leu Thr Ala Pro Leu
 -10 -5 1

Trp Leu Ala Pro Phe Ser Thr Leu Pro Arg
 5 10

(2) INFORMATION FOR SEQ ID NO: 260:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -39..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq LCVSRQLLTGART/LF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 260:

Met Thr Ile His Val Leu Arg Lys Cys Cys Gln Met Gly Arg Leu Asn
 -35 -30 -25

Asn Glu Trp Leu Pro Gly Leu Val Ile Pro Leu Cys Val Ser Arg Gln
 -20 -15 -10

Leu Leu Thr Gly Ala Arg Thr Leu Phe Gln Leu Gln Asn Gly Pro Ala
 -5 1 5

(2) INFORMATION FOR SEQ ID NO: 261:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq IAALLGLLQLRFK/AE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 261:

```

Met Gln Ala Ala Ser Phe Gly Arg Gly Arg Asn Gly Leu Asp Asn Trp
-30                -25                -20                -15

Gly Ile Ala Ala Leu Leu Gly Leu Leu Gln Leu Arg Phe Lys Ala Glu
                -10                -5                1

```

(2) INFORMATION FOR SEQ ID NO: 262:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5
seq ENLLLCCHRCTNC/QR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 262:

```

Met Ser Pro Ser Leu Gly Asp Arg Cys Ser Ser Trp Leu His Leu Val
                -40                -35                -30

Ser His Leu Glu Ser Ile Ser Gly Pro Leu Leu Asn Ile Pro Glu Asn
                -25                -20                -15

Leu Leu Leu Cys Cys His Arg Cys Thr Asn Cys Gln Arg His His Phe
-10                -5                1                5

Cys Ser Val Trp

```

(2) INFORMATION FOR SEQ ID NO: 263:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -24..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.4
seq YFLLPCLINLAIG/VK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 263:

```
Met Ser Gly Ala Glu Pro Thr Thr Phe Ile Arg Tyr Phe Leu Leu Pro
          -20                -15                -10
Cys Leu Ile Asn Leu Ala Ile Gly Val Lys Trp Lys Thr Ala Trp Lys
          -5                  1                  5
Arg Gly Glu Arg Gln Leu Asn Asn Thr Val Phe Phe Phe Phe Phe
    10                15                20
```

(2) INFORMATION FOR SEQ ID NO: 264:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -22..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.4
seq QLLESFLLSTIPT/SY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 264:

```
Met Val Tyr Asp Tyr Phe Ile Ser Gln Gln Leu Leu Phe Ser Phe Leu
          -20                -15                -10
Leu Ser Thr Ile Pro Thr Ser Tyr His Leu Ser Leu Thr Cys Gln Arg
    -5                  1                  5                10
```

(2) INFORMATION FOR SEQ ID NO: 265:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3
seq LFLCSCSLSLNQL/LT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 265:

Met Leu Phe Leu Cys Ser Cys Ser Leu Ser Leu Asn Gln Leu Leu Thr
 -10 -5 1

Tyr Ile Phe Val Val Pro Pro Trp
 5 10

(2) INFORMATION FOR SEQ ID NO: 266:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq FLMVLLFRSNKWT/GK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 266:

Met Phe Phe Leu Met Val Leu Leu Phe Arg Ser Asn Lys Trp Thr Gly
-15 -10 -5 1

Lys Val Tyr Gly Ala Leu
 5

(2) INFORMATION FOR SEQ ID NO: 267:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq FLSHVTTSCLASS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 267:

```

Met Leu Pro Leu Gln Gly Leu Cys Thr Cys Tyr Phe Leu His Leu Glu
      -25                      -20                      -15

Phe Leu Ser His Val Thr Thr Ser Leu Ala Ser Ser Ser Ala Pro Ser
      -10                      -5                      1

Pro Lys Pro Ser Val Thr Leu Ser Ser
      5                      10

```

(2) INFORMATION FOR SEQ ID NO: 268:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq HFFLLNTILLFG/CA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 268:

```

Met Tyr Phe Tyr Gly Leu Thr Phe His Phe Phe Leu Leu Leu Asn Thr
      -20                      -15                      -10

```

Ile Leu Leu Phe Gly Cys Ala Arg
-5 1

(2) INFORMATION FOR SEQ ID NO: 269:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq LWASQGS LQDAQS/ER

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 269:

Met Arg Trp Asn Leu Phe Phe Phe Cys Ile Leu Arg Asn Gln Thr Lys
-25 -20 -15
Leu Trp Ala Ser Gln Gly Ser Leu Gln Asp Ala Gln Ser Glu Arg Gly
-10 -5 1
Cys Phe Ser Leu Asn Gln Asp Gly
5 10

(2) INFORMATION FOR SEQ ID NO: 270:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq FIAALFTMAKTWN/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 270:

Met Phe Ile Ala Ala Leu Phe Thr Met Ala Lys Thr Trp Asn Gln Pro
 -10 -5 1

Gly Cys Ser Ser Met Met Gly Trp Ile Lys Lys Met Arg His Met Thr
 5 10 15

(2) INFORMATION FOR SEQ ID NO: 271:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 61 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -44..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq LWVXLPXXXVIAS/VV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 271:

Met Pro Gly Xaa Lys His Phe Leu Arg Val Phe Arg Xaa Ser Ala Xaa
 -40 -35 -30

Arg Ser Val Gly Tyr Xaa Xaa Lys Pro Gly Thr Ser Arg Ala Ser Leu
 -25 -20 -15

Trp Val Xaa Leu Pro Xaa Xaa Xaa Val Ile Ala Ser Val Val Thr Phe
 -10 -5 1

Ser Xaa His Met Thr Leu Gly Phe Asp Leu Thr Ala Ala
 5 10 15

(2) INFORMATION FOR SEQ ID NO: 272:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -49..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq VALGPLFVTGHFA/GE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 272:

```

Met Arg Leu Glu Ser Pro Asp Glu Asn Phe Ala Val Val Gln Glu His
      -45                      -40                      -35

Ala Ile His His Ile Asp Gly Pro Leu Arg Arg Phe Leu Leu Leu Glu
      -30                      -25                      -20

Val His Glu Pro Val Ala Leu Gly Pro Leu Phe Val Thr Gly His Phe
      -15                      -10                      -5

Ala Gly Glu Asp Val Ala Glu Arg Arg Glu Asp Val Val Gln Arg Leu
      1                      5                      10                      15

Val Val Asp Gly Leu Ala Gln Val Leu Asp Glu Asp Val Ala His Pro
      20                      25                      30

```

(2) INFORMATION FOR SEQ ID NO: 273:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4
seq EVLLPTVLRGSYC/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 273:

```

Met Ala Gly Ser Pro Asp Arg Glu Val Leu Leu Pro Thr Val Leu Arg
-20                      -15                      -10                      -5

Gly Ser Tyr Cys Phe Ser His His Gly
      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 274:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -52..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4
seq RHLFLFEISLVFS/KE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 274:

Met His Val Ser Met Leu Glu Gly Phe Asp Glu Asn Leu Asp Val Gln
-50 -45 -40
Gly Glu Leu Ile Leu Gln Asp Ala Phe Gln Val Trp Asp Pro Lys Ser
-35 -30 -25
Leu Ile Arg Lys Gly Arg Glu Arg His Leu Phe Leu Phe Glu Ile Ser
-20 -15 -10 -5
Leu Val Phe Ser Lys Glu Ile Lys Asp Ser Ser Glu His Asn Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO: 275:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 65 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -39..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9
seq VLAIGLLHIVLLS/IP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 275:

Met Asn Val Gly Thr Ala His Ser Glu Val Asn Xaa Asn Thr Arg Val
-35 -30 -25

Met Lys Xaa Arg Gly Ile Trp Leu Ser Tyr Val Leu Ala Ile Gly Leu
 -20 -15 -10
Leu His Ile Val Leu Leu Ser Ile Pro Phe Val Ser Val Xaa Val Val
 -5 1 5
Trp Thr Leu Thr Asn Leu Ile His Asn Met Gly Met Tyr Ile Phe Leu
10 15 20 25
His

(2) INFORMATION FOR SEQ ID NO: 276:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (D) DEVELOPMENTAL STAGE: Fetal
 - (F) TISSUE TYPE: brain
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: -16..-1
 - (C) IDENTIFICATION METHOD: Von Heijne matrix
 - (D) OTHER INFORMATION: score 3.8
seq FVXAIXXYIPTNS/VQ
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 276:

Met Leu Ser Phe Val Xaa Ala Ile Xaa Xaa Tyr Ile Pro Thr Asn Ser
-15 -10 -5
Val Gln Glu Phe Leu
1 5

(2) INFORMATION FOR SEQ ID NO: 277:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 40 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide

(B) LOCATION: -32..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.8
seq TFINITLWLGSIC/QR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 277:

Met Asp Glu Tyr Ser Trp Trp Cys His Val Leu Glu Val Val Lys Gly
-30 -25 -20

Gln Met Phe Thr Phe Ile Asn Ile Thr Leu Trp Leu Gly Ser Leu Cys
-15 -10 -5

Gln Arg Phe Phe Tyr Ala Ser Gly
1 5

(2) INFORMATION FOR SEQ ID NO: 278:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -23..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.8
seq FIFLIQIWKTCLS/FS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 278:

Met Arg Arg Lys Gly Gln Gly His Leu Ala Phe Ile Phe Leu Ile Gln
-20 -15 -10

Ile Trp Lys Thr Cys Leu Ser Phe Ser Pro Thr Ser Gly
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 279:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

- (D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -25...-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.8
seq NILFLAVSSFSMP/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 279:

Met Phe Leu Ile Ser Gly His Val His Leu Ile Tyr Asn Ile Leu Phe
-25 -20 -15 -10

Leu Ala Val Ser Ser Phe Ser Met Pro Leu Pro Cys Leu Tyr Arg
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 280:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -64...-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.8
seq LFIVVCVICVTLN/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 280:

Met Thr Pro Arg Ile Leu Ser Glu Val Gln Phe Ser Ala Phe Cys Pro
-60 -55 -50

Tyr Trp Thr Ile Ala Arg Ile Leu Glu Arg Val Gly Ser Ala Cys Phe
-45 -40 -35

Arg Leu Glu Leu Cys Ala Ala Ile Val Gly Tyr Phe Val Leu Asp Val
-30 -25 -20

Arg Thr Phe Leu Phe Ile Val Val Cys Val Ile Cys Val Thr Leu Asn
-15 -10 -5

Phe Pro Arg Xaa Xaa Phe Leu Cys Leu Ser Ser Leu Thr Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 281:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 67 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7
seq CSLLSGWGQLLRC/VQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 281:

Met Cys Ser Leu Leu Ser Gly Trp Gly Gln Leu Leu Arg Cys Val Gln
 -10 -5 1

Thr Pro Ala Glu Pro Arg Asp Val Asn Lys Lys Xaa Glu Lys Lys Glu
 5 10 15

Lys Tyr Met Pro Leu Val Asp Ser Leu Cys Gly Gly Leu Gly Thr Arg
 20 25 30

Asn Ser Asp Cys Leu Arg Gly Gly Ala Gly Arg Gly Arg Asp Gly Arg
 35 40 45 50

Arg Ile Arg

(2) INFORMATION FOR SEQ ID NO: 282:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -37..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7
seq LIPFNFSASGLCA/CS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 282:

Met Leu Phe Ser Phe Cys Phe Pro Val His Phe Trp Asn Pro Ser Ser
-35 -30 -25

Leu Phe Pro Pro Ser Ser Val Ser Leu Ile Pro Phe Asn Phe Ser Ala
-20 -15 -10

Ser Gly Leu Cys Ala Cys Ser Arg Thr Phe Thr His Met Gly
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 283:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (D) DEVELOPMENTAL STAGE: Fetal
 - (F) TISSUE TYPE: brain
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: -15..-1
 - (C) IDENTIFICATION METHOD: Von Heijne matrix
 - (D) OTHER INFORMATION: score 3.6
seq WILRILFVIGSXL/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 283:

Met Thr Trp Ile Leu Arg Ile Leu Phe Val Ile Gly Ser Xaa Leu Glu
-15 -10 -5 1

Lys Leu Trp Asn Ile Leu Val Ser Tyr Ile Phe
5 10

(2) INFORMATION FOR SEQ ID NO: 284:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 56 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: -48..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.6
seq ILCIFLGLLIIRC/FK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 284:

Met Ser Ser Thr Tyr Cys Gly Asn Ser Ser Ala Lys Met Ser Val Asn
-45 -40 -35

Glu Val Ser Ala Phe Ser Leu Ser Leu Glu Gln Lys Thr Gly Phe Ala
-30 -25 -20

Phe Val Gly Ile Leu Cys Ile Phe Leu Gly Leu Leu Ile Ile Arg Cys
-15 -10 -5

Phe Lys Ile Leu Leu Xaa Gln Ser
1 5

(2) INFORMATION FOR SEQ ID NO: 285:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq LTMLSMIVGATCY/AM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 285:

Met Thr Asp Ile Trp Leu Thr Met Leu Ser Met Ile Val Gly Ala Thr
-15 -10 -5

Cys Tyr Ala Met Ile Gly
1

(2) INFORMATION FOR SEQ ID NO: 286:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -17..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.6
seq WIYAFISLGYILG/SG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 286:

Met Xaa Xaa Cys Trp Ile Tyr Ala Phe Ile Ser Leu Gly Tyr Ile Leu
-15 -10 -5
Gly Ser Gly Ile Val Gly Leu Phe Gly Asn Phe Met Phe Lys Leu Leu
1 5 10 15
Arg Asn Cys Gln Thr Val Phe Gln Asp Gly Tyr Ala Ile Leu Pro Phe
20 25 30
Pro Pro Thr Gly
35

(2) INFORMATION FOR SEQ ID NO: 287:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -31..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 3.5
seq QLALSWPPXCRV/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 287:

Met Phe Ile Arg Thr Leu Lys Thr Thr Val Leu Pro Phe Met Arg Thr
-30 -25 -20
Ala Pro Gln Leu Ala Leu Ser Trp Val Pro Pro Xaa Cys Arg Val Ser
-15 -10 -5 1
Pro Trp Asp Ser Pro Leu Lys Leu Tyr Cys Leu Gln Pro Gln
5 10 15

(2) INFORMATION FOR SEQ ID NO: 288:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -24..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq SVLIFCLLPYIYH/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 288:

Met Arg Thr Gly Ala Glu Met Arg Thr Asn Ser Ser Val Leu Ile Phe
-20 -15 -10

Cys Leu Leu Pro Tyr Ile Tyr His Phe Phe Pro Ala Ser
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 289:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq EGLELGFSHRTFA/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 289:

Met Ile Val Ile Pro Ser Trp Leu Glu Asn Glu Gly Leu Glu Leu Gly
-20 -15 -10

Phe Ser His Arg Thr Phe Ala Phe Pro Val Thr His Ala Ser Ser Gln
-5 1 5

Tyr Ile Trp Met Asn Xaa Leu Thr Arg Thr Thr Val Ala Ile Ser Val
10 15 20 25

Tyr Phe Trp Thr His Thr Gly
30

(2) INFORMATION FOR SEQ ID NO: 290:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq VISVFLSFLPSYP/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 290:

Met Leu Lys Lys Glu Ile Ala His His Ser Pro Ser Leu Val Ser Cys
-40 -35 -30

Pro Val Cys Thr Thr Lys Tyr Arg Thr Leu Arg Leu Leu Arg Val Ile
-25 -20 -15

Ser Val Phe Leu Ser Phe Leu Pro Ser Tyr Pro Gly Phe Ser Met Gln
-10 -5 1 5

(2) INFORMATION FOR SEQ ID NO: 291:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.5
seq CAYSLPGVALTIG/RQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 291:

Met Thr Xaa Pro Ser Arg Ala Gln Thr Val Asp Xaa Gly Ile Ala Lys
-30 -25 -20 -15
His Cys Ala Tyr Ser Leu Pro Gly Val Ala Leu Thr Leu Gly Arg Gln
-10 -5 1

(2) INFORMATION FOR SEQ ID NO: 292:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.8
seq LGLLCALLPQHGG/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 292:

Met Pro Phe Arg Leu Leu Ile Pro Leu Gly Leu Leu Cys Ala Leu Leu
-20 -15 -10
Pro Gln His His Gly Ala Pro Gly Pro Asp Xaa
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 293:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -18..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 9.2
 seq VFLCSLLAPMVLA/SA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 293:

Met Xaa Leu Val Leu Val Phe Leu Cys Ser Leu Leu Ala Pro Met Val
 -15 -10 -5

Leu Ala Ser Ala Ala Glu Lys Glu Xaa Xaa Met Xaa Pro Phe His Tyr
 1 5 10

Asp Tyr Gln Thr Leu
 15

(2) INFORMATION FOR SEQ ID NO: 294:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 89 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -30..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 9
 seq FFLLLLFRGCLIG/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 294:

Met Ala Leu Arg Arg Pro Pro Arg Leu Arg Leu Cys Ala Arg Leu Pro
 -30 -25 -20 -15

Asp Phe Phe Leu Leu Leu Leu Phe Arg Gly Cys Leu Ile Gly Ala Val
 -10 -5 1

Asn Leu Lys Ser Ser Asn Arg Thr Pro Val Val Gln Glu Phe Glu Ser
 5 10 15

Val Glu Leu Ser Cys Ile Ile Thr Asp Ser Gln Thr Ser Asp Pro Arg
 20 25 30

Ile Glu Trp Lys Lys Ile Gln Asp Glu Gln Thr Thr Tyr Val Phe Phe
 35 40 45 50

Asp Asn Lys Ile Gln Gly Asp Leu Ala
 55

(2) INFORMATION FOR SEQ ID NO: 295:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -60..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.2
seq VLLTLLLI AFIL/II

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 295:

```

Met Gly Gly Asn Gly Ser Thr Cys Lys Pro Asp Thr Glu Arg Gln Gly
-60                               -55                               -50                               -45

Thr Leu Ser Thr Ala Ala Pro Thr Thr Ser Pro Ala Pro Cys Leu Ser
                               -40                               -35                               -30

Asn His His Asn Lys Lys His Leu Ile Leu Ala Phe Cys Ala Gly Val
                               -25                               -20                               -15

Leu Leu Thr Leu Leu Leu Ile Ala Phe Ile Phe Leu Ile Ile Xaa Ser
                               -10                               -5                               1

Tyr Arg Lys Tyr His Ser Lys Pro Gln Ala Pro Asp Pro His Ser Asp
  5                               10                               15                               20

Pro Pro Ala Xaa Leu Ser Xaa Ile Pro Gly Glu Ile Thr Tyr Leu Cys
                               25                               30                               35

Gln His Asn Phe Gln Thr Leu Xaa Xaa Lys Arg Ala Ile Thr Trp Leu
  40                               45                               50

Arg

```

(2) INFORMATION FOR SEQ ID NO: 296:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -39..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.2
seq SALAKLLLTCCSA/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 296:

Met Gly Thr Ala Asp Ser Asp Glu Met Ala Pro Glu Ala Pro Gln His
 -35 -30 -25

Thr His Ile Asp Val His Ile His Gln Glu Ser Ala Leu Ala Lys Leu
 -20 -15 -10

Leu Leu Thr Cys Cys Ser Ala Leu Arg Pro Arg Ala Thr Gln Ala Arg
 -5 1 5

Gly Ser Ser Arg Leu Leu Val Ala Ser Trp Val Met Gln Ile Val Leu
 10 15 20 25

Gly Ile Xaa Ser Ala Val Xaa Gly Gly Phe Phe Tyr Ile Arg Asp Xaa
 30 35 40

Thr Leu Xaa Val Thr Ser Gly Ala Ala Ile Trp Thr Gly Ala Val Ala
 45 50 55

Val

(2) INFORMATION FOR SEQ ID NO: 297:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.6
seq SLLSFLFARVNLG/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 297:

Met Ser Leu Leu Ser Phe Leu Phe Ala Arg Val Asn Leu Gly Ser Pro
 -10 -5 1

Leu Ser Ala Asn Gly
 5

(2) INFORMATION FOR SEQ ID NO: 298:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2
seq WWCCPARLTLTSG/WP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 298:

```

Met Ala Arg Ser Pro Leu Arg Arg Arg Gly Arg Pro Thr Trp Ser Leu
      -40              -35              -30

Ser Thr Pro Arg Pro Gly Ser Pro Thr Ser Ser Ser Arg Ser Trp Trp
      -25              -20              -15

Cys Cys Pro Ala Arg Leu Thr Leu Thr Ser Gly Trp Pro Ala Thr Pro
      -10              -5              1              5

Arg Arg Phe Ser Thr Thr Ser Thr Cys Ser Ile Ala Pro Ser Arg Ser
      10              15              20

Ser Ala Gln Glu Arg Arg Val Arg Arg Trp Pro Cys Ala Thr His Ser
      25              30              35

```

(2) INFORMATION FOR SEQ ID NO: 299:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7

seq TLLLACHLQLEVG/VV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 299:

Met Lys Ile Ile Thr Thr Thr Leu Leu Leu Ala Cys His Leu Gln Leu
 -15 -10 -5
Glu Val Gly Val Val Val Gly Gly Glu Val Asp Met Ala Thr Leu Gln
 1 5 10
Ile Thr Thr Ala Ser
 15

(2) INFORMATION FOR SEQ ID NO: 300:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -37..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7
seq FLGVLALLGYLAV/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 300:

Met Leu Met Pro Val Val Gly Arg Gly Asn Gly Ile Pro Gln Thr Val
 -35 -30 -25
Ser Glu Trp Leu Arg Leu Leu Pro Phe Leu Gly Val Leu Ala Leu Leu
 -20 -15 -10
Gly Tyr Leu Ala Val Arg Pro Gly
 -5 1

(2) INFORMATION FOR SEQ ID NO: 301:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -49..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq PPFFLCLQCFTRG/FE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 301:

```

Met Asp Arg Leu Gly Ser Phe Ser Asn Asp Pro Ser Asp Lys Pro Pro
      -45                      -40                      -35

Cys Arg Gly Cys Ser Ser Tyr Leu Met Glu Pro Tyr Ile Lys Cys Ala
      -30                      -25                      -20

Glu Cys Gly Pro Pro Pro Phe Phe Leu Cys Leu Gln Cys Phe Thr Arg
      -15                      -10                      -5

Gly Phe Glu Tyr Lys Lys His Gln Ser Asp His Thr Tyr Glu Ile Met
      1                      5                      10                      15

Ala Gly Cys Ser Gln Ser Asn Val His Gln Asp Gln Gly Gly Gln
      20                      25                      30

```

(2) INFORMATION FOR SEQ ID NO: 302:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 111 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -87..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6
seq GLLLLYMVYLTIV/EP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 302:

```

Met Ser Asp Val Asn Val Ser Ala Leu Pro Ile Lys Lys Asn Ser Gly
      -85                      -80                      -75

His Ile Tyr Asn Lys Asn Ile Ser Gln Lys Asp Cys Asp Cys Leu His
      -70                      -65                      -60

Val Val Glu Pro Met Pro Val Arg Gly Pro Asp Val Glu Ala Tyr Cys
      -55                      -50                      -45                      -40

```

Leu Arg Cys Glu Cys Lys Tyr Glu Glu Arg Ser Ser Val Thr Ile Lys
 -35 -30 -25
 Val Thr Ile Ile Ile Tyr Leu Ser Ile Leu Gly Leu Leu Leu Tyr
 -20 -15 -10
 Met Val Tyr Leu Thr Leu Val Glu Pro Ile Leu Xaa Arg Arg Leu Phe
 -5 1 5
 Gly His Ala Gln Leu Ile Gln Ser Asp Asp Xaa Ile Gly Gly Leu
 10 15 20

(2) INFORMATION FOR SEQ ID NO: 303:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -46..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq ALSLSLSMAPPNP/GP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 303:

Met Ala Xaa Cys Arg Arg Cys Arg Ser Gln Arg Arg Ser His Cys Cys
 -45 -40 -35
 Gln Asp Arg Arg Leu Arg Arg Pro Arg Leu Thr Leu Trp Arg His His
 -30 -25 -20 -15
 Thr Ala Leu Ser Leu Ser Leu Ser Met Ala Pro Pro Asn Pro Gly Pro
 -10 -5 1

(2) INFORMATION FOR SEQ ID NO: 304:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -36..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5
seq LALTALSVXRKXS/XX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 304:

Met Thr Arg Leu Gly Gly Lys Gly Gly Gln Gln Phe Pro Pro Gly Gln
-35 -30 -25

Lys Ile Ile Ser Lys Asp Ile Leu Ala Leu Thr Ala Leu Ser Val Xaa
-20 -15 -10 -5

Arg Lys Xaa Ser Xaa Xaa Xaa Xaa Xaa Thr Ser Lys Glu Thr Xaa
1 5 10

Asp Asn Gln Asp Ser Val Lys Glu Asn Arg Glu Lys Asp Leu Leu Asp
15 20 25

Ile Ile Lys Gly Thr Lys Val Glu Leu Ser Thr Val Asn Val Gln Thr
30 35 40

Thr Lys Pro Pro Asn Arg Ser Ser Leu Lys Ser Tyr Asn Trp Arg Ala
45 50 55 60

(2) INFORMATION FOR SEQ ID NO: 305:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 12.4
seq ALLLGALLGTAWA/RR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 305:

Met Lys Gly Trp Gly Trp Leu Ala Leu Leu Leu Gly Ala Leu Leu Gly
-20 -15 -10 -5

Thr Ala Trp Ala Arg Arg Ser Arg
1

(2) INFORMATION FOR SEQ ID NO: 306:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 11.4
seq LLCLLLLFGGGDP/RR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 306:

Met His Arg Leu Leu Cys Leu Leu Leu Leu Phe Gly Gly Gly Asp Pro
-15 -10 -5

Arg Arg Arg Ala Glu Ile Arg Leu Gln Ala Thr Ile Cys Ser Arg Pro
1 5 10 15

Leu Arg Lys Thr Thr Ser Gly Arg Gly Gly Pro Pro Trp
20 25

(2) INFORMATION FOR SEQ ID NO: 307:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 128 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 11.1
seq QLLALFFFLPFCLC/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 307:

Met Leu Trp Arg Gln Leu Ile Tyr Trp Gln Leu Leu Ala Leu Phe Phe
-20 -15 -10

Leu Pro Phe Cys Leu Cys Gln Asp Glu Tyr Met Glu Ser Pro Gln Thr
 -5 1 5 10
 Gly Gly Leu Pro Pro Asp Cys Ser Lys Cys Cys His Gly Asp Tyr Ser
 15 20 25
 Phe Arg Gly Tyr Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Ile
 30 35 40
 Pro Gly Asn His Gly Asn Asn Gly Asn Asn Gly Ala Thr Gly His Glu
 45 50 55
 Gly Ala Lys Gly Glu Lys Gly Asp Lys Gly Asp Leu Gly Pro Arg Gly
 60 65 70
 Glu Arg Gly Gln His Gly Pro Lys Gly Glu Lys Gly Tyr Pro Gly Ile
 75 80 85 90
 Pro Pro Glu Leu Gln Ile Ala Phe Met Ala Ser Leu Xaa Pro Thr Ser
 95 100 105

(2) INFORMATION FOR SEQ ID NO: 308:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -17..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.2
seq LLLLVAASAMVRS/XA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 308:

Met Arg Leu Leu Leu Leu Leu Val Ala Ala Ser Ala Met Val Arg
 -15 -10 -5
 Ser Xaa Ala Ser Ala Asn Leu Gly Gly Val Pro Ser Lys Arg Leu Lys
 1 5 10 15
 Met Gln Tyr Thr Thr
 20

(2) INFORMATION FOR SEQ ID NO: 309:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 amino acids

(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -25..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 9
seq ALLVLLGVAAALC/VR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 309:

Met Ser Ser Gly Xaa Glu Leu Leu Trp Pro Gly Ala Ala Leu Leu Val
-25 -20 -15 -10
Leu Leu Gly Val Ala Ala Ser Leu Cys Val Arg Cys Ser Arg Pro Gly
-5 1 5
Ala Lys Arg Ser Glu Lys Ile Tyr Gln Gln Arg Ser Leu Arg Glu Asp
10 15 20
Gln Gln Ser Phe Thr Gly Ser Arg Thr Tyr Ser Leu Val Gly Gln Ala
25 30 35
Trp Pro Gly Pro Leu Ala Asp Met Ala Pro Thr Arg Lys Asp Lys Leu
40 45 50 55
Leu Gln Phe Xaa Pro Ser Leu Glu Xaa Pro Ser Ile Phe Gln Xaa Xaa
60 65 70
Glu Xaa Gln Pro Val Cys Val Cys Ala Ala His Ala Gln Val Gln Xaa
75 80 85
Xaa Gln Arg Lys Ser Thr Ser Arg Glu Val Cys Val Arg Thr Asn Arg
90 95 100
Ala Leu Arg Gly
105

(2) INFORMATION FOR SEQ ID NO: 310:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 125 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -78..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.8
seq SCLGLTLMPFASS/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 310:

```

Met Thr Lys Glu Ile Phe Phe Phe Thr Val Glu Leu Val Cys Glu Asn
      -75                      -70                      -65

Lys Glu Leu Cys Ser Ser Pro Arg Trp Arg Asn Ala Ile Gln Lys Ser
      -60                      -55                      -50

Asn Phe Ser Lys Val Thr Ser Phe Phe Met Ser Cys His His Phe Lys
      -45                      -40                      -35

Gly Leu Ala Pro Leu Pro His Val Tyr Thr Gln Gly Asn Cys Arg Pro
-30                      -25                      -20                      -15

Ile Ser Cys Leu Gly Leu Thr Leu Met Pro Phe Ala Ser Ser Phe Pro
      -10                      -5                      1

Glu Val Lys Val Pro Val Met Tyr Ser His Arg Asn Ile Phe Gln Leu
      5                      10                      15

Phe Met Ser Phe Thr Thr Lys Lys Lys Ile Gln Ser Gly Trp Ser Thr
      20                      25                      30

Thr Leu Ser Ile Phe Leu Val Arg Asn Phe Leu Leu Ile
      35                      40                      45

```

(2) INFORMATION FOR SEQ ID NO: 311:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5
seq YFRALCLPRGAWG/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 311:

```

Met Thr Thr Asp Ile Gly Cys Leu Tyr Phe Arg Ala Leu Cys Leu Pro

```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 313:

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -27...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8
seq VFVCSSVLGQSWG/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 315:

Met Tyr Thr Phe Arg Lys Leu Ser Pro Tyr Leu Asn Lys Ile Val Phe
-25 -20 -15

Val Cys Ser Ser Val Leu Gly Gln Ser Trp Gly Gly Phe Phe Ser Asn
-10 -5 1 5

Leu Ser Glu Thr Leu Ser Ala Thr Leu Phe Asn Gly
10 15

(2) INFORMATION FOR SEQ ID NO: 316:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -21...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4
seq TFCLIFGLGAVWG/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 316:

Met Glu Ser Arg Val Leu Leu Arg Thr Phe Cys Leu Ile Phe Gly Leu
-20 -15 -10

Gly Ala Val Trp Gly Leu Val
-5 1

(2) INFORMATION FOR SEQ ID NO: 317:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 62 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -37..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.4
seq ILIFLGFFLGFLFH/QF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 317:

Met Leu Val Leu Lys Lys His Ser Val Asn Ile Ala Ala Gln Thr Cys
-35 -30 -25
Phe Lys Phe Asn Phe Ile Phe Arg Ile Leu Ile Phe Leu Gly Phe Phe
-20 -15 -10
Leu Gly Leu Phe His Gln Phe Leu Phe Leu Phe Leu Phe Ala Gly Asn
-5 1 5 10
Leu Ser Ser Tyr Leu Leu Lys Gln Ser Lys Ile Gln Ala Arg
15 20 25

(2) INFORMATION FOR SEQ ID NO: 318:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -32..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 6.3
seq TVVLCVGCSTVLC/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 318:

Met Asp Val Lys Cys Pro Gly Cys Tyr Lys Ile Thr Thr Val Phe Ser
-30 -25 -20
His Ala Gln Thr Val Val Leu Cys Val Gly Cys Ser Thr Val Leu Cys
-15 -10 -5
Gln Pro Thr Gly Gly Lys Ala Arg Leu Thr Glu Gly Cys Ser Phe Arg

1

5

10

15

Arg Lys

(2) INFORMATION FOR SEQ ID NO: 319:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2
seq ILSVLHALPAGIA/WS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 319:

Met Cys Ile Ile Leu Ser Val Leu His Ala Leu Pro Ala Gly Ile Ala
-15 -10 -5

Trp Ser Arg Glu Lys Gly
1 5

(2) INFORMATION FOR SEQ ID NO: 320:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -60..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9
seq TWLLLGALPASE/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 320:

Met Leu Val Val Glu Ala Ser Ser Ser Val Arg Leu Ala Ser Ser Glu
 -60 -55 -50 -45

Val Thr Ser Trp Ser Ile Leu Val Thr Pro Ser Ala Ser Thr Pro Ile
 -40 -35 -30

Ile Ser Leu Ser Ala Gly Pro Leu Arg Thr Pro Ser His Ser Lys Thr
 -25 -20 -15

Trp Leu Leu Leu Gly Ala Leu Glu Pro Ala Ser Glu Arg Pro Cys Ser
 -10 -5 1

Ser Val Leu Arg Ser Arg
 5 10

(2) INFORMATION FOR SEQ ID NO: 321:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -27..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7
seq LSLQLIAFPTVSC/EI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 321:

Met Tyr Ser Phe Pro Thr Thr Val Val Glu Glu Ile Leu Ser Leu Ser
 -25 -20 -15

Leu Gln Leu Ile Ala Phe Pro Thr Val Ser Cys Glu Ile Leu Leu Glu
 -10 -5 1 5

Ile Thr Arg

(2) INFORMATION FOR SEQ ID NO: 322:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -16..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.6
 seq LLPLRSLALVRE/SR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 322:

Met Leu Met Leu Leu Pro Leu Arg Ser Leu Leu Ala Leu Val Arg Glu
 -15 -10 -5

Ser Arg Ala Arg
 1

(2) INFORMATION FOR SEQ ID NO: 323:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 103 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Cerebellum

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -25..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 5.5
 seq AQLFACLLRLGTQ/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 323:

Met Val Pro Leu Val Ala Val Val Ser Gly Pro Arg Ala Gln Leu Phe
 -25 -20 -15 -10

Ala Cys Leu Leu Arg Leu Gly Thr Gln Gln Val Gly Pro Leu Gln Leu
 -5 1 5

His Thr Gly Ala Ser His Ala Ala Arg Asn His Tyr Glu Val Leu Val
 10 15 20

Leu Gly Gly Gly Ser Gly Gly Ile Thr Met Ala Ala Arg Met Lys Arg
 25 30 35

Lys Val Gly Ala Glu Asn Val Ala Ile Val Glu Pro Ser Glu Arg His
 40 45 50 55

Phe Tyr Gln Pro Ile Trp Thr Leu Val Gly Ala Gly Ala Xaa Asn Cys

60

65

70

Pro His Leu Val Val Pro Arg
75

(2) INFORMATION FOR SEQ ID NO: 324:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq AFVIACVLSLIST/IY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 324:

Met Asp Asn Arg Phe Ala Thr Ala Phe Val Ile Ala Cys Val Leu Ser
-20 -15 -10 -5

Leu Ile Ser Thr Ile Tyr Met Ala Arg
1 5

(2) INFORMATION FOR SEQ ID NO: 325:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -42..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq WTLTSLDGHLL/EP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 325:

Met Pro Glu Tyr Cys Gly Asn Glu Val Thr Pro Thr Glu Ala Ala Gln
 -40 -35 -30

Ala Pro Glu Val Thr Tyr Glu Ala Glu Gly Ser Leu Trp Thr Leu
 -25 -20 -15

Leu Leu Thr Ser Leu Asp Gly His Leu Leu Glu Pro Asp Ala Glu Tyr
 -10 -5 1 5

Leu His Trp Leu Leu Thr Asn Ile Pro Gly Asn Arg Val Ala Glu Gly
 10 15 20

Gln Val Thr Cys Pro Tyr Leu Pro Pro Phe Pro Ala Arg Gly Ser Gly
 25 30 35

Ile His Arg Leu Ala Phe Leu Leu Phe Lys Gln Asp Gln Pro Ile Asp
 40 45 50

Phe Ser Glu Asp Ala Arg Pro Ser Pro Cys Tyr Gln Leu Xaa Gln Arg
 55 60 65 70

Thr Phe Arg Thr Phe Asp Phe Tyr
 75

(2) INFORMATION FOR SEQ ID NO: 326:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrénals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4
seq RVLCPAAGAVRA/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 326:

Met Asn Arg Val Leu Cys Ala Pro Ala Ala Gly Ala Val Arg Ala Leu
 -15 -10 -5 1

Arg Leu Ile Gly Trp Ala Ser Arg Ser Leu His Pro Leu Pro Gly Lys
 5 10 15

(2) INFORMATION FOR SEQ ID NO: 327:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids

(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -23..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.3
seq MLALLLTAALIFF/AI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 327:

Met Ala Phe Thr Phe Ala Ala Phe Cys Tyr Met Leu Ala Leu Leu Leu
-20 -15 -10

Thr Ala Ala Leu Ile Phe Phe Ala Ile Trp His Ile Ile Ala Phe Asp
-5 1 5

Glu Leu Lys Thr Asp Tyr Lys Asn Pro Ile Asp Gln Leu
10 15 20

(2) INFORMATION FOR SEQ ID NO: 328:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -17..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 5.3
seq XEXLLAFHHDCEA/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 328:

Met Xaa Xaa Xaa Xaa Glu Xaa Leu Leu Ala Phe His His Asp Cys Glu
-15 -10 -5

Ala Ser Pro Ala Thr Trp Asn Leu Ser Pro Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO: 329:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -36..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2
seq PLRLNLLILIEG/SV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 329:

Met Gly Pro Tyr Asn Val Ala Val Pro Ser Asp Val Ser His Ala Arg
-35 -30 -25

Phe Tyr Phe Leu Phe His Arg Pro Leu Arg Leu Leu Asn Leu Leu Ile
-20 -15 -10 -5

Leu Ile Glu Gly Ser Val Val Phe Tyr Gln Leu Tyr Ser Leu Leu Arg
1 5 10

Ser Glu Lys Trp Asn His Thr Leu Ser Met Ala Leu Ile Leu Phe Cys
15 20 25

Asn Tyr Tyr Val Leu Phe Lys Leu Leu Arg Asp Gln
30 35 40

(2) INFORMATION FOR SEQ ID NO: 330:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2

seq IGVGLYLLASAAA/FY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 330:

Met Met Asn Phe Arg Gln Arg Met Gly Trp Ile Gly Val Gly Leu Tyr
 -20 -15 -10

Leu Leu Ala Ser Ala Ala Ala Phe Tyr Tyr Val Phe Glu Ile Ser Glu
 -5 1 5

Thr Tyr Asn Arg Leu Ala Leu Glu His Ile Gln Gln His Xaa Gly
 10 15 20

(2) INFORMATION FOR SEQ ID NO: 331:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 130 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -118..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1
seq AFXVVCWLGPCEA/MH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 331:

Met Leu Phe Ala Ser Gly Gly Phe Xaa Val Lys Leu Tyr Asp Ile Glu
 -115 -110 -105

Gln Gln Gln Ile Arg Asn Ala Leu Glu Asn Ile Arg Lys Glu Met Lys
 -100 -95 -90

Leu Leu Glu Gln Ala Gly Ser Leu Lys Gly Ser Leu Ser Val Glu Glu
 -85 -80 -75

Gln Leu Ser Leu Ile Ser Gly Cys Pro Asn Ile Gln Glu Ala Val Glu
 -70 -65 -60 -55

Gly Ala Met His Ile Gln Glu Cys Val Pro Glu Asp Leu Glu Leu Lys
 -50 -45 -40

Lys Lys Ile Phe Ala Gln Leu Asp Ser Ile Ile Asp Glu Ser Ser Asp
 -35 -30 -25

Leu Lys Arg Phe Xaa Phe Leu Ser His Ala Phe Xaa Val Val Cys Trp
 -20 -15 -10

Leu Gly Pro Cys Glu Ala Met His Arg Gly Ser Ser Cys Glu Ser Ala

-5

1

5

10

Ile Leu

(2) INFORMATION FOR SEQ ID NO: 332:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -22..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.1
seq LCSLPLSPSAVCP/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 332:

Met Gln Cys Phe Leu Gly Gly Leu Gly Leu Cys Ser Leu Pro Leu Ser
-20 -15 -10

Pro Ser Ala Val Cys Pro Ala Pro Thr Ser Ala Pro Trp Trp Glu Gly
-5 1 5 10

Ala Leu

(2) INFORMATION FOR SEQ ID NO: 333:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 63 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -13..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.9
seq MSSFLLSFSQSLS/NV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 333:

```

Met Ser Ser Phe Leu Leu Ser Phe Ser Gln Ser Leu Ser Asn Val Pro
      -10                      -5                      1

Ser Ala Leu Gln Xaa Pro Gln Ile Thr Phe Phe Gln His Pro Leu Ser
      5                      10                      15

Ser Val Met Pro Val Trp Thr Cys Ser Val Val Pro Cys Asp Lys Thr
      20                      25                      30                      35

Xaa Gln Tyr Ser Tyr Cys Phe Tyr Cys Val Leu Gly Thr Val Lys
      40                      45                      50

```

(2) INFORMATION FOR SEO ID NO: 334:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -13..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.9
seq MLTASLAFQLVDG/VS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 334:

Met Leu Thr Ala Ser Leu Ala Phe Gln Leu Val Asp Gly Val Ser Trp
-10 -5 1

Asn Phe Ser Val Ser Lys Met Leu Ala Ser Pro Ser Thr Ser Gly Gln
5 10 15

Leu Ser Gln Phe Gly Ala Ser Leu Tyr Gly Gln Gln Ser Ala Leu Gly
20 25 30 35

Leu Pro Met Arg Gly Met Ser Asn Asn Thr Pro Gln Leu Asn Arg Ser
40 45 50

Leu Ser Gln Xaa Leu Ser Tyr Arg Ala Thr Ser Arg
55 60

(2) INFORMATION FOR SEQ ID NO: 335:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig peptide

(B) LOCATION: -25..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.9
seq LSAFNFLVCLSLG/RG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 335:

Met Tyr Xaa Arg Arg Glu Leu Ser Ile Leu Cys Ile Leu Ser Ala Phe
-25 -20 -15 -10

Asn Phe Leu Val Cys Leu Ser Leu Gly Arg Gly
 -5 1

(2) INFORMATION FOR SEQ ID NO: 336:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

.(A) NAME/KEY: sig peptide

(B) LOCATION: -22..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8
seq IVFGVSWVMLVYS/AS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 336:

Met Gly Leu Ser Ala Met Asp Thr Ser Ile Val Phe Gly Val Ser Trp
-20 -15 -10

Val Met Leu Val Tyr Ser Ala Ser Phe Arg Arg Cys Xaa
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 337:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 69 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -65..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8
seq IILFSAIVGFIYG/YV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 337:

Met Tyr Phe Trp Arg Asp Val Ala Val Ser Leu Asp Thr Leu Trp Ala
-65 -60 -55 -50

Leu Pro Arg Gln Gln Pro Gly Leu Gly Asn Asn Arg Val Leu Gly Leu
-45 -40 -35

Leu Ser Gly Thr Asn Lys Asp Tyr Lys Gly Gln Lys Leu Ala Glu Gln
-30 -25 -20

Met Phe Gln Gly Ile Ile Leu Phe Ser Ala Ile Val Gly Phe Ile Tyr
-15 -10 -5

Gly Tyr Val Ala Ala
1

(2) INFORMATION FOR SEQ ID NO: 338:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -26..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.7
seq LLICXLXIGTATP/VR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 338:

Met His Trp Gly Lys Arg Trp Xaa Leu Xaa Xaa Gly Gly Leu Leu Ile
-25 -20 -15

Cys Xaa Leu Xaa Ile Gly Thr Ala Thr Pro Val Arg Xaa Pro Asn Gly
-10 -5 1 5

Arg Gln Val Leu Val Pro Xaa Gly Tyr Pro Arg Pro Gly Leu Gly Ala
 10 15 20

Val Gly Cys Gly Glu Ala
 25

(2) INFORMATION FOR SEQ ID NO: 339:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: -29..-1
 - (C) IDENTIFICATION METHOD: Von Heijne matrix
 - (D) OTHER INFORMATION: score 4.6
seq ALGLXTCLSVLFG/YA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 339:

Met Ala Xaa Arg Tyr Asn Arg Leu Thr Val Leu Ala Gly Ala Xaa Leu
 -25 -20 -15

Ala Leu Gly Leu Xaa Thr Cys Leu Ser Val Leu Phe Gly Tyr Ala Pro
 -10 -5 1

Gln Ser Ser Pro
 5

(2) INFORMATION FOR SEQ ID NO: 340:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 amino acids
 - (B) TYPE: AMINO ACID
 - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo Sapiens
 - (F) TISSUE TYPE: Substantia nigra
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: -29..-1

- (C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.5
seq FMTCILCRPPISS/CV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 340:

Met Gly Phe Thr Gly Phe Phe Thr Ala Thr Cys Phe Ile Ser Lys Val
 -25 -20 -15
Phe Met Thr Cys Ile Leu Cys Arg Pro Pro Ile Ser Ser Cys Val Leu
 -10 -5 1
Glu Cys Gly
 5

(2) INFORMATION FOR SEQ ID NO: 341:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(D) DEVELOPMENTAL STAGE: Fetal
(F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -35..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.4
seq LIVLLPVLFFSLK/NF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 341:

Met Ile Met Tyr Leu Phe Val Ile Cys Val Ile Phe Glu Ile Ile Arg
-35 -30 -25 -20
Asn Tyr Ala Phe Ser Ile Leu Ile Val Leu Leu Pro Val Leu Phe Phe
 -15 -10 -5
Ser Leu Lys Asn Phe Ile Leu Ser Thr Gln
 1 5

(2) INFORMATION FOR SEQ ID NO: 342:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -37..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.3
seq LWEKLTLLSPGIA/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 342:

Met Ser Thr Val Gly Leu Xaa His Phe Pro Xaa Pro Leu Thr Arg Ile
-35 -30 -25
Cys Pro Ala Pro Trp Gly Leu Arg Leu Trp Glu Lys Leu Thr Leu Leu
-20 -15 -10
Ser Pro Gly Ile Ala Val Thr Pro Val Gln Met Ala Gly Lys Lys Asp
-5 1 5 10
Xaa Pro Ala Leu Leu Ser Leu Asp Glu Asn
15 20

(2) INFORMATION FOR SEQ ID NO: 343:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
(B) TYPE: AMINO ACID
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
(B) LOCATION: -13..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.3
seq MLALAXHLSTVES/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 343:

Met Leu Ala Leu Ala Xaa His Leu Ser Thr Val Glu Ser Glu Lys Gln
-10 -5 1
Lys Leu Arg Ala Gln Val Arg Arg Leu Cys Gln Glu Asn Gln Trp Leu
5 10 15
Arg Asp Glu Leu Ala Gly Thr Gln Gln Arg Leu Gln Arg Ser Glu Gln
20 25 30 35

Ala Val Ala Gln Leu Glu Glu Glu Lys Lys His Leu Glu Phe Leu Gly
40 45 50
Gln Leu Arg Gln Tyr Asp Glu Asp Gly His Thr Ser Glu Ala Gly
55 60 65

(2) INFORMATION FOR SEQ ID NO: 344:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -34..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3
seq QLFAFLNLLPVEA/DI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 344:

Met Leu Leu Ser Ile Gly Met Leu Met Leu Xaa Ala Thr Gln Val Tyr
-30 -25 -20
Thr Ile Leu Thr Val Gln Leu Phe Ala Phe Leu Asn Leu Leu Pro Val
-15 -10 -5
Glu Ala Asp Ile Leu Ala Tyr Asn Phe Glu Asn Ala Ser Gln Thr Phe
1 5 10
Asp Asp Leu Pro Ala
15

(2) INFORMATION FOR SEQ ID NO: 345:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: brain

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -37..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.3
 seq LWEKLTLLSPGIA/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 345:

```

Met Ser Thr Val Gly Leu Phe His Phe Pro Thr Pro Leu Thr Arg Ile
  -35                      -30                      -25

Cys Pro Ala Pro Trp Gly Leu Arg Leu Trp Glu Lys Leu Thr Leu Leu
  -20                      -15                      -10

Ser Pro Gly Ile Ala Val Thr Pro Val Gln Met Ala Gly Lys Lys Asp
  -5                      1                      5                      10

Tyr Pro Ala Leu Leu Ser Leu Asp Glu Xaa Glu Leu Glu Glu Gln Phe
          15                      20                      25

Val Lys Gly His Gly Pro Gly Gly Gln Ala Thr Arg
      30                      35
  
```

(2) INFORMATION FOR SEQ ID NO: 346:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 165 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
 (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: -33..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 4.2
 seq LLNFLGLWSWICK/KW

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 346:

```

Met Glu Leu Thr Ile Phe Ile Leu Arg Leu Ala Ile Tyr Ile Leu Thr
  -30                      -25                      -20

Phe Pro Leu Tyr Leu Leu Asn Phe Leu Gly Leu Trp Ser Trp Ile Cys
  -15                      -10                      -5

Lys Lys Trp Phe Pro Tyr Phe Leu Val Arg Phe Thr Val Ile Tyr Asn
  1                      5                      10                      15

Glu Gln Met Ala Ser Lys Lys Arg Glu Leu Phe Ser Asn Leu Gln Glu
      20                      25                      30
  
```

Phe Ala Gly Pro Ser Gly Lys Leu Ser Leu Leu Glu Val Gly Cys Gly
 35 40 45
 Thr Gly Ala Asn Phe Lys Phe Tyr Pro Pro Gly Cys Arg Val Thr Cys
 50 55 60
 Ile Asp Pro Asn Pro Asn Phe Glu Lys Phe Leu Ile Lys Ser Ile Ala
 65 70 75
 Glu Asn Arg His Leu Gln Phe Glu Arg Phe Val Val Ala Ala Gly Glu
 80 85 90 95
 Asn Met His Gln Val Ala Asp Gly Ser Val Asp Val Val Val Cys Thr
 100 105 110
 Leu Val Leu Cys Ser Val Lys Asn Gln Glu Arg Ile Leu Arg Glu Val
 115 120 125
 Cys Arg Val Leu Arg
 130

(2) INFORMATION FOR SEQ ID NO: 347:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 49 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -37..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1
seq LLLYLCCMINIHH/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 347:

Met Ser Leu Leu His Gly Asn Lys Met Cys Val Thr Ile Arg Pro Thr
 -35 -30 -25
 Gly Gln Pro Leu Asn Gly Asp Leu Leu Leu Leu Tyr Leu Cys Cys Met
 -20 -15 -10
 Ile Asn Ile His His Leu Pro Pro Val Val Leu Pro Arg Thr Pro Gln
 -5 1 5 10
 Gly

(2) INFORMATION FOR SEQ ID NO: 348:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: -17..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 3.9
 seq ISYFIAFPNLSQA/EL
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 348:

Met Ser Phe Asn Ile Ser Tyr Phe Ile Ala Phe Pro Asn Leu Ser Gln
 -15 -10 -5

Ala Glu Leu Thr His Pro Arg Cys Ser Tyr Thr Gly Leu Ser Ser Ser
 1 5 10 15

Cys Gly Phe Gln Leu Ser Asp Thr Pro His Arg
 20 25

(2) INFORMATION FOR SEQ ID NO: 349:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 amino acids
 (B) TYPE: AMINO ACID
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo Sapiens
 (D) DEVELOPMENTAL STAGE: Fetal
 (F) TISSUE TYPE: brain
- (ix) FEATURE:
 (A) NAME/KEY: sig_peptide
 (B) LOCATION: -20..-1
 (C) IDENTIFICATION METHOD: Von Heijne matrix
 (D) OTHER INFORMATION: score 3.7
 seq LTIILLPVHLLIT/IY
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 349:

Met Lys Leu Lys Xaa Asn Val Leu Thr Ile Ile Leu Leu Pro Val His
-20 -15 -10 -5

Leu Leu Ile Thr Ile Tyr Ser Ala Leu Ile

1

5

(2) INFORMATION FOR SEQ ID NO: 350:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 56 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -14..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.7
seq AALVTVLFTGVRR/LH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 350:

Met Ala Ala Leu Val Thr Val Leu Phe Thr Gly Val Arg Arg Leu His
-10 -5 1

Cys Ser Ala Xaa Leu Gly Arg Ala Ala Ser Gly Xaa Tyr Ser Arg Asn
5 10 15

Trp Leu Pro Thr Pro Pro Ala Thr Gly Pro Leu Pro Ser Ser Gln Thr
20 25 30

Gly His Met Arg Met Ala Ala Arg
35 40

(2) INFORMATION FOR SEQ ID NO: 351:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 77 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Surrrenals

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: -42..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.7
seq ILMRDFSPSGIFG/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 351:

```

Met Ala Ser Val Gly Glu Cys Pro Ala Pro Val Pro Val Lys Asp Lys
  -40                -35                -30

Lys Leu Leu Glu Val Lys Leu Gly Glu Leu Pro Ser Trp Ile Leu Met
  -25                -20                -15

Arg Asp Phe Ser Pro Ser Gly Ile Phe Gly Ala Phe Gln Arg Gly Tyr
  -10                -5                1                5

Xaa Arg Tyr Tyr Asn Xaa Tyr Ile Asn Val Xaa Lys Gly Ser Ile Ser
          10                15                20

Gly Ile Xaa Met Val Leu Ala Cys Tyr Val Leu Phe Ser
      25                30                35

```

(2) INFORMATION FOR SEQ ID NO: 352:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Surrrenals

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.9
seq ALLVLVTVALASA/HH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 352:

```

Met Leu Ala Leu Leu Val Leu Val Thr Val Ala Leu Ala Ser Ala His
  -15                -10                -5                1

His Gly Gly Glu His Phe Glu Gly Ala
      5                10

```

(2) INFORMATION FOR SEQ ID NO: 353:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens
(F) TISSUE TYPE: Substantia nigra

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION: -21..-1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 8.9
seq VLLFSGFWGLAMG/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 353:

Met Arg Ile Ile Ser Arg Gln Ile Val Leu Leu Phe Ser Gly Phe Trp
-20 -15 -10

Gly Leu Ala Met Gly Ala Phe Pro Ser Ser Val Gln Ile Gly Gly Leu
-5 1 5 10

Phe Ile Arg Asn Thr Asp Gln Glu Tyr Thr Ala Phe Arg Leu Ala Ile
15 20 25

Phe Leu His Asn Thr Ser Pro Gly
30 35

THIS PAGE BLANK (USPTO)